

QUANTUM FIELD THEORY

P.J. Mulders

Department of Theoretical Physics, Department of Physics and Astronomy,
Faculty of Sciences, VU University,
1081 HV Amsterdam, the Netherlands

E-mail: mulders@few.vu.nl



October 2010 (version 5.21)

Contents

1	Introduction	1
1.1	Quantum field theory	1
1.2	Units	2
1.3	Conventions for vectors and tensors	4
2	Relativistic wave equations	8
2.1	The Klein-Gordon equation	8
2.2	Mode expansion of solutions of the KG equation	9
2.3	Symmetries of the Klein-Gordon equation	10
3	Groups and their representations	13
3.1	The rotation group and $SU(2)$	13
3.2	Representations of symmetry groups	15
3.3	The Lorentz group	18
3.4	The generators of the Poincaré group	20
3.5	Representations of the Poincaré group	21
4	The Dirac equation	27
4.1	The Lorentz group and $SL(2, C)$	27
4.2	Spin 1/2 representations of the Lorentz group	29
4.3	General representations of γ matrices and Dirac spinors	31
4.4	Plane wave solutions	34
4.5	γ gymnastics and applications	36
5	Vector fields and Maxwell equations	39
5.1	Fields for spin 1	39
5.2	The electromagnetic field	39
5.3	The electromagnetic field and topology*	41
6	Classical lagrangian field theory	44
6.1	Euler-Lagrange equations	44
6.2	Lagrangians for spin 0, 1/2 and 1 fields	46
6.3	Symmetries and conserved (Noether) currents	48
6.4	Space-time symmetries	49
6.5	(Abelian) gauge theories	50
7	Quantization of fields	54
7.1	Canonical quantization	54
7.2	Creation and annihilation operators	56
7.3	The real scalar field	57
7.4	The complex scalar field	60

7.5	The Dirac field	60
7.6	The electromagnetic field	62
8	Discrete symmetries	66
8.1	Parity	66
8.2	Charge conjugation	67
8.3	Time reversal	68
8.4	Bi-linear combinations	69
8.5	Form factors	70
9	Path integrals and quantum mechanics	73
9.1	Time evolution as path integral	73
9.2	Functional integrals	75
9.3	Time ordered products of operators and path integrals	78
9.4	An application: time-dependent perturbation theory	79
9.5	The generating functional for time ordered products	81
9.6	Euclidean formulation	82
10	Feynman diagrams for scattering amplitudes	85
10.1	Generating functionals for free scalar fields	85
10.2	Generating functionals for interacting scalar fields	89
10.3	Interactions and the S-matrix	92
10.4	Feynman rules	95
10.5	Some examples	100
11	Scattering theory	103
11.1	kinematics in scattering processes	103
11.2	Crossing symmetry	106
11.3	Cross sections and lifetimes	107
11.4	Unitarity condition	109
11.5	Unstable particles	111
12	The standard model	114
12.1	Non-abelian gauge theories	114
12.2	Spontaneous symmetry breaking	118
12.3	The Higgs mechanism	122
12.4	The standard model $SU(2)_W \otimes U(1)_Y$	123
12.5	Family mixing in the Higgs sector and neutrino masses	128

References

As most directly related books to these notes, I refer to the book of Srednicki [1] and Ryder [2]. Other text books of Quantum Field Theory that are useful are given in refs [3-6]. The notes contain all essential information, but are rather compact. Besides the books, there is also material available via the web pages of colleagues, e.g. those of Jan Smit (UvA), Pierre van Baal (UL), Bert Schellekens (RU and NIKHEF) and Gerard 't Hooft (UU).

The schedule in the Fall of 2010 is Chapters 1 through 10 in period 2 (7 weeks in November and December 2010) and Chapters 11 and 12 in period 3 (3 weeks in January 2011).

1. M. Srednicki, *Quantum Field Theory*, Cambridge University Press, 2007.
2. L.H. Ryder, *Quantum Field Theory*, Cambridge University Press, 1985.
3. M.E. Peskin and D.V. Schroeder, *An introduction to Quantum Field Theory*, Addison-Wesley, 1995.
4. M. Veltman, *Diagrammatica*, Cambridge University Press, 1994.
5. S. Weinberg, *The quantum theory of fields*; Vol. I: Foundations, Cambridge University Press, 1995; Vol. II: Modern Applications, Cambridge University Press, 1996.
6. C. Itzykson and J.-B. Zuber, *Quantum Field Theory*, McGraw-Hill, 1980.

Corresponding chapters in books of Ryder, Peskin & Schroeder and Srednicki.

These notes Section	Ryder	Peskin & Schroeder	Srednicki
1.1			1
1.2			1
1.3	2.1		1
2.1	2.2		1
2.2			3
2.3			
3.1	2.3		
3.2	2.3		
3.3	2.3	3.1	2
3.4	2.3	3.1	2
3.5	2.7		
4.1			
4.2	2.3, 2.4	3.2	
4.3	2.5	3.4	
4.4	2.5	3.3	
4.5		3.4	
5.1	2.8		
5.2	3.4		
6.1	3.1, 3.2	2.2	
6.2	3.2, 3.3		36
6.3	3.2	2.2	22
6.4	3.2		
6.5	3.3		

These notes Section	Ryder	Peskin & Schroeder	Srednicki
7.1	4	2.3	3
7.2	4		3
7.3	4	2.3, 2.4	3
7.4	4	2.3, 2.4	
7.5	4	3.5	37
7.6	4	2.3	
8.1		3.6	
8.2		3.6	
8.3		3.6	
8.4		3.6	
8.5			
9.1	5.1	9.1	6
9.2	5.4, 6.2, 6.7	9.2, 9.5	7
9.3	5.5		6
9.4		4.2	
9.5	5.5		6
9.6			
10.1	6.1, 6.3	4.1, 4.2, 9.2	8
10.2	6.4, 6.5, 6.6	4.3, 4.4	9
10.3	6.8	4.4, 4.6	
10.4	6.7	4.7, 4.8	
10.5		5	
11.1		4.5	
11.2	6.10	4.5	
11.3		7.3	
11.4		7.3	
12.1		20.1	
12.2	8.1, 8.2	20.1	
12.3	8.3	20.1	
12.4	8.5	20.2, 20.3	
12.5			

Corrections 2010/2011)

Chapter 1

Introduction

1.1 Quantum field theory

In *quantum field theory* the theories of quantum mechanics and special relativity are united. In quantum mechanics a special role is played by *Planck's constant* h , usually given divided by 2π ,

$$\begin{aligned}\hbar \equiv h/2\pi &= 1.054\,571\,68\,(18) \times 10^{-34} \text{ J s} \\ &= 6.582\,119\,15\,(56) \times 10^{-22} \text{ MeV s.}\end{aligned}\tag{1.1}$$

In the limit that the action S is much larger than \hbar , $S \gg \hbar$, quantum effects do not play a role anymore and one is in the *classical* domain. In special relativity a special role is played by *the velocity of light* c ,

$$c = 299\,792\,458 \text{ m s}^{-1}.\tag{1.2}$$

In the limit that $v \ll c$ one reaches the *non-relativistic* domain.

In the framework of classical mechanics as well as quantum mechanics the position of a particle is a well-defined concept and the position coordinates can be used as *dynamical variables* in the description of the particles and their interactions. In quantum mechanics, the position can in principle be determined at any time with any accuracy, being eigenvalues of the position operators. One can talk about states $|\mathbf{r}\rangle$ and the wave function $\psi(\mathbf{r}) = \langle \mathbf{r} | \psi \rangle$. In this *coordinate representation* the position operators \mathbf{r}_{op} simply acts as

$$\mathbf{r}_{\text{op}} \psi(\mathbf{r}) = \mathbf{r} \psi(\mathbf{r}).\tag{1.3}$$

The uncertainty principle tells us that in this representation the momenta cannot be fully determined. Corresponding position and momentum operators do not commute. They satisfy the well-known (canonical) operator commutation relations

$$[r_i, p_j] = i\hbar \delta_{ij},\tag{1.4}$$

where δ_{ij} is the Kronecker δ function. Indeed, the action of the momentum operator in the coordinate representation is not as simple as the position operator. It is given by

$$\mathbf{p}_{\text{op}} \psi(\mathbf{r}) = -i\hbar \nabla \psi(\mathbf{r}).\tag{1.5}$$

One can also choose a representation in which the momenta of the particles are the dynamical variables. The corresponding states are $|\mathbf{p}\rangle$ and the wave functions $\tilde{\psi}(\mathbf{p}) = \langle \mathbf{p} | \psi \rangle$ are the Fourier transforms of the coordinate space wave functions,

$$\tilde{\psi}(\mathbf{p}) = \int d^3r \exp\left(-\frac{i}{\hbar} \mathbf{p} \cdot \mathbf{r}\right) \psi(\mathbf{r}),\tag{1.6}$$

and

$$\psi(\mathbf{r}) = \int \frac{d^3p}{(2\pi\hbar)^3} \exp\left(\frac{i}{\hbar}\mathbf{p}\cdot\mathbf{r}\right) \tilde{\psi}(\mathbf{p}). \quad (1.7)$$

The existence of a limiting velocity, however, leads to new fundamental limitations on the possible measurements of physical quantities. Let us consider the measurement of the position of a particle. This position cannot be measured with infinite precision. Any device that wants to locate the position of say a particle within an interval Δx will contain momentum components $p \propto \hbar/\Delta x$. Therefore if we want $\Delta x \leq \hbar/mc$ (where m is the rest mass of the particle), momenta of the order $p \propto mc$ and energies of the order $E \propto mc^2$ are involved. It is then possible to create a particle - antiparticle pair and it is no longer clear of which particle we are measuring the position. As a result, we find that the original particle cannot be located better than within a distance \hbar/mc , its *Compton wavelength*,

$$\Delta x \geq \frac{\hbar}{mc}. \quad (1.8)$$

For a moving particle $mc^2 \rightarrow E$ (or by considering the Lorentz contraction of length) one has $\Delta x \geq \hbar c/E$. If the particle momentum becomes relativistic, one has $E \approx pc$ and $\Delta x \geq \hbar/p$, which says that a particle cannot be located better than its *de Broglie wavelength*.

Thus the coordinates of a particle cannot act as dynamical variables (since these must have a precise meaning).

Some consequences are that only in cases where we restrict ourselves to distances $\gg \hbar/mc$, the concept of a wave function becomes a meaningful (albeit approximate) concept. For a massless particle one gets $\Delta x \gg \hbar/p = \lambda/2\pi$, i.e. the coordinates of a photon only become meaningful in cases where the typical dimensions are much larger than the wavelength.

For the momentum or energy of a particle we know that in a finite time Δt , the energy uncertainty is given by $\Delta E \geq \hbar/\Delta t$. This implies that the momenta of particles can only be measured exactly when one has an infinite time available. For a particle in interaction, the momentum changes with time and a measurement over a long time interval is meaningless. The only case in which the momentum of a particle can be measured exactly is when the particle is free and stable against decay. In this case the momentum is conserved and one can let Δt become infinitely large.

The result thus is that *the only observable quantities that can serve as dynamical coordinates are the momenta (and further the internal degrees of freedom like polarizations, ...)* of free particles. These are the particles in the initial and final state of a scattering process. The theory will not give an observable meaning to the time dependence of interaction processes. The description of such a process as occurring in the course of time is just as unreal as classical paths are in non-relativistic quantum mechanics.

The main problem in Quantum Field Theory is to determine the probability amplitudes between well-defined initial and final states of a system of free particles. The set of such amplitudes $\langle \mathbf{p}'_1, \mathbf{p}'_2; \text{out} | \mathbf{p}_1, \mathbf{p}_2; \text{in} \rangle \equiv \langle \mathbf{p}'_1, \mathbf{p}'_2; \text{in} | S | \mathbf{p}_1, \mathbf{p}_2; \text{in} \rangle$ determines the *scattering matrix* or S-matrix.

Another point that needs to be emphasized is the meaning of particle in the above context. Actually, the better name might be 'degree of freedom'. If the energy is low enough to avoid excitation of internal degrees of freedom, an atom is a perfect example of a particle. In fact, it is the behavior under Poincaré transformations or in the limit $v \ll c$ Gallilei transformations that determine the description of a particle state, in particular the free particle state.

1.2 Units

It is important to choose an appropriate set of units when one considers a specific problem, because physical sizes and magnitudes only acquire a meaning when they are considered in relation to each other. This is true specifically for the domain of atomic, nuclear and high energy physics, where the typical numbers are difficult to conceive on a macroscopic scale. They are governed by a few fundamental units and constants, which have been discussed in the previous section, namely \hbar and

c. By making use of these fundamental constants, we can work with less units. For instance, the quantity *c* is used to define the meter. We could as well have set *c* = 1. This would mean that one of the two units, meter or second, is eliminated, e.g. given a length *l* the quantity *l/c* has the dimension of time and one finds 1 m = 0.33 × 10⁻⁸ s or eliminating the second one would use that, given a time *t*, the quantity *ct* has dimension of length and hence 1 s = 3 × 10⁸ m.

Table 1.1: *Physical quantities and their canonical dimensions d, determining units (energy)^d.*

quantity	quantity with dimension energy ^d	dimension	canonical dimension <i>d</i>
time <i>t</i>	<i>t/ħ</i>	(energy) ⁻¹	-1
length <i>l</i>	<i>l/(ħc)</i>	(energy) ⁻¹	-1
energy <i>E</i>	<i>E</i>	(energy) ¹	1
momentum <i>p</i>	<i>pc</i>	(energy) ¹	1
angular momentum <i>ℓ</i>	<i>ℓ/ħ</i>	(energy) ⁰	0
mass <i>m</i>	<i>mc²</i>	(energy) ¹	1
area <i>A</i>	<i>A/(ħc)²</i>	(energy) ⁻²	-2
force <i>F</i>	<i>F ħc</i>	(energy) ²	2
charge (squared) <i>e²</i>	<i>α = e²/4πϵ₀ ħc</i>	(energy) ⁰	0
Newton's constant <i>G_N</i>	<i>G_N/(ħc⁵)</i>	(energy) ⁻²	-2
velocity <i>v</i>	<i>v/c</i>	(energy) ⁰	0

In field theory, it turns out to be convenient to work with units such that \hbar and *c* are set to one. All length, time and energy or mass units then can be expressed in one unit and powers thereof, for which one can use energy (see table 1.1). The elementary unit that is most relevant depends on the domain of applications, e.g. the eV for atomic physics, the MeV or GeV for nuclear physics and the GeV or TeV for high energy physics. To convert to other units of length or time we use appropriate combinations of \hbar and *c*, e.g. for lengths

$$\hbar c = 0.197\,326\,968\,(17)\,\text{GeV fm} \quad (1.9)$$

or for order of magnitude estimates $\hbar c \approx 0.2\,\text{GeV fm} = 200\,\text{eV nm}$, implying (when $\hbar = c = 1$) that $1\,\text{fm} = 10^{-15}\,\text{m} \approx 5\,\text{GeV}^{-1}$. For areas, e.g. cross sections, one needs

$$\hbar^2 c^2 = 0.389\,379\,323\,(67)\,\text{GeV}^2\,\text{mbarn} \quad (1.10)$$

(1 barn = 10⁻²⁸ m² = 10² fm²). For times one needs

$$\hbar = 6.582\,119\,15\,(56) \times 10^{-22}\,\text{MeV s}, \quad (1.11)$$

implying (when $\hbar = c = 1$) that $1\,\text{s} \approx 1.5 \times 10^{24}\,\text{GeV}^{-1}$. Depending on the specific situation, of course masses come in that one needs to know or look up, e.g. those of the electron or proton,

$$m_e = 9.109\,382\,6\,(16) \times 10^{-31}\,\text{kg} = 0.510\,998\,918\,(44)\,\text{MeV}/c^2, \quad (1.12)$$

$$m_p = 1.672\,621\,71\,(29) \times 10^{-27}\,\text{kg} = 0.938\,272\,029\,(80)\,\text{GeV}/c^2. \quad (1.13)$$

Furthermore one encounters the strength of the various interactions. In some cases like the electromagnetic and strong interactions, these can be written as dimensionless quantities, e.g. for electromagnetism the *fine structure constant*

$$\alpha = \frac{e^2}{4\pi\epsilon_0\hbar c} = 1/137.035\,999\,11\,(46). \quad (1.14)$$

For weak interactions and gravity one has quantities with a dimension, e.g. for gravity Newton's constant,

$$\frac{G_N}{\hbar c^5} = 6.7087(10) \times 10^{-39} \text{ GeV}^{-2}. \quad (1.15)$$

By putting this quantity equal to 1, one can also eliminate the last dimension. All masses, lengths and energies are compared with the Planck mass or length (see exercises). Having many particles, the concept of temperature becomes relevant. A relation with energy is established via the average energy of a particle being of the order of kT , with the Boltzmann constant given by

$$k = 1.3806505(24) \times 10^{-23} \text{ J/K} = 8.617343(15) \times 10^{-5} \text{ eV/K}. \quad (1.16)$$

Quantities that do not contain \hbar or c are classical quantities, e.g. the mass of the electron m_e . Quantities that contain only \hbar are expected to play a role in non-relativistic quantum mechanics, e.g. the Bohr radius, $a_\infty = 4\pi\epsilon_0\hbar^2/m_e e^2$ or the Bohr magneton $\mu_e = e\hbar/2m_e$. Quantities that only contain c occur in classical relativity, e.g. the electron rest energy $m_e c^2$ and the classical electron radius $r_e = e^2/4\pi\epsilon_0 m_e c^2$. Quantities that contain both \hbar and c play a role in relativistic quantum mechanics, e.g. the electron Compton wavelength $\lambda_e = \hbar/m_e c$. It remains useful, however, to use \hbar and c to simplify the calculation of quantities.

1.3 Conventions for vectors and tensors

We start with vectors in Euclidean 3-space $E(3)$. A vector \mathbf{x} can be expanded with respect to a basis $\hat{\mathbf{e}}_i$ ($i = 1, 2, 3$ or $i = x, y, z$),

$$\mathbf{x} = \sum_{i=1}^3 x_i \hat{\mathbf{e}}_i = x_i \hat{\mathbf{e}}_i, \quad (1.17)$$

to get the three components of a vector, x_i . When a repeated index appears, such as on the right hand side of this equation, summation over this index is assumed (Einstein summation convention). Choosing an orthonormal basis, the metric in $E(3)$ is given by $\hat{\mathbf{e}}_i \cdot \hat{\mathbf{e}}_j = \delta_{ij}$, where the Kronecker delta is given by

$$\delta_{ij} = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j, \end{cases} \quad (1.18)$$

The inner product of two vectors is given by

$$\mathbf{x} \cdot \mathbf{y} = x_i y_i \hat{\mathbf{e}}_i \cdot \hat{\mathbf{e}}_j = x_i y_j \delta_{ij} = x_i y_i. \quad (1.19)$$

The inner product of a vector with itself gives its length squared. A vector can be rotated, $\mathbf{x}' = R\mathbf{x}$ or $x'_i = R_{ij}x_j$ leading to a new vector with different components. Actually, *rotations* are those *real, linear* transformations that do not change the length of a vector. Tensors of rank n are objects with n components that transform according to $T'_{i_1 \dots i_n} = R_{i_1 j_1} \dots R_{i_n j_n} T_{j_1 \dots j_n}$. A vector is a tensor of rank 1. The inner product of two vectors is a rank 0 tensor or scalar. The Kronecker delta is a constant rank-2 tensor. It is an invariant tensor that does not change under rotations. The only other invariant constant tensor in $E(3)$ is the Levi-Civita tensor

$$\epsilon_{ijk} = \begin{cases} 1 & \text{if } ijk \text{ is an even permutation of } 123 \\ -1 & \text{if } ijk \text{ is an odd permutation of } 123 \\ 0 & \text{otherwise.} \end{cases} \quad (1.20)$$

that can be used in the cross product of two vectors $\mathbf{z} = \mathbf{x} \times \mathbf{y}$, in which case $z_i = \epsilon_{ijk} x_j y_k$. Useful relations are

$$\epsilon_{ijk} \epsilon_{lmn} = \begin{vmatrix} \delta_{il} & \delta_{im} & \delta_{in} \\ \delta_{jl} & \delta_{jm} & \delta_{jn} \\ \delta_{kl} & \delta_{km} & \delta_{kn} \end{vmatrix}, \quad (1.21)$$

$$\epsilon_{ijk} \epsilon_{imn} = \delta_{jm} \delta_{kn} - \delta_{jn} \delta_{km}, \quad (1.22)$$

$$\epsilon_{ijk} \epsilon_{ijl} = 2 \delta_{kl}. \quad (1.23)$$

We note that for Euclidean spaces (with a positive definite metric) vectors and tensors there is only one type of indices. No difference is made between upper or lower. So we could have used all upper indices in the above equations. When 3-dimensional space is considered as part of Minkowski space, however, we will use upper indices for the three-vectors.

In special relativity we start with a four-dimensional real vector space $E(1,3)$ with basis \hat{n}_μ ($\mu = 0,1,2,3$). Vectors are denoted $x = x^\mu \hat{n}_\mu$. The length (squared) of a vector is obtained from the scalar product,

$$x^2 = x \cdot x = x^\mu x^\nu \hat{n}_\mu \cdot \hat{n}_\nu = x^\mu x^\nu g_{\mu\nu}. \quad (1.24)$$

The quantity $g_{\mu\nu} \equiv \hat{n}_\mu \cdot \hat{n}_\nu$ is the metric tensor, given by $g_{00} = -g_{11} = -g_{22} = -g_{33} = 1$ (the other components are zero). For four-vectors in Minkowski space we will use the notation with upper indices and write $x = (t, \mathbf{x}) = (x^0, x^1, x^2, x^3)$, where the coordinate $t = x^0$ is referred to as the time component, x^i are the three space components. Because of the different signs occurring in $g_{\mu\nu}$, it is convenient to distinguish lower indices from upper indices. The lower indices are constructed in the following way, $x_\mu = g_{\mu\nu} x^\nu$, and are given by $(x_0, x_1, x_2, x_3) = (t, -\mathbf{x})$. One has

$$x^2 = x^\mu x_\mu = t^2 - \mathbf{x}^2. \quad (1.25)$$

The scalar product of two different vectors x and y is denoted

$$x \cdot y = x^\mu y^\nu g_{\mu\nu} = x^\mu y_\mu = x_\mu y^\mu = x^0 y^0 - \mathbf{x} \cdot \mathbf{y}. \quad (1.26)$$

Within Minkowski space the *real, linear* transformations that do not change the length of a four-vector are called the *Lorentz transformations*. These transformations do change the components of a vector, denoted as $V'^\mu = \Lambda^\mu_{\nu} V^\nu$. The (invariant) lengths often have special names, such as *eigentime* τ for the position vector $\tau^2 \equiv x^2 = t^2 - \mathbf{x}^2$. The invariant distance between two points x and y in Minkowski space is determined from the length $ds^\mu = (x - y)^\mu$. The *real, linear* transformations that leave the length of a vector invariant are called (*homogeneous*) *Lorentz transformations*. The transformations that leave invariant the distance $ds^2 = dt^2 - (dx^2 + dy^2 + dz^2)$ between two points are called inhomogeneous Lorentz transformations or *Poincaré transformations*. The Poincaré transformations include Lorentz transformations and translations.

Unlike in Euclidean space, the invariant length or distance (squared) is not positive definite. One can distinguish:

- $ds^2 > 0$ (timelike intervals); in this case an inertial system exists in which the two points are at the same space point and in that frame ds^2 just represents the time difference $ds^2 = dt^2$;
- $ds^2 < 0$ (spacelike intervals); in this case an inertial system exists in which the two points are at the same time and ds^2 just represents minus the spatial distance squared $ds^2 = -d\mathbf{x}^2$;
- $ds^2 = 0$ (lightlike or null intervals); the points lie on the lightcone and they can be connected by a light signal.

Many other four vectors and tensors transforming like $T'^{\mu_1 \dots \mu_n} = \Lambda^{\mu_1}_{\nu_1} \dots \Lambda^{\mu_n}_{\nu_n} T^{\nu_1 \dots \nu_n}$ can be constructed. In Minkowski space, one must distinguish tensors with upper or lower indices and one can have mixed tensors. Relations relating tensor expressions, independent of a coordinate system, are

called covariant. Examples are the scalar products above but also relations like $p^\mu = m dx^\mu/d\tau$ for the momentum four vector. Note that in this equation one has on left- and righthandside a four vector because τ is a scalar quantity! The equation with $t = x^0$ instead of τ simply would not make sense! The momentum four vector, explicitly written as $(p^0, \mathbf{p}) = (E, \mathbf{p})$, is timelike with invariant length (squared) $p^2 = p \cdot p = p^\mu p_\mu = E^2 - \mathbf{p}^2 = m^2$, where m is called the *mass* of the system.

The derivative ∂_μ is defined $\partial_\mu = \partial/\partial x^\mu$ and we have a four vector ∂ with components

$$(\partial_0, \partial_1, \partial_2, \partial_3) = \left(\frac{\partial}{\partial t}, \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right) = \left(\frac{\partial}{\partial t}, \nabla \right). \quad (1.27)$$

It is easy to convince oneself of the nature of the indices in the above equation, because one has

$$\partial_\mu x^\nu = g_\mu^\nu. \quad (1.28)$$

Note that g_μ^ν with one upper and lower index, constructed via the metric tensor itself, $g_\mu^\nu = g_{\mu\rho} g^{\rho\nu}$ and is in essence a 'Kronecker delta', $g_0^0 = g_1^1 = g_2^2 = g_3^3 = 1$. The length squared of ∂ is the d'Alembertian operator, defined by

$$\square = \partial^\mu \partial_\mu = \frac{\partial^2}{\partial t^2} - \nabla^2. \quad (1.29)$$

The value of the antisymmetric tensor $\epsilon^{\mu\nu\rho\sigma}$ is determined in the same way as for ϵ^{ijk} , starting from

$$\epsilon^{0123} = 1. \quad (1.30)$$

(Note that there are different conventions around and sometimes the opposite sign is used). It is an invariant tensor, not affected by Lorentz transformations. The product of two epsilon tensors is given by

$$\epsilon^{\mu\nu\rho\sigma} \epsilon^{\mu'\nu'\rho'\sigma'} = - \begin{vmatrix} g^{\mu\mu'} & g^{\mu\nu'} & g^{\mu\rho'} & g^{\mu\sigma'} \\ g^{\nu\mu'} & g^{\nu\nu'} & g^{\nu\rho'} & g^{\nu\sigma'} \\ g^{\rho\mu'} & g^{\rho\nu'} & g^{\rho\rho'} & g^{\rho\sigma'} \\ g^{\sigma\mu'} & g^{\sigma\nu'} & g^{\sigma\rho'} & g^{\sigma\sigma'} \end{vmatrix}, \quad (1.31)$$

$$\epsilon^{\mu\nu\rho\sigma} \epsilon_\mu^{\nu'\rho'\sigma'} = - \begin{vmatrix} g^{\nu\nu'} & g^{\nu\rho'} & g^{\nu\sigma'} \\ g^{\rho\nu'} & g^{\rho\rho'} & g^{\rho\sigma'} \\ g^{\sigma\nu'} & g^{\sigma\rho'} & g^{\sigma\sigma'} \end{vmatrix}, \quad (1.32)$$

$$\epsilon^{\mu\nu\rho\sigma} \epsilon_{\mu\nu}^{\rho'\sigma'} = -2 \left(g^{\rho\rho'} g^{\sigma\sigma'} - g^{\rho\sigma'} g^{\sigma\rho'} \right), \quad (1.33)$$

$$\epsilon^{\mu\nu\rho\sigma} \epsilon_{\mu\nu\rho}^{\sigma'} = -6 g^{\sigma\sigma'}, \quad (1.34)$$

$$\epsilon^{\mu\nu\rho\sigma} \epsilon_{\mu\nu\rho\sigma} = -24. \quad (1.35)$$

The first identity, for instance, is easily proven for $\epsilon^{0123} \epsilon^{0123}$ from which the general case can be obtained by making permutations of indices on the lefthandside and permutations of rows or columns on the righthandside. Each of these permutations leads to a minus sign, but more important has the same effect on lefthandside and righthandside. For the contraction of a vector with the antisymmetric tensor one often uses the shorthand notation

$$\epsilon^{ABCD} = \epsilon^{\mu\nu\rho\sigma} A_\mu B_\nu C_\rho D_\sigma. \quad (1.36)$$

Exercises

Exercise 1.1

- (a) In the the Hydrogen atom (quantum system) the scale is set by the Bohr radius, $a_\infty = 4\pi\epsilon_0 \hbar^2 / m_e e^2$. Relate this quantity to the electron Compton wavelength λ_e via the dimensionless fine structure constant α .

- (b) Relate the classical radius of the electron (a relativistic concept), $r_e = e^2/4\pi\epsilon_0 m_e c^2$ to the Compton wavelength.
- (c) Calculate the Compton wavelength of the electron and the quantities under (a) and (b) using the value of $\hbar c$, α and $m_e c^2 = 0.511$ MeV. This demonstrates how a careful use of units can save a lot of work. One does not need to know \hbar , c , ϵ_0 , m_e , e , but only appropriate combinations.
- (d) Use the value of the gravitational constant $G_N/\hbar c^5 = 6.71 \times 10^{-39}$ GeV⁻² to construct a mass M_{pl} (Planck mass). Compare it with the proton mass and use Eq. 1.13 to give its actual value in kg. Also construct and calculate the Planck length L_{pl} , which is the Compton wavelength for the Planck mass.
- (e) Note that $\hbar/m = \lambda c$ and use this to calculate in a simple way (avoiding putting in the value of e) the Bohr magneton $\mu_e = e\hbar/2m_e$ and the nuclear magneton $\mu_p = e\hbar/2m_p$ in electronvolt per Tesla (eV/T).
[Note: what is the MKS unit for V/T?]

Exercise 1.2

Prove the identity $\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) = (\mathbf{A} \cdot \mathbf{C}) \mathbf{B} - (\mathbf{A} \cdot \mathbf{B}) \mathbf{C}$ using the properties of the tensor ϵ^{ijk} given in section 1.3.

Exercise 1.3

Prove the following relation

$$\epsilon^{\mu\nu\rho\sigma} g^{\alpha\beta} = \epsilon^{\alpha\nu\rho\sigma} g^{\mu\beta} + \epsilon^{\mu\alpha\rho\sigma} g^{\nu\beta} + \epsilon^{\mu\nu\alpha\sigma} g^{\rho\beta} + \epsilon^{\mu\nu\rho\alpha} g^{\sigma\beta}.$$

by a simple few-line reasoning [For instance: If $\{\mu, \nu, \rho, \sigma\}$ is a permutation of $\{0, 1, 2, 3\}$ the index α can only be equal to *one* of the indices in $\epsilon^{\mu\nu\rho\sigma}$, ...].

Exercise 1.4

Lightcone coordinates for a four vector a (which we will denote with square brackets as $[a^-, a^+, a^1, a^2]$ or $[a^-, a^+, \mathbf{a}_T]$) are defined through

$$a^\pm \equiv (a^0 \pm a^3)/\sqrt{2}.$$

- (a) Express the scalar product $a \cdot b$ in lightcone coordinates and deduce from this the values of g_{++} , g_{--} , g_{+-} and g_{-+} . How are a_+ and a_- related to a^+ and a^- .
- (b) The coordinates (a^0, a^1, a^2, a^3) are the expansion coefficients using the basis vectors $\hat{n}_0, \hat{n}_1, \hat{n}_2, \hat{n}_3$; These are

$$\begin{aligned}\hat{n}_1 &= (0, 1, 0, 0) \\ \hat{n}_2 &= (0, 0, 1, 0) \\ \hat{n}_0 &= (1, 0, 0, 0) \\ \hat{n}_3 &= (0, 0, 0, 1)\end{aligned}$$

Also for the coordinates $[a^-, a^+, a^1, a^2]$ we can find basis vectors $\hat{n}_-, \hat{n}_+, \hat{n}_1, \hat{n}_2$. Note that $a \cdot \hat{n}_\alpha = a_\alpha$ and derive from that the components \hat{n}_α^μ of the four vector \hat{n}_α . Use that to give the components of the basis vectors \hat{n}_+ and \hat{n}_- .

Chapter 2

Relativistic wave equations

2.1 The Klein-Gordon equation

In this chapter, we just want to play a bit with covariant equations and study their behavior under Lorentz transformations. The Schrödinger equation in quantum mechanics is the operator equation corresponding to the non-relativistic expression for the energy,

$$E = \frac{\mathbf{p}^2}{2M}, \quad (2.1)$$

under the substitution (in coordinate representation)

$$E \longrightarrow E_{\text{op}} = i\frac{\partial}{\partial t}, \quad \mathbf{p} \longrightarrow \mathbf{p}_{\text{op}} = -i\nabla. \quad (2.2)$$

Acting on the wave function one finds for a free particle,

$$i\frac{\partial}{\partial t}\psi(\mathbf{r}, t) = -\frac{\nabla^2}{2M}\psi(\mathbf{r}, t). \quad (2.3)$$

Equations 2.1 and 2.3 are not covariant. But the replacement 2.2, written as $p_\mu \longrightarrow i\partial_\mu$ is covariant (the same in every frame of reference). Thus a covariant equation can be obtained by starting with the (covariant) equation for the invariant length of the four vector (E, \mathbf{p}) ,

$$p^2 = p^\mu p_\mu = E^2 - \mathbf{p}^2 = M^2, \quad (2.4)$$

where M is the particle mass. Substitution of operators gives the Klein-Gordon (KG) equation for a real or complex function ϕ ,

$$(\square + M^2)\phi(\mathbf{r}, t) = \left(\frac{\partial^2}{\partial t^2} - \nabla^2 + M^2\right)\phi(\mathbf{r}, t) = 0. \quad (2.5)$$

Although it is straightforward to find the solutions of this equation, namely plane waves characterized by a wave number \mathbf{k} ,

$$\phi_{\mathbf{k}}(\mathbf{r}, t) = \exp(-ik^0 t + i\mathbf{k} \cdot \mathbf{r}), \quad (2.6)$$

with $(k^0)^2 = \mathbf{k}^2 + M^2$, the interpretation of this equation as a single-particle equation in which ϕ is a complex wave function poses problems because the energy spectrum is not bounded from below and the probability is not positive definite.

The energy spectrum is not bounded from below: considering the above stationary plane wave solutions, one obtains

$$k^0 = \pm\sqrt{\mathbf{k}^2 + M^2} = \pm E_{\mathbf{k}}, \quad (2.7)$$

i.e. there are solutions with negative energy.

Probability is not positive: in quantum mechanics one has the probability and probability current

$$\rho = \psi^* \psi \quad (2.8)$$

$$\mathbf{j} = -\frac{i}{2M} (\psi^* \nabla \psi - (\nabla \psi^*) \psi) \equiv -\frac{i}{2M} \psi^* \overleftrightarrow{\nabla} \psi. \quad (2.9)$$

They satisfy the continuity equation,

$$\frac{\partial \rho}{\partial t} = -\nabla \cdot \mathbf{j}, \quad (2.10)$$

which follows directly from the Schrödinger equation. This continuity equation can be written down covariantly using the components (ρ, \mathbf{j}) of the four-current j ,

$$\partial_\mu j^\mu = 0. \quad (2.11)$$

Therefore, relativistically the density is not a scalar quantity, but rather the zero component of a four vector. The appropriate current corresponding to the KG equation (see Exercise 2.2) is

$$j^\mu = i \phi^* \overleftrightarrow{\partial}^\mu \phi \quad \text{or} \quad (\rho, \mathbf{j}) = \left(i \phi^* \overleftrightarrow{\partial}_0 \phi, -i \phi^* \overleftrightarrow{\nabla} \phi \right). \quad (2.12)$$

It is easy to see that this current is conserved if ϕ (and ϕ^*) satisfy the KG equation. The KG equation, however, is a second order equation and ϕ and $\partial\phi/\partial t$ can be fixed arbitrarily at a given time. This leads to the existence of negative densities.

As we will see later both problems are related and have to do with the existence of particles and antiparticles, for which we need the interpretation of ϕ itself as an operator, rather than as a wave function. This operator has all possible solutions in it multiplied with creation (and annihilation) operators. At that point the dependence on position \mathbf{r} and time t is just a dependence on numbers/parameters on which the operator depends, just as the dependence on time was in ordinary quantum mechanics. Then, there are no longer fundamental objections to mix up space and time, which is what Lorentz transformations do. And, it is simply a matter of being careful to find a consistent (covariant) theory.

2.2 Mode expansion of solutions of the KG equation

Before quantizing fields, having the KG equation as a space-time symmetric (classical) equation, we want the most general solution. For this we note that an arbitrary solution for the field ϕ can always be written as a superposition of plane waves,

$$\phi(x) = \int \frac{d^4 k}{(2\pi)^4} 2\pi \delta(k^2 - M^2) e^{-i k \cdot x} \tilde{\phi}(k) \quad (2.13)$$

with (in principle complex) coefficients $\tilde{\phi}(k)$. The integration over k -modes clearly is covariant and restricted to the ‘mass’-shell (as required by Eq. 2.5). It is possible to rewrite it as an integration over positive energies only but this gives two terms (use the result of exercise 2.3),

$$\phi(x) = \int \frac{d^3 k}{(2\pi)^3 2E_k} \left(e^{-i k \cdot x} \tilde{\phi}(E_k, \mathbf{k}) + e^{i k \cdot x} \tilde{\phi}(-E_k, -\mathbf{k}) \right). \quad (2.14)$$

Introducing $\tilde{\phi}(E_k, \mathbf{k}) \equiv a(\mathbf{k})$ and $\tilde{\phi}(-E_k, -\mathbf{k}) \equiv b^*(\mathbf{k})$ one has

$$\phi(x) = \int \frac{d^3 k}{(2\pi)^3 2E_k} \left(e^{-i k \cdot x} a(\mathbf{k}) + e^{i k \cdot x} b^*(\mathbf{k}) \right) = \phi_+(x) + \phi_-(x). \quad (2.15)$$

In Eqs 2.14 and 2.15 one has eliminated k^0 and in both equations $k \cdot x = E_k t - \mathbf{k} \cdot \mathbf{x}$. The coefficients $a(\mathbf{k})$ and $b^*(\mathbf{k})$ are the amplitudes of the two independent solutions (*two*, after restricting the energies to be positive). They are referred to as mode and anti-mode amplitudes (or because of their origin positive and negative energy modes). The choice of a and b^* allows an easier distinction between the cases that ϕ is real ($a = b$) or complex (a and b are independent amplitudes).

2.3 Symmetries of the Klein-Gordon equation

We arrived at the Klein-Gordon equation by constructing a covariant operator $(\partial_\mu \partial^\mu + M^2)$ acting on a complex function ϕ . Performing some Lorentz transformation $x \rightarrow x' = \Lambda x$, one thus must have that the function $\phi \rightarrow \phi'$ such that

$$\phi'(x') = \phi(x) \quad \text{or} \quad \phi'(x) = \phi(\Lambda^{-1}x). \quad (2.16)$$

The consequence of this is discussed in Exercise 2.6

We will explicitly discuss the example of a discrete symmetry, for which we consider space inversion, i.e. changing the sign of the spatial coordinates, which implies

$$(x^\mu) = (t, \mathbf{x}) \rightarrow (t, -\mathbf{x}) \equiv (\tilde{x}^\mu). \quad (2.17)$$

Transforming everywhere in the KG equation $x \rightarrow \tilde{x}$ one obtains

$$\left(\tilde{\partial}_\mu \tilde{\partial}^\mu + M^2 \right) \phi(\tilde{x}) = 0. \quad (2.18)$$

Since $a \cdot b = \tilde{a} \cdot \tilde{b}$, it is easy to see that

$$(\partial_\mu \partial^\mu + M^2) \phi(\tilde{x}) = 0, \quad (2.19)$$

implying that for each solution $\phi(x)$ there exists a corresponding solution with the same energy, $\phi^P(x) \equiv \phi(\tilde{x})$ (P for parity). It is easy to show that

$$\begin{aligned} \phi^P(x) = \phi(\tilde{x}) &= \int \frac{d^3k}{(2\pi)^3 2E_k} \left(e^{-i k \cdot \tilde{x}} a(\mathbf{k}) + e^{i k \cdot \tilde{x}} b^*(\mathbf{k}) \right) \\ &= \int \frac{d^3k}{(2\pi)^3 2E_k} \left(e^{-i \tilde{k} \cdot x} a(\mathbf{k}) + e^{i \tilde{k} \cdot x} b^*(\mathbf{k}) \right) \\ &= \int \frac{d^3k}{(2\pi)^3 2E_k} \left(e^{-i k \cdot x} a(-\mathbf{k}) + e^{i k \cdot x} b^*(-\mathbf{k}) \right), \end{aligned} \quad (2.20)$$

or since one can define

$$\phi^P(x) \equiv \int \frac{d^3k}{(2\pi)^3 2E_k} \left(e^{-i k \cdot x} a^P(\mathbf{k}) + e^{i k \cdot x} b^{P*}(\mathbf{k}) \right), \quad (2.21)$$

one has for the mode amplitudes $a^P(\mathbf{k}) = a(-\mathbf{k})$ and $b^P(\mathbf{k}) = b(-\mathbf{k})$. This shows how parity transforms \mathbf{k} -modes into $-\mathbf{k}$ modes.

Another symmetry is found by complex conjugating the KG equation. It is trivial to see that

$$(\partial_\mu \partial^\mu + M^2) \phi^*(x) = 0, \quad (2.22)$$

showing that with each solution there is a corresponding *charge conjugated* solution $\phi^C(x) = \phi^*(x)$. In terms of modes one has

$$\begin{aligned} \phi^C(x) = \phi^*(x) &= \int \frac{d^3k}{(2\pi)^3 2E_k} \left(e^{-i k \cdot x} b(\mathbf{k}) + e^{i k \cdot x} a^*(\mathbf{k}) \right) \\ &\equiv \int \frac{d^3k}{(2\pi)^3 2E_k} \left(e^{-i k \cdot x} a^C(\mathbf{k}) + e^{i k \cdot x} b^{C*}(\mathbf{k}) \right), \end{aligned} \quad (2.23)$$

i.e. for the mode amplitudes $a^C(\mathbf{k}) = b(\mathbf{k})$ and $b^C(\mathbf{k}) = a(\mathbf{k})$. For the real field one has $a^C(\mathbf{k}) = a(\mathbf{k})$. This shows how charge conjugation transforms 'particle' modes into 'antiparticle' modes and vice versa.

Exercises

Exercise 2.1

Show that for a conserved current ($\partial_\mu j^\mu = 0$) the charge in a finite volume, $Q_V \equiv \int_V d^3x j^0(x)$, satisfies

$$\dot{Q}_V = - \int_S ds \cdot \mathbf{j},$$

and thus for any normalized solution the full 'charge', letting $V \rightarrow \infty$, is conserved, $\dot{Q} = 0$.

Exercise 2.2

Show that if ϕ and ϕ^* are solutions of the KG equation, that

$$j^\mu = i \phi^* \overleftrightarrow{\partial}^\mu \phi$$

is a conserved current

Note: $A \overleftrightarrow{\partial}_\mu B \equiv A \partial_\mu B - (\partial_\mu A) B$.

Exercise 2.3

Show that¹

$$\int \frac{d^4k}{(2\pi)^4} 2\pi \delta(k^2 - M^2) \theta(k^0) F(k^0, \mathbf{k}) = \int \frac{d^3k}{(2\pi)^3 2E_k} F(E_k, \mathbf{k}),$$

where $E_k = \sqrt{\mathbf{k}^2 + M^2}$.

Exercise 2.4

write down the 3-dimensional Fourier transform $\tilde{\phi}(\mathbf{k}, t) \equiv \int d^3x \phi(\mathbf{x}, t) \exp(-i\mathbf{k} \cdot \mathbf{x})$ and its time derivative $i\partial_0 \tilde{\phi}(\mathbf{k}, t)$ for the mode expansion in Eq. 2.15. Use these to show that

$$\begin{aligned} a(\mathbf{k}) &= e^{iE_k t} i \overleftrightarrow{\partial}_0 \tilde{\phi}(\mathbf{k}, t) = e^{iE_k t} (i\partial_0 + E_k) \tilde{\phi}(\mathbf{k}, t), \\ b(\mathbf{k}) &= e^{iE_k t} i \overleftrightarrow{\partial}_0 \tilde{\phi}^*(-\mathbf{k}, t), \end{aligned}$$

Note that $a(\mathbf{k})$ and $b(\mathbf{k})$ are independent of t .

¹Use the following property of delta functions

$$\delta(f(x)) = \sum_{\text{zeros } x_n} \frac{1}{|f'(x_n)|} \delta(x - x_n).$$

Exercise 2.5

- (a) As an introduction to parts (b) and (c), show that for two fields ϕ_1 (with coefficients a_1 and b_1^*) and ϕ_2 (with coefficients a_2 and b_2^*) one has

$$\int d^3x \phi_1^*(x) \phi_2(x) = \int \frac{d^3k}{(2\pi)^3 4E_k^2} \left(a_1^*(\mathbf{k})a_2(\mathbf{k}) + b_1(\mathbf{k})b_2^*(\mathbf{k}) + a_1^*(\mathbf{k})b_2^*(-\mathbf{k}) e^{2iE_k t} + b_1(\mathbf{k})a_2(-\mathbf{k}) e^{-2iE_k t} \right).$$

- (b) Express now the full charge Q_V (exercise 2.1) for a complex scalar field current (exercise 2.2) in terms of the $a(\mathbf{k})$ and $b(\mathbf{k})$ using the expansion in Eq. 2.15.

After you have found the result, note that replacing $a^*(\mathbf{k}) \rightarrow a_{\text{op}}^\dagger(\mathbf{k})$ and $a(\mathbf{k}) \rightarrow a_{\text{op}}(\mathbf{k})$ with a_{op} and a_{op}^\dagger being annihilation and creation operators (as known from a harmonic oscillator) and the same for the coefficients $b(\mathbf{k})$, one sees that Q is expressed as an infinite sum over number operators $a^\dagger a$ (particles) and $b^\dagger b$ (anti-particles). This will be the topic of chapter 7.

- (c) Similarly, also using the result in (a), express the quantities

$$E = \int d^3x \left((\partial_0 \phi)^* (\partial_0 \phi) + \nabla \phi^* \cdot \nabla \phi + M^2 \phi^* \phi \right),$$

$$P^i = \int d^3x (\partial^{\{0} \phi)^* (\partial^{i\}} \phi),$$

in terms of the $a(\mathbf{k})$ and $b(\mathbf{k})$. Note that $a^{\{\mu b^{\nu\}}$ indicates symmetrization, $a^{\{\mu b^{\nu\}} \equiv a^\mu b^\nu + a^\nu b^\mu$. We will encounter these quantities later as energy and momentum.

Exercise 2.6

Write down the mode expansion for the Lorentz transformed scalar field $\phi'(x)$ and show that it implies that the Lorentz transformed modes satisfy $a'(\Lambda \mathbf{k}) = a(\mathbf{k})$.

Exercise 2.7

To solve the problem with positive and negative energies and get a positive definite density, an attempt to construct a first order differential equation for the time evolution would be to write,

$$i \frac{\partial}{\partial t} \psi(\mathbf{r}, t) = (-i \boldsymbol{\alpha} \cdot \nabla + m \beta) \psi(\mathbf{r}, t),$$

where ψ is a wave function and the nature of $\boldsymbol{\alpha}$ and β are left open for the moment.

- (a) Show that relativistic invariance, i.e. making sure that each component of ψ satisfies the Klein-Gordon equation (in terms of differential operators $\partial_\mu \partial^\mu + m^2 = 0$) requires the anticommutation relation²

$$\{\alpha^i, \alpha^j\} = 2 \delta^{ij} I, \quad \{\alpha^i, \beta\} = 0, \quad \text{and} \quad \beta^2 = I.$$

From this one concludes that $\boldsymbol{\alpha}$ and β must be matrix-valued and ψ must be a multi-component wave function.

- (b) Show that one has current conservation for the current

$$j^0 = \psi^\dagger \psi \quad \text{and} \quad \mathbf{j} = \psi^\dagger \boldsymbol{\alpha} \psi.$$

²We denote the commutator as $[A, B] = AB - BA$ and the anticommutator as $\{A, B\} = AB + BA$.

Chapter 3

Groups and their representations

The simple systems that we want to describe in a relativistically invariant way are free particles with spin, e.g. electrons. In this section we will investigate the requirements imposed by Poincaré invariance. In particular, we want to investigate if there exist objects other than just a scalar (real or complex) field ϕ , e.g. two-component fields in analogy to the two-component wave functions used to include spin in a quantum mechanical description of an electron in the atom.

Since the KG equation expresses just the relativistic relation between energy and momentum, we also want it to hold for particles with spin. In quantum mechanics spin is described by a vector, i.e. it has 3 components that we know the behavior of under rotations. In a relativistic theory, however, the symmetry group describing rotations is embedded in the Lorentz group, and we must study the representations of the Lorentz group. Particles with spin then will be described by certain spinors. The KG equation will actually remain valid, in particular each component of these spinors will satisfy this equation.

Before proceeding with the Lorentz group we will first discuss the rotation group as an example of a Lie group with and a group we are familiar with in ordinary quantum mechanics.

3.1 The rotation group and $SU(2)$

The rotation groups $SO(3)$ and $SU(2)$ are examples of *Lie groups*, that is groups characterized by a finite number of real parameters, in which the parameter space forms locally a Euclidean space. A general rotation — we will consider $SO(3)$ as an example — is of the form


$$\begin{pmatrix} V'_x \\ V'_y \\ V'_z \end{pmatrix} = \begin{pmatrix} \cos \varphi & \sin \varphi & 0 \\ -\sin \varphi & \cos \varphi & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} V_x \\ V_y \\ V_z \end{pmatrix} \quad (3.1)$$

for a rotation around the z -axis or shorthand $\mathbf{V}' = R(\varphi, \hat{z})\mathbf{V}$. The parameter-space of $SO(3)$ is a sphere with radius π . Any rotation can be uniquely written as $R(\varphi, \hat{\mathbf{n}})$ where $\hat{\mathbf{n}}$ is a unit vector and φ is the rotation angle, $0 \leq \varphi \leq \pi$, provided we identify the antipodes, i.e. $R(\pi, \hat{\mathbf{n}}) \equiv R(\pi, -\hat{\mathbf{n}})$. Locally this parameter-space is 3-dimensional and correspondingly one has three generators. For an infinitesimal rotation around the z -axis one has

$$R(\delta\varphi, \hat{z}) = 1 + i \delta\varphi L_z \quad (3.2)$$

with as generator

$$L_z = \frac{1}{i} \left. \frac{\partial R(\varphi, \hat{z})}{\partial \varphi} \right|_{\varphi=0} = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (3.3)$$


 Figure 3.1: the parameter spaces of $SO(3)$ (left) and $SU(2)$ (right).

In the same way we can consider rotations around the x- and y-axes that are generated by

$$L_x = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad L_y = \begin{pmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ -i & 0 & 0 \end{pmatrix}, \quad (3.4)$$

or $(L_k)_{ij} = -i \epsilon_{ijk}$. It is straightforward to check that any (finite) rotation can be obtained from a combination of infinitesimal rotations, for rotations around z for instance,

$$R(\varphi, \hat{z}) = \lim_{N \rightarrow \infty} \left[R\left(\frac{\varphi}{N}, \hat{z}\right) \right]^N. \quad (3.5)$$

Rotations in general do not commute, which reflects itself in the noncommutation of the generators. They satisfy the commutation relations

$$[L_i, L_j] = i \epsilon_{ijk} L_k. \quad (3.6)$$

Summarizing, the rotations in $SO(3)$ can be generated from infinitesimal rotations that can be expressed in terms of a basis of three generators L_x , L_y and L_z . These generators form a *three-dimensional Lie algebra* $SO(3)$. With matrix commutation this algebra satisfies the requirements for a Lie algebra, namely that there exists a bilinear product $[\]$ that satisfies

- $\forall x, y \in \underline{A} \Rightarrow [x, y] \in \underline{A}$.
- $[x, x] = 0$ (thus $[x, y] = -[y, x]$).
- $[x, [y, z]] + [y, [z, x]] + [z, [x, y]] = 0$ (Jacobi identity).

Next, we turn to the group $SU(2)$ of *special* ($\det A = 1$) *unitary* ($A^\dagger = A^{-1}$) 2×2 matrices. These matrices can be defined as acting on 2-component spinors ($\chi \rightarrow A\chi$) or equivalently as acting on 2×2 matrices ($B \rightarrow ABA^\dagger$). It is straightforward to check that the conditions require

$$\begin{aligned} A &= \begin{pmatrix} a_0 + i a_3 & +i a_1 + a_2 \\ +i a_1 - a_2 & a_0 - i a_3 \end{pmatrix} = a_0 \mathbf{1} + i \mathbf{a} \cdot \boldsymbol{\sigma} \\ &= a_0 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + i a_1 \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + i a_2 \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} + i a_3 \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \end{aligned} \quad (3.7)$$

with real a 's and $\sum_{i=0}^3 (a_i)^2 = 1$. One way of viewing the parameter space, thus is as the surface of a sphere in 4 Euclidean dimensions. Locally this is a 3-dimensional Euclidean space and $SU(2)$, therefore, is a 3-dimensional Lie-group. Writing $a_0 = \cos(\varphi/2)$ and $\mathbf{a} = \hat{\mathbf{n}} \sin(\varphi/2)$ we have¹

$$A = A(\varphi, \hat{\mathbf{n}}) = \mathbf{1} \cos\left(\frac{\varphi}{2}\right) + i (\boldsymbol{\sigma} \cdot \hat{\mathbf{n}}) \sin\left(\frac{\varphi}{2}\right) = \exp\left(i \frac{\varphi}{2} \boldsymbol{\sigma} \cdot \hat{\mathbf{n}}\right), \quad (3.8)$$

¹Note that $(\boldsymbol{\sigma} \cdot \mathbf{a})(\boldsymbol{\sigma} \cdot \mathbf{b}) = \mathbf{a} \cdot \mathbf{b} + i \boldsymbol{\sigma} \cdot (\mathbf{a} \times \mathbf{b})$, and thus $(\boldsymbol{\sigma} \cdot \hat{\mathbf{n}})^2 = 1$.

where we have used the (for operators new!) definition

$$e^A \equiv \sum_{n=0}^{\infty} \frac{A^n}{n!} = 1 + A + \frac{1}{2!} A^2 + \dots \quad (3.9)$$

The parameter-space of $SU(2)$, thus, also can be considered as a *filled* 3-sphere, now with radius 2π and with *all* points at the surface identified (see figure). The infinitesimal generators of $SU(2)$ are obtained by considering infinitesimal transformations, i.e. for fixed $\hat{\mathbf{n}}$,

$$A(\varphi, \hat{\mathbf{n}}) \approx \mathbf{1} + i \varphi \mathbf{J} \cdot \hat{\mathbf{n}}, \quad (3.10)$$

with

$$\mathbf{J} \cdot \hat{\mathbf{n}} \equiv \frac{1}{i} \left. \frac{\partial A(\varphi, \hat{\mathbf{n}})}{\partial \varphi} \right|_{\varphi=0} = \frac{\boldsymbol{\sigma}}{2} \cdot \hat{\mathbf{n}}. \quad (3.11)$$

Thus $\sigma_x/2$, $\sigma_y/2$ and $\sigma_z/2$ form the basis of the Lie-algebra $SU(2)$. They satisfy

$$\left[\frac{\sigma_i}{2}, \frac{\sigma_j}{2} \right] = i \epsilon_{ijk} \frac{\sigma_k}{2}. \quad (3.12)$$

One, thus, immediately sees that the Lie algebras are identical, $SU(2) \simeq SO(3)$, i.e. one has a Lie algebra isomorphism that is linear and preserves the bilinear product.

There exists a corresponding mapping of the groups given by

$$\begin{aligned} \mu : \quad SU(2) &\longrightarrow SO(3) \\ A(\varphi, \hat{\mathbf{n}}) &\longrightarrow R(\varphi, \hat{\mathbf{n}}) & 0 \leq \varphi \leq \pi \\ &\longrightarrow R(2\pi - \varphi, \hat{\mathbf{n}}) & \pi \leq \varphi \leq 2\pi. \end{aligned}$$

The relation

$$A(\boldsymbol{\sigma} \cdot \mathbf{a}) A^{-1} = \boldsymbol{\sigma} \cdot R_A \mathbf{a} \quad (3.13)$$

(valid for any vector \mathbf{a}) can be used to establish the homomorphism (Check that it satisfies the requirements of a homomorphism, in particular that $AB \rightarrow R_{AB} = R_A R_B$). Near the identity, the above mapping corresponds to the trivial mapping of the Lie algebras. In the full parameter space, however, the $SU(2) \rightarrow SO(3)$ mapping is a 2 : 1 mapping where both $A = \pm \mathbf{1}$ are mapped into $R = I$.

3.2 Representations of symmetry groups

The presence of symmetries simplifies the description of a physical system and is at the heart of physics. Suppose we have a system described by a Hamiltonian H . The existence of symmetries means that there are operators g belonging to a symmetry group G that commute with the Hamiltonian,

$$[g, H] = 0 \quad \text{for } g \in G. \quad (3.14)$$

For a Lie group, it is sufficient that the generators commute with H , since any finite transformation can be constructed from the infinitesimal ones, sometimes in more than one way (but this will be discussed later), i.e.

$$[\underline{g}, H] = 0 \quad \text{for } \underline{g} \in \underline{G}. \quad (3.15)$$

The next issue is to find the appropriate symmetry operators in the Hilbert space. For instance in a function space translations,

$$\phi'(\mathbf{r}) = \phi(\mathbf{r} + \mathbf{a}) = \phi(\mathbf{r}) + \mathbf{a} \cdot \nabla \phi + \frac{1}{2!} (\mathbf{a} \cdot \nabla)^2 \phi + \dots,$$

are generated by the derivative acting on the functions, more precise $\phi(\mathbf{r} + \mathbf{a}) = U(\mathbf{a})\phi(\mathbf{r})$ where

$$U(\mathbf{a}) = \exp(+i\mathbf{a}\mathbf{p}_{\text{op}}), \quad (3.16)$$

with $\mathbf{p}_{\text{op}} = -i\nabla$. Similarly one has the time translation operator $\phi(t + \tau) = U(\tau)\phi(t)$,

$$U(\tau) = \exp(-i\tau H), \quad (3.17)$$

with $H = i\partial/\partial t$ and the rotation operator, $\phi(r, \theta, \varphi + \alpha) = U(\alpha, \hat{z})\phi(r, \theta, \varphi)$,

$$U(\alpha, \hat{z}) = \exp(+i\alpha \ell_z), \quad (3.18)$$

with $\ell_z = -i\partial/\partial\varphi = -i(x\partial/\partial y - y\partial/\partial x)$ or $\ell_{\text{op}} = \mathbf{r}_{\text{op}} \times \mathbf{p}_{\text{op}}$. The above set of operators H , \mathbf{p}_{op} and ℓ_{op} work in Hilbert space and their commutators indeed follow the required commutation relations as symmetry operators. In particular the quantum operators ℓ_{op} obey in quantum mechanics commutation relations $[\ell_i, \ell_j] = i\epsilon_{ijk}\ell_k$ that are identical to those in Eq. 3.6. It may seem trivial, but it is at the heart of being able to construct a suitable Hilbert space, e.g. for the function space, just starting off with the (basic) canonical commutation relations $[r_i, p_j] = i\delta_{ij}$. The latter is sufficient to get the commutation relations of the (quantum) rotation operators.

Note that the same requirements would apply to classical mechanics. Somehow the nontrivial structure of rotations should show up and it, indeed, does in the nontrivial behavior of Poisson brackets of quantities $A(x, p)$ and $B(x, p)$ (we will come back to this also in Chapter 7). Indeed, the Poisson bracket operation satisfies all properties of a Lie algebra.

In many cases one can simply construct a suitable Hilbert space or part of it by considering the representations Φ of a group G . These are mappings of G into a finite dimensional vector space, preserving the group structure. The finite dimensional vector space just represents new degrees of freedom. In order to find local representations Φ of a Lie-group G , it is sufficient to consider the representations $\underline{\Phi}$ of the Lie-algebra \underline{G} . These are mappings from \underline{G} into the same finite dimensional vector space (its dimension is the dimension of the representation), which preserve the Lie-algebra structure, i.e. the commutation relations. The most well-known example is spin (or the total angular momentum of a system) in non-relativistic quantum mechanics as representations of $SU(2)$.

To get explicit representations, one looks among the generators for a maximal commuting set of operators, for rotations the operator J_z and the (quadratic Casimir) operator \mathbf{J}^2 . Casimir operators commute with all the generators and the eigenvalue of \mathbf{J}^2 can be used to label the representation (j). Within the $(2j + 1)$ -dimensional representation space $V^{(j)}$ one can label the eigenstates $|j, m\rangle$ with eigenvalues of J_z . The other generators J_x and J_y (or $J_{\pm} \equiv J_x \pm iJ_y$) then transform between the states in $V^{(j)}$. From the algebra one derives $\mathbf{J}^2|j, m\rangle = j(j + 1)|j, m\rangle$, $J_z|j, m\rangle = m|j, m\rangle$, while $J_{\pm}|j, m\rangle = \sqrt{j(j + 1) - m(m \pm 1)}|j, m \pm 1\rangle$ with $2j + 1$ being integer and $m = j, j - 1, \dots, -j$.

Explicit representations using the basis states $|j, m\rangle$ with m -values running from the highest to the lowest, $m = j, j - 1, \dots, -j$ one has for $j = 0$:

$$J_z = \begin{pmatrix} 0 \end{pmatrix}, \quad J_+ = \begin{pmatrix} 0 \end{pmatrix}, \quad J_- = \begin{pmatrix} 0 \end{pmatrix},$$

for $j = 1/2$:

$$J_z = \begin{pmatrix} 1/2 & 0 \\ 0 & -1/2 \end{pmatrix}, \quad J_+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad J_- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix},$$

for $j = 1$:

$$J_z = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad J_+ = \begin{pmatrix} 0 & \sqrt{2} & 0 \\ 0 & 0 & \sqrt{2} \\ 0 & 0 & 0 \end{pmatrix}, \quad J_- = \begin{pmatrix} 0 & 0 & 0 \\ \sqrt{2} & 0 & 0 \\ 0 & \sqrt{2} & 0 \end{pmatrix},$$

and for $j = 3/2$:

$$J_z = \begin{pmatrix} 3/2 & 0 & 0 & 0 \\ 0 & 1/2 & 0 & 0 \\ 0 & 0 & -1/2 & 0 \\ 0 & 0 & 0 & -3/2 \end{pmatrix}, \quad J_+ = \begin{pmatrix} 0 & \sqrt{2} & 0 & 0 \\ 0 & 0 & \sqrt{3} & 0 \\ 0 & 0 & 0 & \sqrt{2} \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad J_- = \begin{pmatrix} 0 & 0 & 0 & 0 \\ \sqrt{2} & 0 & 0 & 0 \\ 0 & \sqrt{3} & 0 & 0 \\ 0 & 0 & \sqrt{2} & 0 \end{pmatrix}.$$

If rotations leave H invariant, all states in the representation space have the same energy, or equivalently the Hilbert space can be written as a direct product space of spaces $V^{(j)}$.

For $j = 1$ another commonly used representation starts with three Cartesian basis states ϵ related to the previous basis via

$$|1, 1\rangle \equiv \epsilon_1 \equiv -\frac{1}{\sqrt{2}}(\epsilon_x + i\epsilon_y), \quad |1, 0\rangle \equiv \epsilon_0 \equiv \epsilon_z, \quad |1, -1\rangle \equiv \epsilon_{-1} \equiv \frac{1}{\sqrt{2}}(\epsilon_x - i\epsilon_y).$$

The spin matrices for that Cartesian basis $\{\epsilon_x, \epsilon_y, \epsilon_z\}$ are

$$J_x = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad J_y = \begin{pmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ -i & 0 & 0 \end{pmatrix}, \quad J_z = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

From the (hermitean) representation $\underline{\Phi}(g)$ of \underline{G} , one obtains the unitary representations $\Phi(g) = \exp(i\underline{\Phi}(g))$ of the full group G . The matrix elements of these unitary representations are known as D -functions. For an the elements A of $SU(2)$ or R of $SO(3)$ these are commonly parametrized with Euler angles, $U_E(\varphi, \theta, \chi) \equiv e^{-i\varphi J_z} e^{-i\theta J_y} e^{-i\chi J_z}$,

$$\langle j, m' | U_E(\varphi, \theta, \chi) | j, m \rangle = D_{m'm}^{(j)}(\varphi, \theta, \chi) = e^{im'\varphi} d_{m'm}^{(j)}(\theta) e^{-im\chi}. \quad (3.19)$$

Infinitesimally (around the identity) the D -functions for $SU(2)$ and $SO(3)$ are the same, e.g.

$$d_{m'm}^{(j)}(\theta) \approx \delta_{m'm} - i\theta(J^2)_{m'm}. \quad (3.20)$$

By moving through the parameter space the D -functions can be extended to global functions for all allowed angles. For those global representations, however, the topological structure of the group is important. If the group is *simply connected*, that is any closed curve in the parameter space can be contracted to a point, any point in the parameter space can be reached in a unique way and any local (infinitesimal) representation can be extended to a global one. This works for $SU(2)$. The group $SO(3)$, however, is not simply connected. There exist two different types of paths, contractable and paths that run from a point at the surface to its antipode. For an element in the group G the corresponding point in the parameter space can be reached in two ways. For a decent (well-defined) global representation, however, the extension from a local one must be unique. For $SO(3)$, the possibility thus exist that some extensions will not be well-defined representations. This turns out to be the case for all half-integer representations of the Lie algebra. Of all groups of which the Lie algebras are homeomorphic, the simply connected group is called the (*universal*) *covering group*, i.e. $SU(2)$ is the covering group of $SO(3)$.

Given a representation, one can look at the *conjugate* representation. Consider the $j = 1/2$ representation of $SU(2)$. If a transformation U acts on χ , the conjugate transformation U^* acts on χ^* . The jump $U \rightarrow U^*$ implies for the generators $\sigma/2 \rightarrow -\sigma^*/2$. For $SU(2)$, however, the conjugate representation is not a new one. Because there exists a matrix ϵ such that $-\sigma^* = \epsilon\sigma\epsilon^{-1}$ one immediately sees that appropriate linear combinations of conjugate states, to be precise the states $\epsilon\chi$ transform via $-\sigma^*$. Explicitly, if $\chi \rightarrow \sigma\chi$, then $\epsilon\chi \rightarrow \epsilon\sigma\chi = \epsilon\sigma\epsilon^{-1}\epsilon\chi = -\sigma^*\epsilon\chi$. Therefore the representation and conjugate representation are *equivalent* in this case (see Exercise 3.3).

3.3 The Lorentz group

In the previous section spin has been introduced as a representation of the rotation group $SU(2)$ without worrying much about the rest of the symmetries of the world. We considered the generators and looked for representations in finite dimensional spaces, e.g. $\sigma/2$ in a two-dimensional (spin 1/2) case. In this section we consider the Poincaré group, consisting of the Lorentz group and translations. To derive some of the properties of the Lorentz group, it is convenient to use a vector notation for the points in Minkowski space. Writing x as a column vector and the metric tensor in matrix form,

$$G = \begin{pmatrix} g_{00} & g_{01} & g_{02} & g_{03} \\ g_{10} & g_{11} & g_{12} & g_{13} \\ g_{20} & g_{21} & g_{22} & g_{23} \\ g_{30} & g_{31} & g_{32} & g_{33} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad (3.21)$$

the scalar product can be written as

$$x^2 = x^T G x \quad (3.22)$$

(Note that x^T is a row vector).

Denoted in terms of column vectors and 4-dimensional matrices one writes for the poincaré transformations $x' = \Lambda x + a$, explicitly

$$x'^{\mu} = (\Lambda)^{\mu\nu} x^{\nu} + a^{\mu} \quad \text{or} \quad x'^{\mu} = \Lambda^{\mu}_{\nu} x^{\nu} + a^{\mu}. \quad (3.23)$$

The proper tensor structure of the matrix element $(\Lambda)^{\mu\nu}$ is a tensor with one upper and one lower index. Invariance of the length of a vector requires for the Lorentz transformations

$$x'^2 = g_{\mu\nu} x'^{\mu} x'^{\nu} = g_{\mu\nu} \Lambda^{\mu}_{\rho} x^{\rho} \Lambda^{\nu}_{\sigma} x^{\sigma} = x^2 = g_{\rho\sigma} x^{\rho} x^{\sigma} \quad (3.24)$$

or

$$\Lambda^{\mu}_{\rho} g_{\mu\nu} \Lambda^{\nu}_{\sigma} = g_{\rho\sigma}, \quad (3.25)$$

which as a matrix equation with $(\Lambda)^{\mu\nu} = \Lambda^{\mu}_{\nu}$ and $(G)^{\mu\nu} = g_{\mu\nu}$ gives

$$(\Lambda^T)^{\rho\mu} (G)^{\mu\nu} (\Lambda)^{\nu\sigma} = (G)^{\rho\sigma}. \quad (3.26)$$

Thus one has the (pseudo-orthogonality) relation

$$\Lambda^T G \Lambda = G \Leftrightarrow G \Lambda^T G = \Lambda^{-1} \Leftrightarrow \Lambda G \Lambda^T = G. \quad (3.27)$$

From this property, it is easy to derive some properties of the matrices Λ :

(i) $\det(\Lambda) = \pm 1$.

proof: $\det(\Lambda^T G \Lambda) = \det(G) \rightarrow (\det \Lambda)^2 = 1$.

($\det \Lambda = +1$ is called *proper*, $\det \Lambda = -1$ is called *improper*).

(ii) $|\Lambda^0_0| \geq 1$.

proof: $(\Lambda^T G \Lambda)^{00} = (G)^{00} = 1 \rightarrow \Lambda^{\mu}_0 g_{\mu\nu} \Lambda^{\nu}_0 = 1 \rightarrow (\Lambda^0_0)^2 - \sum_i (\Lambda^i_0)^2 = 1$.

Using (i) and (ii) the Lorentz transformations can be divided into 4 *classes* (with disconnected parameter spaces)

	$\det \Lambda$	Λ^0_0	
L^{\uparrow}_+	+1	≥ 1	proper orthochrone
L^{\downarrow}_+	+1	≤ -1	proper non-orthochrone
L^{\uparrow}_-	-1	≥ 1	improper orthochrone
L^{\downarrow}_-	-1	≤ -1	improper non-orthochrone

$$(iii) \sum_{i=1}^3 (\Lambda_0^i)^2 = \sum_{i=1}^3 (\Lambda_i^0)^2.$$

proof: use $\Lambda^T G \Lambda = G$ and $\Lambda G \Lambda^T = G$.

Note that Lorentz transformations generated from the identity must belong to L_+^\uparrow , since $I \in L_+^\uparrow$ and $\det \Lambda$ and Λ_0^0 change continuously along a path from the identity. In L_+^\uparrow , one distinguishes rotations and boosts. Rotations around the z-axis are given by $\Lambda_R(\varphi, \hat{z}) = \exp(i\varphi J^3)$, infinitesimally given by $\Lambda_R(\varphi, \hat{z}) \approx I + i\varphi J^3$. Thus

$$\begin{pmatrix} V^{0'} \\ V^{1'} \\ V^{2'} \\ V^{3'} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \varphi & \sin \varphi & 0 \\ 0 & -\sin \varphi & \cos \varphi & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} V^0 \\ V^1 \\ V^2 \\ V^3 \end{pmatrix} \quad \longrightarrow \quad J^3 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 \\ 0 & i & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \quad (3.28)$$

Boosts along the z-direction are given by $\Lambda_B(\eta, \hat{z}) = \exp(-i\eta K^3)$, infinitesimally given by $\Lambda_B(\eta, \hat{z}) \approx I - i\eta K^3$. Thus

$$\begin{pmatrix} V^{0'} \\ V^{1'} \\ V^{2'} \\ V^{3'} \end{pmatrix} = \begin{pmatrix} \cosh \eta & 0 & 0 & \sinh \eta \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \sinh \eta & 0 & 0 & \cosh \eta \end{pmatrix} \begin{pmatrix} V^0 \\ V^1 \\ V^2 \\ V^3 \end{pmatrix} \quad \longrightarrow \quad K^3 = \begin{pmatrix} 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 \end{pmatrix}. \quad (3.29)$$

The parameter η runs from $-\infty < \eta < \infty$. Note that the velocity $\beta = v = v/c$ and the Lorentz contraction factor $\gamma = (1 - \beta^2)^{-1/2}$ corresponding to the boost are related to η as $\gamma = \cosh \eta$, $\beta\gamma = \sinh \eta$. Using these explicit transformations, we have found the generators of rotations, $\mathbf{J} = (J^1, J^2, J^3)$, and those of the boosts, $\mathbf{K} = (K^1, K^2, K^3)$, which satisfy the commutation relations (check!)

$$\begin{aligned} [J^i, J^j] &= i \epsilon^{ijk} J^k, \\ [J^i, K^j] &= i \epsilon^{ijk} K^k, \\ [K^i, K^j] &= -i \epsilon^{ijk} J^k. \end{aligned}$$

The first two sets of commutation relations exhibit the rotational behavior of \mathbf{J} and \mathbf{K} as vectors in $E(3)$ under rotations. From the commutation relations one sees that the boosts (*pure* Lorentz transformations) do not form a group, since the generators \mathbf{K} do not form a closed algebra. The commutator of two boosts in different directions (e.g. the difference of first performing a boost in the y-direction and thereafter in the x-direction and the boosts in reversed order) contains a rotation (in the example around the z-axis). This is the origin of the Thomas precession.

Returning to the global group, it is easy to find the following typical examples from each of the four classes,

$$I = \begin{pmatrix} +1 & 0 & 0 & 0 \\ 0 & +1 & 0 & 0 \\ 0 & 0 & +1 & 0 \\ 0 & 0 & 0 & +1 \end{pmatrix} \in L_+^\uparrow \quad (\text{identity}) \quad (3.30)$$

$$I_t = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & +1 & 0 & 0 \\ 0 & 0 & +1 & 0 \\ 0 & 0 & 0 & +1 \end{pmatrix} \in L_-^\downarrow \quad (\text{time inversion}) \quad (3.31)$$

$$I_s = \begin{pmatrix} +1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \in L_-^\uparrow \quad (\text{space inversion}) \quad (3.32)$$

$$I_s I_t = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \in L_+^\downarrow \quad (\text{space-time inversion}) \quad (3.33)$$

These four transformations form the Vierer group (group of Klein). Multiplying the proper orthochrone transformations with one of them gives all Lorentz transformations.

Summarizing, the Lorentz transformations form a (Lie) group ($\Lambda_1 \Lambda_2$ and Λ^{-1} are again Lorentz transformations). There are six generators. Of the four parts only L_+^\uparrow forms a group. This is a normal subgroup and the factor group is the Vierer group. The extension to the Poincaré group is straightforward. Also this group can be divided into four parts, P_+^\uparrow etc.

3.4 The generators of the Poincaré group

The transformations belonging to P_+^\uparrow are denoted as (Λ, a) , infinitesimally approximated by $(I + \omega, \epsilon)$, explicitly reading

$$\begin{aligned} x'^\mu &= (\Lambda)^{\mu\nu} x^\nu + a^\mu \stackrel{\text{inf}}{=} ((I)^{\mu\nu} + (\omega)^{\mu\nu}) x^\nu + \epsilon^\mu \\ &= \Lambda^\mu{}_\nu x^\nu + a^\mu = (g^\mu{}_\nu + \omega^\mu{}_\nu) x^\nu + \epsilon^\mu. \end{aligned} \quad (3.34)$$

The condition $\Lambda^T G \Lambda = G$ yields

$$(g^\rho{}_\mu + \omega^\rho{}_\mu) g_{\rho\sigma} (g^\sigma{}_\nu + \omega^\sigma{}_\nu) = g_{\mu\nu} \quad \implies \quad \omega_{\nu\mu} + \omega_{\mu\nu} = 0, \quad (3.35)$$

thus $(\omega)^{ij} = -(\omega)^{ji}$ and $(\omega)^{0i} = (\omega)^{i0}$. We therefore find (again) that there are six generators for the Lorentz group, three of which only involve spatial coordinates (rotations) and three others involving time components (boosts).

We now want to find a covariant form of the six generators of the Lorentz transformations, which are obtained by writing the infinitesimal parameters $(\omega)^{\mu\nu}$ in terms of six antisymmetric matrices $(M^{\alpha\beta})^{\mu\nu}$. One can easily convince oneself that

$$(\omega)^{\mu\nu} = \omega^\mu{}_\nu = -\frac{i}{2} \omega_{\alpha\beta} (M^{\alpha\beta})^{\mu\nu}, \quad (3.36)$$

$$(M^{\alpha\beta})^{\mu\nu} = i (g^{\alpha\mu} g^\beta{}_\nu - g^\alpha{}_\nu g^{\beta\mu}) \quad (3.37)$$

The algebra of the generators of the Lorentz transformations can be obtained by an explicit calculation²,

$$[M^{\mu\nu}, M^{\rho\sigma}] = -i (g^{\mu\rho} M^{\nu\sigma} - g^{\nu\rho} M^{\mu\sigma}) - i (g^{\mu\sigma} M^{\rho\nu} - g^{\nu\sigma} M^{\rho\mu}). \quad (3.38)$$

Explicitly, we have for the (infinitesimal) rotations (around z-axis)

$$\Lambda_R = I + i \varphi^3 J^3 = I - i \omega_{12} M^{12}, \quad (3.39)$$

with $\varphi^i = -\frac{1}{2} \epsilon^{ijk} \omega^{jk}$ and $J^i = \frac{1}{2} \epsilon^{ijk} M^{jk}$ (e.g. $J^3 = M^{12}$) and for the (infinitesimal) boosts (along z-axis),

$$\Lambda_B = I + i \eta^3 K^3 = I - i \omega_{30} M^{30}, \quad (3.40)$$

with $\eta^i = \omega^{i0}$ and $K^i = M^{i0}$ (e.g. $K^3 = M^{30}$). Thus the two vector operators \mathbf{J} and \mathbf{K} form under Lorentz transformations an antisymmetric tensor $M^{\mu\nu}$.

In order to find the commutation relations including the translation generators, we continue using covariance arguments. We will just require Poincaré invariance for the generators themselves. The

²These commutation relations are the same as those for the 'quantummechanics' operators $i(x^\mu \partial^\nu - x^\nu \partial^\mu)$

infinitesimal form of any representation of the Poincaré group (thus including the translations) is of the form

$$U(I + \omega, \epsilon) = 1 - \frac{i}{2} \omega_{\alpha\beta} M^{\alpha\beta} + i \epsilon_{\alpha} P^{\alpha}. \quad (3.41)$$

At this point, the generators $M^{\alpha\beta}$ no longer exclusively act in Minkowski space. The generators of the translations are the (momentum) operators P^{μ} . The requirement that P^{μ} transforms as a four-vector (for which we know the explicit behavior from the defining four-dimensional representation), leads to

$$U(\Lambda, a) P^{\mu} U^{\dagger}(\Lambda, a) = (\Lambda^{\mu}_{\nu}) P^{\nu} \quad \text{or} \quad U^{\dagger}(\Lambda, a) P^{\mu} U(\Lambda, a) = (\Lambda^{-1})^{\mu}_{\nu} P^{\nu}, \quad (3.42)$$

(to decide on where the inverse is, check that $U_1 U_2$ corresponds with $\Lambda_1 \Lambda_2$). Infinitesimally,

$$(1 + i \epsilon_{\gamma} P^{\gamma} - \frac{i}{2} \omega_{\alpha\beta} M^{\alpha\beta}) P^{\mu} (1 - i \epsilon_{\delta} P^{\delta} + \frac{i}{2} \omega_{\rho\sigma} M^{\rho\sigma}) = P^{\mu} + \omega^{\mu}_{\nu} P^{\nu}, \quad (3.43)$$

giving the following commutation relations by equating the coefficients of ϵ_{μ} and $\omega_{\mu\nu}$,

$$[P^{\mu}, P^{\nu}] = 0, \quad (3.44)$$

$$[M^{\mu\nu}, P^{\rho}] = -i (g^{\mu\rho} P^{\nu} - g^{\nu\rho} P^{\mu}). \quad (3.45)$$

Note that the commutation relations among the generators $M^{\mu\nu}$ in Eq. 3.38 could have been obtained in the same way. They just state that $M^{\mu\nu}$ transforms as a tensor with two Lorentz indices. Explicitly, writing the generator $P = (H/c, \mathbf{P})$ in terms of the *Hamiltonian* and the *three-momentum* operators, the tensor $M^{\mu\nu}$ in terms of boosts $cK^i = M^{i0}$ and rotations $J^i = \frac{1}{2} \epsilon^{ijk} M^{jk}$, one obtains

$$\begin{aligned} [P^i, P^j] &= [P^i, H] = [J^i, H] = 0, \\ [J^i, J^j] &= i \epsilon^{ijk} J^k, \quad [J^i, P^j] = i \epsilon^{ijk} P^k, \quad [J^i, K^j] = i \epsilon^{ijk} K^k, \\ [K^i, H] &= i P^i, \quad [K^i, K^j] = -i \epsilon^{ijk} J^k / c^2, \quad [K^i, P^j] = i \delta^{ij} H / c^2. \end{aligned} \quad (3.46)$$

We have here reinstated c , because one then sees that by letting $c \rightarrow \infty$ the commutation relations of the *Galilei group*, known from non-relativistic quantum mechanics are obtained.

3.5 Representations of the Poincaré group

In order to label the states in an irreducible representation we construct a maximal set of commuting operators. These define states with specified quantum numbers that are eigenvalues of these generators. For instance, the generators J^2 and J^3 in the case of the rotation group. Taking any one of the states in an irreducible representation, other states in that representation are obtained by the action of operators outside the maximal commuting set. For instance, the generators J_{\pm} in the case of the rotation group.

Of the generators of the Poincaré group we choose first of all the generators P^{μ} , that commute among themselves, as part of the set. The eigenvalues of these will be the four-momentum p^{μ} of the state,

$$P^{\mu} |p, \dots\rangle = p^{\mu} |p, \dots\rangle, \quad (3.47)$$

where p^{μ} is a set of four arbitrary real numbers. To find other generators that commute with P^{μ} we look for Lorentz transformations that leave the four vector p^{μ} invariant. These form a group called the little group associated with that four vector.

$$\begin{aligned} \Lambda^{\mu}_{\nu} p^{\nu} &= p^{\mu} + \omega^{\mu}_{\nu} p^{\nu} = p^{\mu} \\ \Rightarrow \omega_{\mu\nu} p^{\nu} &= 0 \\ \Rightarrow \omega_{\mu\nu} &= \epsilon_{\mu\nu\rho\sigma} p^{\rho} s^{\sigma}, \end{aligned} \quad (3.48)$$

where s is an arbitrary vector of which the length and the component along p^σ are irrelevant, which can thus be chosen spacelike. The elements of the little group thus are

$$\begin{aligned} U(\Lambda(p)) &= \exp\left(-\frac{i}{2}\omega_{\mu\nu}M^{\mu\nu}\right) \\ &= \exp\left(-\frac{i}{2}\epsilon_{\mu\nu\rho\sigma}M^{\mu\nu}p^\rho s^\sigma\right) \end{aligned} \quad (3.49)$$

and

$$\begin{aligned} U(\Lambda(p))|p, \dots\rangle &= \exp\left(-\frac{i}{2}\epsilon_{\mu\nu\rho\sigma}M^{\mu\nu}p^\rho s^\sigma\right)|p, \dots\rangle \\ &= \exp(-is_\mu W^\mu)|p, \dots\rangle \end{aligned} \quad (3.50)$$

where the Pauli-Lubanski operators W^μ are given by

$$W_\mu = -\frac{1}{2}\epsilon_{\mu\nu\rho\sigma}M^{\nu\rho}P^\sigma. \quad (3.51)$$

The following properties follow from the fact that W^μ is a four-vector by construction, of which the components generate the little group of p^μ or (for the third of the following relations) from explicit calculation

$$[M_{\mu\nu}, W_\alpha] = -i(g_{\mu\alpha}W_\nu - g_{\nu\alpha}W_\mu), \quad (3.52)$$

$$[W_\mu, P_\nu] = 0, \quad (3.53)$$

$$[W_\mu, W_\nu] = i\epsilon_{\mu\nu\rho\sigma}W^\rho P^\sigma. \quad (3.54)$$

This will enable us to pick a suitable commuting 'spin' operator. From the algebra of the generators one finds that

$$P^2 = P_\mu P^\mu \quad \text{and} \quad W^2 = W_\mu W^\mu \quad (3.55)$$

commute with all generators and therefore are invariants under Poincaré transformations. These operators are the Casimir operators of the algebra and can be used to define the representations of P_+^\dagger , for which we distinguish the following cases

$$\begin{array}{ll} p^2 = m^2 > 0 & p^0 > 0 \\ p^2 = 0 & p^0 > 0 \\ p^\mu \equiv 0 & \\ p^2 = m^2 > 0 & p^0 < 0 \\ p^2 = 0 & p^0 < 0 \\ p^2 < 0 & \end{array}$$

Only the first two cases correspond to physical states, the third case represents the *vacuum*, while the others have no physical significance.

Massive particles: $p^2 = M^2 > 0$, $p^0 > 0$.

Given the momentum four vector p^μ we choose a tetrad consisting of three orthogonal spacelike unit vectors $n_i(p)$, satisfying

$$\begin{aligned} g_{\mu\nu}n_i^\mu(p)n_j^\nu(p) &= -\delta_{ij}, \\ n_i^\mu(p)p_\mu &= 0, \\ \epsilon_{\mu\nu\rho\sigma}p^\mu n_i^\nu n_j^\rho n_k^\sigma &= M\epsilon_{ijk}. \end{aligned} \quad (3.56)$$

We can write

$$W^\mu(p) = \sum_{i=1}^3 W^i(p) n_i^\mu(p) \quad (3.57)$$

(i.e. $W^i(p) = -W \cdot n_i(p)$).

Having made a covariant decomposition, it is sufficient to choose a particular frame to investigate the coefficients in Eq. 3.57. The best insight in the meaning of the operators W^μ is to sit in the rest frame of the particle and put $P^\mu = (M, \mathbf{0})$. In that case the vectors $n_i(p)$ are just the space directions and $W = (0, M\mathbf{J})$ with components $W^i = (M/2)\epsilon^{ijk}M^{jk}$. The commutation relations

$$[W^i(p), W^j(p)] = iM\epsilon^{ijk}W^k(p), \quad (3.58)$$

show that \mathbf{W}/M can be interpreted as the intrinsic spin.

More generally (fully covariantly) one can proceed by defining $S^i(p) \equiv W^i(p)/M$ and obtain

$$[S^i(p), S^j(p)] = -i \frac{n_i^\mu(p)n_j^\nu(p)}{M^2} \epsilon_{\mu\nu\rho\sigma} W^\rho P^\sigma = i\epsilon^{ijk}S^k(p), \quad (3.59)$$

i.e. the $S^i(p)$ form the generators of an $SU(2)$ subgroup that belongs to the maximal set of commuting operators. Noting that

$$\sum_i (S^i(p))^2 = \frac{1}{M^2} W^\mu(p) W^\nu(p) \sum_i n_{i\mu}(p) n_{i\nu}(p), \quad (3.60)$$

and using the completeness relation

$$\sum_i n_i^\mu(p) n_i^\nu(p) = -\left(g^{\mu\nu} - \frac{p^\mu p^\nu}{M^2}\right), \quad (3.61)$$

one sees that

$$\sum_i (S^i(p))^2 = -\frac{W^2}{M^2}. \quad (3.62)$$

Thus W^2 has the eigenvalues $-M^2 s(s+1)$ with $s = 0, \frac{1}{2}, 1, \dots$. Together with the four momentum states thus can be labeled as

$$|M, s; \mathbf{p}, m_s\rangle, \quad (3.63)$$

where $E = \sqrt{\mathbf{p}^2 + M^2}$ and m_s is the z-component of the spin $-s \leq m_s \leq +s$ (in steps of one).

The explicit construction of the wave function can be done using the D-functions (analogous as the rotation functions). We will not do this but construct them as solutions of a wave equation to be discussed explicitly in chapter 4

Massless particles: $p^2 = M^2 = 0$, $p^0 > 0$.

In this case a set of four independent vectors is chosen starting with a reference frame in which $p^\mu = p_0^\mu$ in the following way:

$$\begin{aligned} p_0 &= (p^0, 0, 0, |p^3|), \\ n_1(p_0) &\equiv (0, 1, 0, 0), \\ n_2(p_0) &\equiv (0, 0, 1, 0), \\ s(p_0) &\equiv (1, 0, 0, -1) \end{aligned} \quad (3.64)$$

such that in an arbitrary frame where the four vectors p , $n_1(p)$, $n_2(p)$ and $s(p)$ are obtained by a Lorentz boost from the reference frame one has the property

$$\epsilon_{\mu\nu\rho\sigma} s^\mu n_1^\nu n_2^\rho p^\sigma > 0. \quad (3.65)$$

The above does actually include the massive case. The vector $W^\mu(p)$ now can be expanded

$$W^\mu(p) = W^1(p) n_1^\mu(p) + W^2(p) n_2^\mu(p) + \lambda(p) p^\mu + W^s(p) s^\mu(p), \quad (3.66)$$

where $W \cdot p = 0$ implies that $W^s(p) = -(M^2/p \cdot s) \lambda(p)$. The algebra of W^1 , W^2 and λ is derived from the general commutation relations for $[W_\mu, W_\nu]$, and gives

$$[W^1(p), W^2(p)] = i M^2 \lambda(p), \quad (3.67)$$

$$[\lambda(p), W^1(p)] = i W^2(p), \quad (3.68)$$

$$[\lambda(p), W^2(p)] = -i W^1(p). \quad (3.69)$$

We reproduce the algebra for the massive case with $W^3(p) = M \lambda(p)$. But if $M = 0$ one has a different algebra (Eq. 3.67 has a vanishing right hand side). This algebra is (for instance) isomorphic to the one generated by rotations and translations in a 2-dimensional Euclidean plane (see exercises). The eigenvalues of W^1 and W^2 (in that case corresponding to the translations) thus can take any continuous values, implying continuous values for the spin (actually for $W^i = M S^i$). We don't know of any physical relevance, however, and the eigenvalues of W^1 and W^2 are set to zero (corresponding to the limit $M \rightarrow 0$ for the $W^\mu W_\mu$ eigenvalues of $M^2 s(s+1) \rightarrow 0$). Thus we have

$$P^2, P^\mu, W^\mu(p) = \lambda(p) P^\mu, W^2 \quad (3.70)$$

as a commuting set with zero eigenvalues for P^2 and W^2 . The meaning of $\lambda(p)$ is most easily seen by comparing for a momentum eigenstate

$$W^0(p) = \lambda(p) |\mathbf{p}\rangle$$

with

$$W^0(p) = -\frac{1}{2} \epsilon^{0ijk} M_{ij} p_k = \mathbf{J} \cdot \mathbf{p}.$$

Thus

$$\lambda(p) = \frac{\mathbf{J} \cdot \mathbf{p}}{|\mathbf{p}|}, \quad (3.71)$$

which is called the helicity. Note that angular momentum does not contribute to helicity as $\mathbf{L} \cdot \mathbf{p} = 0$. A massless particle, thus, is characterized by its momentum and the helicity,

$$|\mathbf{p}, \lambda\rangle, \quad (3.72)$$

which can take any *integer* or *half-integer* value.

The vacuum: $p^\mu = 0$.

This state is in physical applications nondegenerate and is denoted by

$$|0\rangle. \quad (3.73)$$

One has $P^\mu |0\rangle = W^\mu |0\rangle = 0$ and the state is invariant under Lorentz transformations, $U(\Lambda, a) |0\rangle = |0\rangle$.

Exercises

Exercise 3.1

- (a) Derive from the relation that defines the mapping of $SU(2)$ matrices A into $SO(3)$ matrices R_A in Eq. 3.13,

$$\boldsymbol{\sigma} \cdot R_A \mathbf{a} \equiv A(\boldsymbol{\sigma} \cdot \mathbf{a})A^{-1}$$

(valid for any vector \mathbf{a}), where $A = \exp(i\varphi \cdot \boldsymbol{\sigma}/2) = \exp(i\varphi \boldsymbol{\sigma} \cdot \hat{\mathbf{n}}/2)$, the action of R_A

$$\begin{aligned} \mathbf{b} \cdot R_A \mathbf{a} &= \mathbf{b} \cdot \mathbf{a} \cos \varphi + \hat{\mathbf{n}} \cdot (\mathbf{b} \times \mathbf{a}) \sin \varphi + 2(\mathbf{b} \cdot \hat{\mathbf{n}})(\mathbf{a} \cdot \hat{\mathbf{n}}) \sin^2(\varphi/2) \\ &= \mathbf{b} \cdot \mathbf{a} + \mathbf{b} \cdot (\hat{\mathbf{a}} \times \mathbf{n}) \sin \varphi - 2(\mathbf{b} \cdot \mathbf{a} - (\mathbf{b} \cdot \hat{\mathbf{n}})(\mathbf{a} \cdot \hat{\mathbf{n}})) \sin^2(\varphi/2) \end{aligned}$$

for any vectors \mathbf{a} and \mathbf{b} .

Note: Recall (or derive) that the Pauli matrices $\boldsymbol{\sigma}$ satisfy

$$\sigma_i \sigma_j = \delta_{ij} + i \epsilon_{ijk} \sigma_k$$

and

$$\begin{aligned} \text{Tr}(\sigma_i \sigma_j) &= 2 \delta_{ij}, \\ \text{Tr}(\sigma_i \sigma_j \sigma_k) &= 2i \epsilon_{ijk}, \\ \text{Tr}(\sigma_i \sigma_j \sigma_k \sigma_l) &= 2(\delta_{ij} \delta_{kl} - \delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk}). \end{aligned}$$

- (b) Use the result under (a) to derive the matrix elements of a rotation around the z-axis and show that it indeed represents the $SO(3)$ rotation $R_A = \exp(i\varphi L_z)$ in which the L_z is one of the (defining) generators of $SO(3)$.

Exercise 3.2

Show that for a unitary operator $U = \exp[i\alpha_k J_k]$ with real coefficients α_k , the operators J_k must be hermitean. Show that if $\det(U) = 1$, the trace of the operator J_k is zero (Convince yourself that $\det(e^A) = e^{\text{Tr}A}$ by diagonalizing A).

Exercise 3.3

Consider the representation $A = \exp(i\boldsymbol{\alpha} \cdot \mathbf{J})$ with the Lie-algebra representations $\mathbf{J} = \boldsymbol{\sigma}/2$ describing spin rotations for a spin 1/2 particle. For a spin 1/2 antiparticle the representation A^* is needed. Writing $A^* = \exp(i\boldsymbol{\alpha} \cdot \mathbf{J}^c)$, what are the matrices \mathbf{J}^c in the Lie-algebra representation describing the spin rotations for two-component 'anti-spinors'. Show that the Lie-algebra representations \mathbf{J} and \mathbf{J}^c are equivalent, i.e. show that *one* matrix ϵ exist such that $\mathbf{J}^c = \epsilon \mathbf{J} \epsilon^\dagger$.

Comment: this establishes that for spin 1/2 one doesn't need to talk about 'anti-spin' for the antiparticle. Both particle and antiparticle have 'spin 1/2'. This is true for $SU(2)$ but it is, for instance, not true for $SU(3)$ (flavor or color properties of particles know two distinct defining 3-dimensional representation, triplet and anti-triplet representation, respectively).

Exercise 3.4

- (a) Show that if the states in a Hilbert space (unitarily) transform as $\phi' = U \phi = e^{iJ} \phi$ and O is an operator acting on ϕ (in that Hilbert space), that the transformation for the operators is given by

$$O' = U O U^{-1} = O + i[J, O] + \dots$$

- (b) Show that this transformation for the position operator in quantum mechanics indeed gives

$$U(\mathbf{a}) \mathbf{r} U^{-1}(\mathbf{a}) = \exp(i\mathbf{a} \cdot \mathbf{p}) \mathbf{r} \exp(-i\mathbf{a} \cdot \mathbf{p}) = \mathbf{r} + \mathbf{a}.$$

and calculate $U(\mathbf{a}) \mathbf{p} U^{-1}(\mathbf{a})$

Exercise 3.5

In classical mechanics (Galilean invariance) boosts describe invariance of physical laws when viewed from a moving frame, i.e. they describe

$$\mathbf{r} \longrightarrow \mathbf{r}' = \mathbf{r} - \mathbf{u} t.$$

- (a) In classical mechanics, boost invariance leads to a conserved quantity. For a single free particle or for the free-moving center of mass system that is the quantity $\mathbf{K} = m\mathbf{r} - t\mathbf{p}$. Determine the commutation relations for the corresponding operators K_i in quantum mechanics, in particular those with the momentum operator p_j and the Hamiltonian $H = mc^2 + \mathbf{p}^2/2m$.
- (b) In quantum mechanics one has the Ehrenfest relation

$$i\hbar \frac{d\langle O \rangle}{dt} = \langle [H, O] \rangle + i\hbar \left\langle \frac{\partial O}{\partial t} \right\rangle.$$

Show that boost invariance for a free-moving particle (the commutation relations under (a) being valid), indeed assures $d\langle K \rangle/dt = 0$, although that in this case is a relation that is found already from the Ehrenfest relation for $O = r_i$.

- (c) Give the unitary operator for boosts, i.e. the operator $U(\mathbf{u})$ that gives

$$\begin{aligned} U(\mathbf{u}) \mathbf{r} U^{-1}(\mathbf{u}) &= \mathbf{r} - \mathbf{u} t, \\ U(\mathbf{u}) \mathbf{p} U^{-1}(\mathbf{u}) &= \mathbf{p} - m \mathbf{u}. \end{aligned}$$

Exercise 3.6

Consider rotations and translations in a plane (two-dimensional Euclidean space). Use a 3-dimensional matrix representation or consider the generators of rotations and translations in the space of functions on the (x-y) plane. From either of these, derive the commutation relations and show that the algebra is isomorphic to (i.e., there is a one to one mapping) the algebra in Eqs 3.67 to 3.69 of the Pauli-Lubanski operators for massless particles.

Exercise 3.7 (optional)

One might wonder if it is actually possible to write down a set of operators that generate the Poincaré transformations, consistent with the (canonical) commutation relations of a quantum theory. This is possible for a single particle. Do this by showing that the set of operators,

$$\begin{aligned} H &= \sqrt{\mathbf{p}^2 c^2 + m^2 c^4}, \\ \mathbf{P} &= \mathbf{p}, \\ \mathbf{J} &= \mathbf{r} \times \mathbf{p} + \mathbf{s}, \\ \mathbf{K} &= \frac{1}{2c^2} (\mathbf{r}H + H\mathbf{r}) - t\mathbf{p} + \frac{\mathbf{p} \times \mathbf{s}}{H + mc^2}. \end{aligned}$$

satisfy the commutation relations of the Poincaré group if the position, momentum and spin operators satisfy the canonical commutation relations, $[r^i, p^j] = i\delta^{ij}$ and $[s^i, s^j] = i\epsilon^{ijk} s^k$; the others vanish, $[r^i, r^j] = [p^i, p^j] = [r^i, s^j] = [p^i, s^j] = 0$.

Hint: for the Hamiltonian, show first the operator identity $[\mathbf{r}, f(\mathbf{p})] = i \nabla_{\mathbf{p}} f(\mathbf{p})$; if you don't want to do this in general, you might just check relations involving \mathbf{J} or \mathbf{K} by taking some (relevant) explicit components.

Comment: extending this to more particles is a highly non-trivial procedure, but it can be done, although the presence of an interaction term $V(\mathbf{r}_1, \mathbf{r}_2)$ inevitably leads to interaction terms in the boost operators. These do not matter in the non-relativistic limit ($c \rightarrow \infty$), that's why many-particle non-relativistic quantum mechanics is 'easy'.

Chapter 4

The Dirac equation

4.1 The Lorentz group and $SL(2, C)$

Instead of the generators \mathbf{J} and \mathbf{K} of the homogeneous Lorentz transformations we can use the (hermitean) combinations

$$\mathbf{A} = \frac{1}{2}(\mathbf{J} + i\mathbf{K}), \quad (4.1)$$

$$\mathbf{B} = \frac{1}{2}(\mathbf{J} - i\mathbf{K}), \quad (4.2)$$

which satisfy the commutation relations

$$[A^i, A^j] = i\epsilon^{ijk}A^k, \quad (4.3)$$

$$[B^i, B^j] = i\epsilon^{ijk}B^k, \quad (4.4)$$

$$[A^i, B^j] = 0. \quad (4.5)$$

This shows that the Lie algebra of the Lorentz group is identical to that of $SU(2) \otimes SU(2)$. This tells us how to find the representations of the group. They will be labeled by two angular momenta corresponding to the A and B groups, respectively, (j, j') . Special cases are the following representations:

$$\text{Type I: } (j, 0) \quad \mathbf{K} = -i\mathbf{J} \quad (\mathbf{B} = 0), \quad (4.6)$$

$$\text{Type II: } (0, j) \quad \mathbf{K} = i\mathbf{J} \quad (\mathbf{A} = 0). \quad (4.7)$$

From the considerations above, it also follows directly that the Lorentz group is homeomorphic with the group $SL(2, C)$, similarly as the homeomorphism between $SO(3)$ and $SU(2)$. The group $SL(2, C)$ is the group of complex 2×2 matrices with determinant one. It is simply connected and forms the covering group of L_+^\uparrow . It is easy to see that these matrices can be written as a product of a unitary matrix U and a hermitean matrix H ,

$$M = \exp\left(\frac{i}{2}\boldsymbol{\varphi} \cdot \boldsymbol{\sigma}\right) \exp\left(\pm\frac{1}{2}\boldsymbol{\eta} \cdot \boldsymbol{\sigma}\right) = \begin{Bmatrix} U(\boldsymbol{\varphi})H(\boldsymbol{\eta}) \\ U(\boldsymbol{\varphi})\overline{H}(\boldsymbol{\eta}) \end{Bmatrix}, \quad (4.8)$$

with $\boldsymbol{\eta} = \eta\hat{\mathbf{n}}$ and $\boldsymbol{\varphi} = \varphi\hat{\mathbf{n}}$, where we restrict (for fixed $\hat{\mathbf{n}}$) the parameters $0 \leq \varphi \leq 2\pi$ and $0 \leq \eta < \infty$. With this choice of parameter-spaces the plus and minus signs are actually relevant. They precisely correspond to the two types of representations that we have seen before, becoming the defining representations of $SL(2, C)$:

$$\text{Type I (denoted } M): \quad \mathbf{J} = \frac{\boldsymbol{\sigma}}{2}, \quad \mathbf{K} = -i\frac{\boldsymbol{\sigma}}{2}, \quad (4.9)$$

$$\text{Type II (denoted } \overline{M}): \quad \mathbf{J} = \frac{\boldsymbol{\sigma}}{2}, \quad \mathbf{K} = +i\frac{\boldsymbol{\sigma}}{2}, \quad (4.10)$$

Let us investigate the defining (two-dimensional) representations of $SL(2, C)$. One defines spinors ξ and η transforming similarly under unitary rotations ($U^\dagger = U^{-1}$, $\bar{U} \equiv (U^\dagger)^{-1} = U$)

$$\begin{aligned} \xi &\rightarrow U\xi, & \eta &\rightarrow U\eta, \\ U(\boldsymbol{\varphi}) &= \exp(i\boldsymbol{\varphi} \cdot \boldsymbol{\sigma}/2) \end{aligned} \quad (4.11)$$

but differently under hermitean boosts ($H^\dagger = H$, $\bar{H} \equiv (H^\dagger)^{-1} = H^{-1}$), namely

$$\begin{aligned} \xi &\rightarrow H\xi, & \eta &\rightarrow \bar{H}\eta, \\ H(\boldsymbol{\eta}) &= \exp(\boldsymbol{\eta} \cdot \boldsymbol{\sigma}/2), \\ \bar{H}(\boldsymbol{\eta}) &= \exp(-\boldsymbol{\eta} \cdot \boldsymbol{\sigma}/2). \end{aligned} \quad (4.12)$$

A practical boost for the spinors is the one transforming from the rest frame to the frame with momentum \mathbf{p} . With $E = M\gamma = M \cosh(\eta)$ and $\mathbf{p} = M\beta\gamma\hat{\mathbf{n}} = M \sinh(\eta) \hat{\mathbf{n}}$ it is given by

$$H(\mathbf{p}) = \exp\left(\frac{\boldsymbol{\eta} \cdot \boldsymbol{\sigma}}{2}\right) = \cosh\left(\frac{\eta}{2}\right) + \boldsymbol{\sigma} \cdot \hat{\mathbf{n}} \sinh\left(\frac{\eta}{2}\right) = \frac{M + E + \boldsymbol{\sigma} \cdot \mathbf{p}}{\sqrt{2M(E + M)}}, \quad (4.13)$$

Also useful is the relation $H^2(\mathbf{p}) = \tilde{\sigma}^\mu p_\mu / M = (E + \boldsymbol{\sigma} \cdot \mathbf{p}) / M$.

The sets of four operators defined by definitions

$$\boldsymbol{\sigma}^\mu \equiv (\mathbf{1}, \boldsymbol{\sigma}), \quad \tilde{\boldsymbol{\sigma}}^\mu \equiv (\mathbf{1}, -\boldsymbol{\sigma}), \quad (4.14)$$

satisfying $\text{Tr}(\boldsymbol{\sigma}^\mu \tilde{\boldsymbol{\sigma}}^\nu) = -2g^{\mu\nu}$ and $\text{Tr}(\boldsymbol{\sigma}^\mu \boldsymbol{\sigma}^\nu) = 2g^\mu{}_\nu = 2\delta^{\mu\nu}$ (the matrices, thus, are not covariant!) can be used to establish the homeomorphism between L_+^\uparrow and $SL(2, C)$. The following relations for rotations and boosts are useful,

$$U \boldsymbol{\sigma}^\mu a_\mu U^{-1} = \boldsymbol{\sigma}^\mu \Lambda_{R\alpha\mu}, \quad H \boldsymbol{\sigma}^\mu a_\mu H^{-1} = \boldsymbol{\sigma}^\mu \Lambda_{B\alpha\mu} \quad (4.15)$$

$$\bar{U} \tilde{\boldsymbol{\sigma}}^\mu a_\mu \bar{U}^{-1} = \tilde{\boldsymbol{\sigma}}^\mu \Lambda_{R\alpha\mu}, \quad \bar{H} \tilde{\boldsymbol{\sigma}}^\mu a_\mu \bar{H}^{-1} = \tilde{\boldsymbol{\sigma}}^\mu \Lambda_{B\alpha\mu}. \quad (4.16)$$

Non-equivalence of M and \bar{M}

One might wonder if type I and type II representations are not equivalent, i.e. if there does not exist a unitary matrix S , such that $\bar{M} = SMS^{-1}$. In fact, there exists the matrix

$$\epsilon = i\sigma^2, \quad (\epsilon^* = \epsilon; \epsilon^{-1} = \epsilon^\dagger = \epsilon^T = -\epsilon), \quad (4.17)$$

that can be used to relate $\tilde{\boldsymbol{\sigma}}^{\mu*} = \epsilon^\dagger \boldsymbol{\sigma}^\mu \epsilon = \epsilon^{-1} \boldsymbol{\sigma}^\mu \epsilon$ and hence

$$U^* = \epsilon^\dagger U \epsilon = \epsilon^\dagger \bar{U} \epsilon, \quad H^* = \epsilon^\dagger \bar{H} \epsilon = \epsilon^{-1} \bar{H} \epsilon. \quad (4.18)$$

As already discussed for $SU(2)$, the first equation shows that the conjugate representation with spinors χ^* transforming according to $-\boldsymbol{\sigma}^*/2$ is equivalent with the ordinary two-dimensional representation ($\pm\epsilon\chi^*$ transforms according to $\boldsymbol{\sigma}/2$). This is not true for the two-component spinors in $SL(2, C)$ or L_+^\uparrow . Eqs 4.18 show that $\pm\epsilon\xi^*$ transforms as a type-II (η) spinor, while $\pm\epsilon\eta^*$ transforms as a type-I spinor. This shows that $M^* \simeq \bar{M}$.

Note that $M^* = \epsilon^\dagger \bar{M} \epsilon$ has its equivalent for the Lorentz transformations $\Lambda(\bar{M})$ and $\Lambda(M^*)$. They are connected via $\Lambda(M^*) = \Lambda^{-1}(\epsilon)\Lambda(\bar{M})\Lambda(\epsilon)$ within L_+^\uparrow because $i\sigma^2 = U(\pi\hat{\mathbf{y}})$ showing that $\epsilon \in SL(2, C)$ and, thus, $\Lambda(\epsilon) \in L_+^\uparrow$. The Lorentz transformations $\Lambda(M)$ and $\Lambda(M^*)$, however, cannot be connected by a Lorentz transformation belonging to L_+^\uparrow . One has

$$\Lambda(M^*) = I_2 \Lambda(M) I_2^{-1}, \quad (4.19)$$

where

$$I_2 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (4.20)$$

The matrix I_2 does not belong to L_+^\uparrow , but to L_-^\uparrow . Thus M^* and M are not equivalent within $SL(2, C)$.

4.2 Spin 1/2 representations of the Lorentz group

Both representations $(0, \frac{1}{2})$ and $(\frac{1}{2}, 0)$ of $SL(2, C)$ are suitable for representing spin 1/2 particles. The angular momentum operators J^i are represented by $\sigma^i/2$, that have the correct commutation relations. Although these operators cannot be used to label the representations, but rather the operators $W^i(p)$ should be used, we have seen that in the rest frame $W^i(p)/M \rightarrow J^i$, and the angular momentum in the rest frame is what we are familiar with as the spin of a particle. We also have seen that the representations $(0, \frac{1}{2})$ and $(\frac{1}{2}, 0)$ are inequivalent, i.e. they cannot be connected by a unitary transformation. Within the Lorentz group, they can be connected, but by a transformation belonging to the class P_-^\uparrow , as we have seen in section 4.1. The representing member of the class P_-^\uparrow is the parity or space inversion operator I_s . A parity transformation changes a^μ into \tilde{a}^μ , where $a = (a^0, \mathbf{a})$ and $\tilde{a} = (a^0, -\mathbf{a})$. It has the same effect as lowering the indices. This provides the easiest way of seeing what is happening, e.g. $\epsilon^{\mu\nu\rho\sigma}$ will change sign, the a^{ij} elements of a tensor will not change sign. Examples (with between brackets the Euclidean assignments) are

$$\begin{aligned} \mathbf{r} &\longrightarrow -\mathbf{r} && \text{(vector),} \\ t &\longrightarrow t && \text{(scalar),} \\ \mathbf{p} &\longrightarrow -\mathbf{p} && \text{(vector),} \\ H &\longrightarrow H && \text{(scalar),} \\ \mathbf{J} &\longrightarrow \mathbf{J} && \text{(axial vector),} \\ \lambda(p) &\longrightarrow -\lambda(p) && \text{(pseudoscalar),} \\ \mathbf{K} &\longrightarrow -\mathbf{K} && \text{(vector).} \end{aligned}$$

The behavior is the same for classical quantities, generators, etc. From the definition of the representations $(0, \frac{1}{2})$ and $(\frac{1}{2}, 0)$ (via operators \mathbf{J} and \mathbf{K}) one sees that under parity

$$(0, \frac{1}{2}) \longrightarrow (\frac{1}{2}, 0). \quad (4.21)$$

In nature parity turns (often) out to be a good quantum number for elementary particle states. For the spin 1/2 representations of the Poincaré group including parity we, therefore, must combine the representations, i.e. consider the four component spinor that transforms under a Lorentz transformation as

$$u = \begin{pmatrix} \xi \\ \eta \end{pmatrix} \longrightarrow \begin{pmatrix} M(\Lambda) & 0 \\ 0 & \overline{M}(\Lambda) \end{pmatrix} \begin{pmatrix} \xi \\ \eta \end{pmatrix}, \quad (4.22)$$

where $\overline{M}(\Lambda) = \epsilon M^*(\Lambda) \epsilon^{-1}$ with ϵ given in Eq. 4.17. For a particle at rest only angular momentum is important and we can choose $\xi(\mathbf{0}, m) = \eta(\mathbf{0}, m) = \chi_m$, the well-known two-component spinor for a spin 1/2 particle. Taking $M(\Lambda) = H(\mathbf{p})$, the boost in Eq. 4.13, we obtain for the two components of u which we will refer to as *chiral right* and *chiral left* components,

$$u(\mathbf{p}, m) = \begin{pmatrix} u_R \\ u_L \end{pmatrix} = \begin{pmatrix} H(\mathbf{p}) & 0 \\ 0 & \overline{H}(\mathbf{p}) \end{pmatrix} \begin{pmatrix} \chi_m \\ \chi_m \end{pmatrix}, \quad (4.23)$$

with

$$H(\mathbf{p}) = \frac{E + M + \boldsymbol{\sigma} \cdot \mathbf{p}}{\sqrt{2M(E + M)}}, \quad (4.24)$$

$$\bar{H}(\mathbf{p}) = \frac{E + M - \boldsymbol{\sigma} \cdot \mathbf{p}}{\sqrt{2M(E + M)}} = H^{-1}(\mathbf{p}). \quad (4.25)$$

It is straightforward to eliminate χ_m and obtain the following constraint on the components of u ,

$$\begin{pmatrix} 0 & H^2(p) \\ H^{-2}(p) & 0 \end{pmatrix} \begin{pmatrix} u_R \\ u_L \end{pmatrix} = \begin{pmatrix} u_R \\ u_L \end{pmatrix}, \quad (4.26)$$

or explicitly in the so-called Weyl representation

$$\begin{pmatrix} -M & E + \boldsymbol{\sigma} \cdot \mathbf{p} \\ E - \boldsymbol{\sigma} \cdot \mathbf{p} & -M \end{pmatrix} \begin{pmatrix} u_R \\ u_L \end{pmatrix} = 0, \quad (4.27)$$

which is an explicit realization of the (momentum space) Dirac equation, which in general is a linear equation in p^μ ,

$$(p_\mu \gamma^\mu - M) u(p) \equiv (\not{p} - M) u(p) = 0, \quad (4.28)$$

where γ^μ are 4×4 matrices called the Dirac matrices¹.

As in section 2 we can use $P_\mu = i\partial_\mu$ as a representation for the momenta (translation operators) in function space. This leads to the Dirac equation for $\psi(x) = u(p)e^{-ip \cdot x}$ in coordinate space,

$$(i\gamma^\mu \partial_\mu - M) \psi(x) = 0, \quad (4.29)$$

which is a covariant (linear) first order differential equation. It is of a form that we also played with in Exercise 2.7. The general requirements for the γ matrices are thus easily obtained. Applying $(i\gamma^\mu \partial_\mu + M)$ from the left gives

$$(\gamma^\mu \gamma^\nu \partial_\mu \partial_\nu + M^2) \psi(x) = 0. \quad (4.30)$$

Since $\partial_\mu \partial_\nu$ is symmetric, this can be rewritten

$$\left(\frac{1}{2} \{ \gamma^\mu, \gamma^\nu \} \partial_\mu \partial_\nu + M^2 \right) \psi(x) = 0. \quad (4.31)$$

To achieve also that the energy-momentum relation $p^2 = M^2$ is satisfied for $u(p)$, one must require that for $\psi(x)$ the Klein-Gordon relation $(\square + M^2) \psi(x) = 0$ is valid for each component separately). From this one obtains the Clifford algebra for the Dirac matrices,


$$\{ \gamma^\mu, \gamma^\nu \} = 2 g^{\mu\nu}, \quad (4.32)$$

suppressing on the RHS the identity matrix in Dirac space. The explicit realization appearing in Eq. 4.27 is known as the Weyl representation. We will discuss another explicit realization of this algebra in the next section.

We first want to investigate if the Dirac equation solves the problem with the negative densities and negative energies. The first indeed is solved. To see this consider the equation for the hermitean conjugate spinor ψ^\dagger (noting that $\gamma_0^\dagger = \gamma_0$ and $\gamma_i^\dagger = -\gamma_i$),

$$\psi^\dagger \left(-i\gamma^0 \overleftarrow{\partial}_0 + i\gamma^i \overleftarrow{\partial}_i - M \right) = 0. \quad (4.33)$$

¹We define for a four vector a the contraction $\not{a} = a^\mu \gamma_\mu$

Figure 4.1: *The Dirac sea of negative energy states and antiparticles.*

Multiplying with γ^0 on the right and pulling it through one finds a covariant equation

$$\bar{\psi}(x) \left(i\gamma^\mu \overleftarrow{\partial}_\mu + M \right) = 0, \quad (4.34)$$

for the *adjoint* spinor $\bar{\psi} \equiv \psi^\dagger \gamma_0$. From the equations for ψ and $\bar{\psi}$ one immediately sees that

$$j^\mu = \bar{\psi} \gamma^\mu \psi \quad (4.35)$$

is a conserved probability current (exercise!). The density

$$j^0 = \bar{\psi} \gamma^0 \psi = \psi^\dagger \psi \quad (4.36)$$

is indeed positive and j^0 can serve as a probability density.

The second problem encountered before, the one of the negative energy states, remains. This can most easily be seen in the particle rest frame, where

$$\gamma^0 p_0 \psi = M \psi \quad \longrightarrow \quad E \psi = M \gamma^0 \psi. \quad (4.37)$$

Using the explicit form of γ^0 in the Weyl representation (see Eq. 4.27), one sees that there are two positive and two negative eigenvalues, $E = +M$ (twice) and $E = -M$ (twice),

This problem was solved by Dirac through the introduction of a negative energy sea. Relying on the Pauli exclusion principle for spin 1/2 particles, Dirac supposed that the negative energy states were already completely filled and the exclusion principle prevents any more particles to enter the sea of negative energy states. The Dirac sea forms the vacuum.

From the vacuum a particle can be removed. This hole forms an *antiparticle* with the same mass, but with properties such that it can be annihilated by the particle (e.g. opposite charge). We will see how this is properly implemented (also for bosons!) when quantizing fields.

4.3 General representations of γ matrices and Dirac spinors

The general algebra for the Dirac matrices is

$$\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}. \quad (4.38)$$

Two often used explicit representations are the following²: The *standard* representation:

$$\gamma^0 = \rho^3 \otimes \mathbf{1} = \begin{pmatrix} \mathbf{1} & 0 \\ 0 & -\mathbf{1} \end{pmatrix}, \quad \gamma^k = i\rho^2 \otimes \sigma^k = \begin{pmatrix} 0 & \sigma^k \\ -\sigma^k & 0 \end{pmatrix}; \quad (4.39)$$

²We use $\rho^i \otimes \sigma^j$ with both ρ and σ being the standard 2×2 Pauli matrices.

The Weyl (or chiral) representation:

$$\gamma^0 = \rho^1 \otimes \mathbf{1} = \begin{pmatrix} 0 & \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix}, \quad \gamma^k = -i\rho^2 \otimes \sigma^k = \begin{pmatrix} 0 & -\sigma^k \\ \sigma^k & 0 \end{pmatrix}. \quad (4.40)$$

Different representations can be related to each other by unitary transformations,

$$\gamma_\mu \longrightarrow S\gamma_\mu S^{-1}, \quad (4.41)$$

$$\psi \longrightarrow S\psi. \quad (4.42)$$

We note that the explicit matrix taking us from the Weyl representation to the standard representation, $(\gamma_\mu)_{S.R.} = S(\gamma_\mu)_{W.R.} S^{-1}$, is

$$S = \frac{1}{\sqrt{2}} \begin{pmatrix} \mathbf{1} & \mathbf{1} \\ \mathbf{1} & -\mathbf{1} \end{pmatrix}. \quad (4.43)$$

For all representations one has

$$\gamma_\mu^\dagger = \gamma_0 \gamma_\mu \gamma_0, \quad (4.44)$$

and an adjoint spinor defined by

$$\bar{\psi} = \psi^\dagger \gamma_0. \quad (4.45)$$

Another matrix which is often used is γ_5 defined as

$$\gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3 = -i\gamma_0\gamma_1\gamma_2\gamma_3 = \frac{i\epsilon_{\mu\nu\rho\sigma}}{4!}\gamma^\mu\gamma^\nu\gamma^\rho\gamma^\sigma. \quad (4.46)$$

It satisfies $\{\gamma_5, \gamma^\mu\} = 0$ and explicitly one has

$$(\gamma_5)_{S.R.} = \rho^1 \otimes \mathbf{1} = \begin{pmatrix} 0 & \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix}, \quad (\gamma_5)_{W.R.} = \rho^3 \otimes \mathbf{1} = \begin{pmatrix} \mathbf{1} & 0 \\ 0 & -\mathbf{1} \end{pmatrix}. \quad (4.47)$$

For instance in the Weyl representation (but valid generally), it is easy to see that

$$P_{R/L} = \frac{1}{2}(1 \pm \gamma_5) \quad (4.48)$$

are projection operators, that project out chiral right/left states, which in the case of the Weyl representation are just the upper and lower components.

Lorentz invariance

The Lorentz transformations can also be written in terms of Dirac matrices. For example, the rotation and boost generators in Weyl representation in Eqs 4.9 and 4.10 are

$$J^3 = M^{12} = \frac{1}{2} \begin{pmatrix} \sigma^3 & 0 \\ 0 & \sigma^3 \end{pmatrix} = \frac{i}{4}[\gamma^1, \gamma^2],$$

$$K^3 = M^{30} = \frac{1}{2} \begin{pmatrix} -i\sigma^3 & 0 \\ 0 & i\sigma^3 \end{pmatrix} = -\frac{i}{4}[\gamma^3, \gamma^0],$$

or in general one has the transformation

$$\psi \longrightarrow L\psi = \exp\left(-\frac{i}{2}\omega_{\rho\sigma}S^{\rho\sigma}\right)\psi, \quad (4.49)$$

with $-iS^{\rho\sigma} = \frac{i}{2}\sigma^{\rho\sigma} = -\frac{1}{4}[\gamma^\rho, \gamma^\sigma]$. We note that $\psi \longrightarrow L\psi$ and $\bar{\psi} \longrightarrow \bar{\psi}L^{-1}$, while $L^{-1}\gamma^\mu L = \Lambda_\nu^\mu\gamma^\nu$. The latter assures Lorentz invariance of the Dirac equation (see Exercise 4.3)

Parity

There are a number of symmetries in the Dirac equation, e.g. parity. It is easy to convince oneself that if $\psi(x)$ is a solution of the Dirac equation,

$$(i\cancel{\partial} - M)\psi(x) = 0, \quad (4.50)$$

one can apply space inversion, $x = (t, \mathbf{x}) \rightarrow \tilde{x} = (t, -\mathbf{x})$ and via a few manipulations obtain again the Dirac equation

$$(i\cancel{\partial} - M)\psi^P(x) = 0, \quad (4.51)$$

but with $\psi^P(x) \equiv \eta_P \gamma_0 \psi(\tilde{x})$ (Exercise 4.6). Note that we have (as expected) explicitly in Weyl representation in Dirac space

$$\psi = \begin{pmatrix} \xi \\ \eta \end{pmatrix} \xrightarrow{\mathcal{P}} \psi^P = \gamma^0 \psi = \begin{pmatrix} \eta \\ \xi \end{pmatrix}. \quad (4.52)$$

Charge conjugation

The existence of positive and negative energy solutions implies another symmetry in the Dirac equation. This symmetry does not change the spin 1/2 nature, but it does, for instance, reverse the charge of the particle. As with parity we look for a transformation, called *charge conjugation*, that brings $\psi \rightarrow \psi^c$, which is again a solution of the Dirac equation. Starting with $(i\cancel{\partial} - M)\psi(x) = 0$ we note that by hermitean conjugating and transposing the Dirac equation one obtains

$$(i\gamma^{\mu T} \partial_\mu + M)\bar{\psi}^T(x) = 0. \quad (4.53)$$

In any representation a matrix C exist, such that

$$C\gamma_\mu^T C^{-1} = -\gamma_\mu, \quad (4.54)$$

e.g.

$$(C)_{S.R.} = i\gamma^2\gamma^0 = -i\rho^1 \otimes \sigma^2 = \begin{pmatrix} 0 & -i\sigma^2 \\ -i\sigma^2 & 0 \end{pmatrix} = \begin{pmatrix} 0 & -\epsilon \\ -\epsilon & 0 \end{pmatrix}, \quad (4.55)$$

$$(C)_{W.R.} = i\gamma^2\gamma^0 = -i\rho^3 \otimes \sigma^2 = \begin{pmatrix} -i\sigma^2 & 0 \\ 0 & i\sigma^2 \end{pmatrix} = \begin{pmatrix} -\epsilon & 0 \\ 0 & \epsilon \end{pmatrix}. \quad (4.56)$$

Thus we find back the Dirac equation,

$$(i\cancel{\partial} - M)\psi^c(x) = 0. \quad (4.57)$$

with the solution

$$\psi^c(x) = \eta_c C\bar{\psi}^T(x), \quad (4.58)$$

where η_c is an arbitrary (unobservable) phase, usually to be taken unity. Some properties of C are $C^{-1} = C^\dagger$ and $C^T = -C$. One has $\bar{\psi}^c = -\psi^T C^{-1}$. In S.R. (or W.R.) and all representations connected via a real (up to an overall phase) matrix S , C is real and one has $C^{-1} = C^\dagger = C^T = -C$ and $[C, \gamma_5] = 0$. The latter implies that the conjugate of a right-handed spinor, ψ_R^c , is a left-handed spinor. Explicitly, in Weyl representation we find in Dirac space

$$\psi = \begin{pmatrix} \xi \\ \eta \end{pmatrix} \xrightarrow{\mathcal{C}} \psi^c = C\bar{\psi}^T = \begin{pmatrix} -\epsilon\eta^* \\ \epsilon\xi^* \end{pmatrix}. \quad (4.59)$$

4.4 Plane wave solutions

For a free massive particle, the best representation to describe particles at rest is the standard representation, in which γ^0 is diagonal (see discussion of negative energy states in section 4.1). The explicit Dirac equation in the standard representation reads

$$\begin{pmatrix} i\frac{\partial}{\partial t} - M & i\boldsymbol{\sigma} \cdot \boldsymbol{\nabla} \\ -i\boldsymbol{\sigma} \cdot \boldsymbol{\nabla} & -i\frac{\partial}{\partial t} - M \end{pmatrix} \psi(x) = 0. \quad (4.60)$$

Looking for positive energy solutions $\propto \exp(-iEt)$ one finds two solutions, $\psi(x) = u^{\pm s}(p) e^{-i p \cdot x}$, with $E = E_p = +\sqrt{\mathbf{p}^2 + M^2}$, where u satisfies

$$\begin{pmatrix} E_p - M & -\boldsymbol{\sigma} \cdot \mathbf{p} \\ \boldsymbol{\sigma} \cdot \mathbf{p} & -(E_p + M) \end{pmatrix} u(p) = 0, \quad \Leftrightarrow \quad (\not{p} - M) u(p) = 0. \quad (4.61)$$

There are also two negative energy solutions, $\psi(x) = v^{\pm s}(p) e^{i p \cdot x}$, where v satisfies

$$\begin{pmatrix} -(E_p + M) & \boldsymbol{\sigma} \cdot \mathbf{p} \\ -\boldsymbol{\sigma} \cdot \mathbf{p} & (E_p - M) \end{pmatrix} v(p) = 0, \quad \Leftrightarrow \quad (\not{p} + M) v(p) = 0. \quad (4.62)$$

Explicit solutions in the standard representation are

$$u(p, s) = \sqrt{E_p + M} \begin{pmatrix} \chi_s \\ \frac{\boldsymbol{\sigma} \cdot \mathbf{p}}{E_p + M} \chi_s \end{pmatrix}, \quad v(p, s) = \sqrt{E_p + M} \begin{pmatrix} \frac{\boldsymbol{\sigma} \cdot \mathbf{p}}{E_p + M} \bar{\chi}_s \\ \bar{\chi}_s \end{pmatrix}, \quad (4.63)$$

where χ_s are two independent ($s = \pm$) two-component spinors.. Note that the spinors in the negative energy modes (antiparticles) could be two different spinors. Choosing $\bar{\chi} = -\epsilon \chi^*$ (the equivalent spin 1/2 conjugate representation), the spinors satisfy $C \bar{u}^T(p, s) = v(p, s)$ and $C \bar{v}^T(p, s) = u(p, s)$. The solutions are normalized to

$$\bar{u}(p, s) u(p, s') = 2M \delta_{ss'}, \quad \bar{v}(p, s) v(p, s') = -2M \delta_{ss'}, \quad (4.64)$$

$$\bar{u}(p, s) v(p, s') = \bar{v}(p, s) u(p, s') = 0, \quad (4.65)$$

$$u^\dagger(p, s) u(p, s') = v^\dagger(p, s) v(p, s') = 2E_p \delta_{ss'}. \quad (4.66)$$

An arbitrary spin 1/2 field can be expanded in the independent solutions. After separating positive and negative energy solutions as done in the case of the scalar field one has

$$\psi(x) = \sum_s \int \frac{d^3 k}{(2\pi)^3 2E_k} (u(k, s) e^{-i k \cdot x} b(\mathbf{k}, s) + v(k, s) e^{i k \cdot x} d^*(\mathbf{k}, s)). \quad (4.67)$$

It is straightforward to find projection operators for the positive and negative energy states

$$P_+ = \sum_s \frac{u(p, s) \bar{u}(p, s)}{2M} = \frac{\not{p} + M}{2M}, \quad (4.68)$$

$$P_- = -\sum_s \frac{v(p, s) \bar{v}(p, s)}{2M} = \frac{-\not{p} + M}{2M}. \quad (4.69)$$

In order to project out spin states, the spin polarization vector in the rest frame is the starting point. In that frame is a spacelike unit vector $s^\mu = (0, \hat{\mathbf{s}})$. In an arbitrary frame one has $s \cdot p = 0$ and $s(p)$ can e.g. be obtained by a Lorentz transformation. It is easy to check that

$$P_s = \frac{1 \pm \gamma_5 \not{s}}{2} = \frac{1}{2} \begin{pmatrix} 1 \pm \boldsymbol{\sigma} \cdot \hat{\mathbf{s}} & 0 \\ 0 & 1 \mp \boldsymbol{\sigma} \cdot \hat{\mathbf{s}} \end{pmatrix} \quad (4.70)$$

(the last equality in the restframe and in standard representation) projects out spin $\pm 1/2$ states (check this in the restframe for $\hat{\mathbf{s}} = \hat{\mathbf{z}}$).

Note that for solutions of the massless Dirac equation $\not{p}\psi = 0$. Therefore, $\gamma_5\not{p}\psi = 0$ but also $\not{p}\gamma_5\psi = -\gamma_5\not{p}\psi = 0$. This means that in the solution space for massless fermions the chirality states, $\psi_{R/L} \equiv P_{R/L}\psi$ are independent solutions. In principle massless fermions can be described by two-component spinors. The chirality projection operators in Eq. 4.48 replace the spin projection operators which are not defined (by lack of a rest frame).

Explicit examples of spinors are useful to illustrate spin eigenstates, helicity states, chirality, etc. For instance with the z-axis as spin quantization axis, one has in standard representation:

$$u(p, +1/2) = \frac{1}{\sqrt{E+M}} \begin{pmatrix} E+M \\ 0 \\ p^3 \\ p^1 + ip^2 \end{pmatrix}, \quad u(p, -1/2) = \frac{1}{\sqrt{E+M}} \begin{pmatrix} 0 \\ E+M \\ p^1 - ip^2 \\ -p^3 \end{pmatrix}, \quad (4.71)$$

$$v(p, +1/2) = \frac{1}{\sqrt{E+M}} \begin{pmatrix} p^1 - ip^2 \\ -p^3 \\ 0 \\ E+M \end{pmatrix}, \quad v(p, -1/2) = \frac{1}{\sqrt{E+M}} \begin{pmatrix} -p^3 \\ -p^1 - ip^2 \\ -(E+M) \\ 0 \end{pmatrix}. \quad (4.72)$$

Helicity states (\mathbf{p} along $\hat{\mathbf{z}}$) in Standard representation are:

$$u(p, \lambda = +1/2) = \begin{pmatrix} \sqrt{E+M} \\ 0 \\ \sqrt{E-M} \\ 0 \end{pmatrix}, \quad u(p, \lambda = -1/2) = \begin{pmatrix} 0 \\ \sqrt{E+M} \\ 0 \\ -\sqrt{E-M} \end{pmatrix}, \quad (4.73)$$

$$v(p, \lambda = +1/2) = \begin{pmatrix} 0 \\ -\sqrt{E-M} \\ 0 \\ \sqrt{E+M} \end{pmatrix}, \quad v(p, \lambda = -1/2) = \begin{pmatrix} -\sqrt{E-M} \\ 0 \\ -\sqrt{E+M} \\ 0 \end{pmatrix}. \quad (4.74)$$

By writing the helicity states in Weyl representation it is easy to project out righthanded (upper components) and lefthanded (lower components). One finds for the helicity states in Weyl representation:

$$u(p, \lambda = +1/2) = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{E+M} + \sqrt{E-M} \\ 0 \\ \sqrt{E+M} - \sqrt{E-M} \\ 0 \end{pmatrix}, \quad u(p, \lambda = -1/2) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ \sqrt{E+M} - \sqrt{E-M} \\ 0 \\ \sqrt{E+M} + \sqrt{E-M} \end{pmatrix},$$

$$v(p, \lambda = +1/2) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ \sqrt{E+M} - \sqrt{E-M} \\ 0 \\ -\sqrt{E+M} - \sqrt{E-M} \end{pmatrix}, \quad v(p, \lambda = -1/2) = \frac{1}{\sqrt{2}} \begin{pmatrix} -\sqrt{E+M} - \sqrt{E-M} \\ 0 \\ +\sqrt{E+M} - \sqrt{E-M} \\ 0 \end{pmatrix}.$$

Also for helicity states $C \bar{u}^T(p, \lambda) = v(p, \lambda)$ and $C \bar{v}^T(p, \lambda) = u(p, \lambda)$. We note that for high energy the positive helicity ($\lambda = +1/2$) is in essence righthanded, while the negative helicity ($\lambda = -1/2$) is in essence lefthanded. The ratio of the components is (directly or inversely) proportional to

$$\frac{\sqrt{E+M} - \sqrt{E-M}}{\sqrt{E+M} + \sqrt{E-M}} = \frac{2M}{(\sqrt{E+M} + \sqrt{E-M})^2} \xrightarrow{M \ll E} \frac{M}{2E},$$

which vanishes in the ultra-relativistic limit $E \gg M$ or in the massless case.

4.5 γ gymnastics and applications

An overview of properties can be found in many text books. Some elementary properties are:

Properties of products of γ -matrices:

- (1) $\gamma^\mu \gamma_\mu = 4$
- (2) $\gamma^\mu \gamma^\nu \gamma_\mu = -2\gamma^\nu$ or $\gamma^\mu \not{a} \gamma_\mu = -2 \not{a}$.
- (3) $\gamma^\mu \gamma^\rho \gamma^\sigma \gamma_\mu = 4 g^{\rho\sigma}$ or $\gamma^\mu \not{a} \not{b} \gamma_\mu = 4 a \cdot b$.
- (4) $\gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma \gamma_\mu = -2\gamma^\sigma \gamma^\rho \gamma^\nu$.
- (5) $\gamma^\mu \gamma^\lambda \gamma^\nu \gamma^\rho \gamma^\sigma \gamma_\mu = 2 [\gamma^\sigma \gamma^\lambda \gamma^\nu \gamma^\rho + \gamma^\rho \gamma^\nu \gamma^\lambda \gamma^\sigma]$.
- (6) $\gamma_5 = i \gamma^0 \gamma^1 \gamma^2 \gamma^3 = -i \epsilon_{\mu\nu\rho\sigma} \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma / 4!$.
- (7) $\gamma_\mu \gamma_5 = -\gamma_5 \gamma_\mu = i \epsilon_{\mu\nu\rho\sigma} \gamma^\nu \gamma^\rho \gamma^\sigma / 3!$.
- (8) $\sigma_{\mu\nu} \equiv (i/2) [\gamma_\mu, \gamma_\nu]$
- (9) $\sigma_{\mu\nu} \gamma_5 = \gamma_5 \sigma_{\mu\nu} = \epsilon_{\mu\nu\rho\sigma} \gamma^\rho \gamma^\sigma / 2$.
- (10) $\gamma^\mu \gamma^\nu = g^{\mu\nu} - i \sigma^{\mu\nu}$
- (11) $\gamma^\mu \gamma^\nu \gamma^\rho = S^{\mu\nu\rho\sigma} \gamma_\sigma + i \epsilon^{\mu\nu\rho\sigma} \gamma_\sigma \gamma_5$ with $S^{\mu\nu\rho\sigma} = (g^{\mu\nu} g^{\rho\sigma} - g^{\mu\rho} g^{\nu\sigma} + g^{\mu\sigma} g^{\nu\rho})$.

Properties of traces of γ -matrices

- (1) $\text{Tr}(\gamma^{\mu_1} \dots \gamma^{\mu_n}) = 0$ if n is odd.
(Use $(\gamma_5)^2 = 1$ and pull one γ_5 through.)
- (2) $\text{Tr}(1) = 4$.
- (3) $\text{Tr}(\gamma^\mu \gamma^\nu) = 4 g^{\mu\nu}$ or $\text{Tr}(\not{a} \not{b}) = 4 a \cdot b$.
- (4) $\text{Tr}(\gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma) = 4 S^{\mu\nu\rho\sigma}$.
- (5) $\text{Tr}(\gamma^{\mu_1} \dots \gamma^{\mu_n}) = g^{\mu_1 \mu_2} \text{Tr}(\gamma^{\mu_3} \dots \gamma^{\mu_n}) - g^{\mu_1 \mu_3} \text{Tr}(\gamma^{\mu_2} \gamma^{\mu_4} \dots \gamma^{\mu_n}) + \dots$
- (6) $\text{Tr}(\gamma_5) = 0$.
- (7) $\text{Tr}(\gamma_5 \gamma^\mu \gamma^\nu) = 0$ or $\text{Tr}(\gamma_5 \not{a} \not{b}) = 0$.
- (8) $\text{Tr}(\gamma_5 \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma) = -4i \epsilon^{\mu\nu\rho\sigma}$.
- (9) $\text{Tr}(\gamma^{\mu_1} \dots \gamma^{\mu_{2n}}) = \text{Tr}(\gamma^{\mu_{2n}} \dots \gamma^{\mu_1})$.

(Use matrices C to transpose the matrices in the expression under the trace and use $\text{Tr} A^T = \text{Tr} A$.)

In order to show the use of the above relations, we will give one example, namely the calculation of the quantity

$$L^{\mu\nu}(k, k') = \frac{1}{2} \sum_{s, s'} (\bar{u}(k, s) \gamma^\mu u(k', s'))^* (\bar{u}(k, s) \gamma^\nu u(k', s')), \quad (4.75)$$

which appears in quantum electrodynamics calculations (see Chapter 10) for the emission of a photon from an electron changing its momentum from k to k' ($k^2 = k'^2 = m^2$). In principle one can take a representation and just calculate the quantity $\bar{u}(k, s) \gamma^\mu u(k', s')$, etc. It is, however, more convenient to use the projection operators introduced earlier. For this we first have to prove (do this) that

$$(\bar{u}(k, s) \gamma^\mu u(k', s'))^* = \bar{u}(k', s') \gamma^\mu u(k, s), \quad (4.76)$$

This leads with explicit Dirac indices to

$$\begin{aligned} L^{\mu\nu} &= \frac{1}{2} \sum_{s, s'} \bar{u}_i(k', s') (\gamma^\mu)_{ij} u_j(k, s) \bar{u}_k(k, s) (\gamma^\nu)_{kl} u_l(k', s') \\ &= \frac{1}{2} \sum_{s, s'} u_l(k', s') \bar{u}_i(k', s') (\gamma^\mu)_{ij} u_j(k, s) \bar{u}_k(k, s) (\gamma^\nu)_{kl} \\ &= \frac{1}{2} (\not{k}' + m)_{li} (\gamma^\mu)_{ij} (\not{k} + m)_{jk} (\gamma^\nu)_{kl} \\ &= \frac{1}{2} \text{Tr} [(\not{k}' + m) \gamma^\mu (\not{k} + m) \gamma^\nu]. \end{aligned}$$

The trace is linear and can be split up in parts containing traces with up to four gamma-matrices, of which only the traces of four and two gamma-matrices are nonzero. They lead to

$$\begin{aligned}
L^{\mu\nu} &= \frac{1}{2} \text{Tr} [(k' + m)\gamma^\mu(k + m)\gamma^\nu] \\
&= 2 [k^\mu k'^\nu + k^\nu k'^\mu - g^{\mu\nu}(k \cdot k' - m^2)] \\
&= 2 k^\mu k'^\nu + 2 k^\nu k'^\mu + (k - k')^2 g^{\mu\nu}.
\end{aligned} \tag{4.77}$$

Exercises

Exercise 4.1

Prove Eq. 4.13,

$$H(\mathbf{p}) = \exp\left(\frac{\boldsymbol{\eta} \cdot \boldsymbol{\sigma}}{2}\right) = \frac{M + E + \boldsymbol{\sigma} \cdot \mathbf{p}}{\sqrt{2M(E + M)}},$$

where $E = M\gamma = M \cosh(\eta)$ and $\mathbf{p} = M\beta\gamma\hat{\mathbf{n}} = M \sinh(\eta) \hat{\mathbf{n}}$. Do this e.g. by proving that

$$H^2(\mathbf{p}) = \exp(\boldsymbol{\eta} \cdot \boldsymbol{\sigma}) = (E + \boldsymbol{\sigma} \cdot \mathbf{p})/M = \tilde{\sigma}^\mu p_\mu/M.$$

Exercise 4.2

Show that $j^\mu = \bar{\psi}\gamma^\mu\psi$ is a conserved current if $\bar{\psi}$ en ψ are solutions of the Dirac equation

Exercise 4.3

Imposing Lorentz invariance on the Dirac equation can be achieved by requiring $L^{-1}\gamma^\mu L = \Lambda_\nu^\mu\gamma^\nu$. Look at the infinitesimal forms $\Lambda_\nu^\mu = g_\nu^\mu + \omega_\nu^\mu$ and $L = 1 - (i/2)\omega_{\rho\sigma}S^{\rho\sigma}$ to obtain

$$\begin{aligned}
[\gamma^\mu, iS^{\rho\sigma}] &= g^{\rho\mu}\gamma^\sigma - g^{\sigma\mu}\gamma^\rho, \\
-iS^{\rho\sigma} &= -\frac{1}{4}[\gamma^\rho, \gamma^\sigma] = \frac{i}{2}\sigma^{\rho\sigma}.
\end{aligned}$$

Exercise 4.4

Show that in

$$P_+ \equiv \sum_{s=\pm 1/2} \frac{u(p, s)\bar{u}(p, s)}{2M} = \frac{\not{p} + M}{2M}$$

both sides are projection operators. Show that the equality holds, e.g. by letting them act on the four basis spinors $u(p, s)$ and $v(p, s)$.

Exercise 4.5

Show starting from the relation $\{\gamma_\mu, \gamma_\nu\} = 2g_{\mu\nu}$ that

$$\begin{aligned}
\gamma_\mu \not{a} \gamma^\mu &= -2 \not{a}, \\
\text{Tr}(\not{a}\not{b}) &= 4 a \cdot b, \\
\text{Tr}(\gamma^{\mu_1} \dots \gamma^{\mu_n}) &= 0 \quad \text{if } n \text{ is odd.}
\end{aligned}$$

Exercise 4.6

Apply space-inversion, $x \rightarrow \tilde{x}$, to the Dirac equation and use this to show that the spinor $\psi^p(x) = \gamma_0\psi(\tilde{x})$, where $\tilde{x} = (t, -\mathbf{x})$ is also a solution of the Dirac equation.

Exercise 4.7

- (a) Show that (in standard representation)

$$\psi(\mathbf{x}, t) = N \begin{pmatrix} j_0(kr)\chi_m \\ i \frac{k}{E+M} \boldsymbol{\sigma} \cdot \hat{\mathbf{r}} j_1(kr)\chi_m \end{pmatrix} e^{-iEt},$$

is a stationary solution of the (free) Dirac equation. In this solution the wave number $k = |\mathbf{k}| = \sqrt{E^2 - M^2}$ and χ_m a two-component spinor while j_0 and j_1 are spherical Bessel functions.

Note: use first the appropriate equation (which?) that is obeyed by each individual component of the Dirac equation to show that the upper two components are allowed solutions.

- (b) Next, we want to confine this solution to a spherical cavity with radius
- R
- . A condition for confining the fermion in the cavity is
- $i\not{r}\psi = \psi$
- . Show that this condition is sufficient to guarantee that
- $n_\mu j^\mu = 0$
- , i.e. there is no current flowing through the surface of the cavity.

Note: To prove this, you need to determine the corresponding confinement condition for $\bar{\psi}$. The conditions for ψ and $\bar{\psi}$ imply that $n_\mu j^\mu = \bar{\psi}\not{r}\psi = 0$.

It turns out that there is actually a whole family of conditions that provide confinement, namely $i\not{r}\psi = e^{i\alpha\gamma_5}\psi$. We will not pursue this chiral freedom at this point and take $\alpha = 0$.

- (c) Apply the confinement condition to find the value of
- k
- for the lowest eigenmode in the spherical cavity. For such a cavity
- $n^\mu = (0, \hat{\mathbf{r}})$
- . Calculate the value of
- k
- for
- $MR = 0$
- ,
- $MR = 1.5$
- and
- $M \rightarrow \infty$
- .

- (d) Plot for these cases the vector densities
- $\psi^\dagger\psi$
- and the scalar densities
- $\bar{\psi}\psi$
- as a function of
- r/R
- .

Exercise 4.8

A negatively charged pion (π^-) decays in a lepton ℓ (ℓ can be an electron or a muon) and a corresponding anti-neutrino ($\bar{\nu}_\ell$). The pion has spin 0, the lepton and neutrino have spin 1/2 and are described by momentum-space spinors. Our starting point is the transition amplitude given by

$$\mathcal{M} = \langle \ell(k_\ell)\bar{\nu}_\ell(k_\nu) | H_{\text{int}} | \pi(p) \rangle = G_F f_\pi \bar{u}(k_\ell)\not{p}(1 - \gamma_5)v(k_\nu).$$

Calculate $|\mathcal{M}|^2$, which is the most important factor needed to find the decay width (inverse lifetime). Note: the outcome shows that the decay into muons is much more likely than the decay into electrons ($m_\mu = 105.6$ MeV, $m_e = 0.511$ MeV). Actually if the lepton would be massless, the decay would even be forbidden. Do you understand why? The calculation of the life-time is completed in chapter 11.

Chapter 5

Vector fields and Maxwell equations

5.1 Fields for spin 1

In section 3.2 we have discussed the vectors ϵ_λ (with $\lambda = 0$ and ± 1) as basis states of the spin 1 representation. In the rest frame plane waves of the form $\epsilon_\lambda e^{ik \cdot x}$, thus, can describe spin 1 waves. The covariant generalization is to describe them with

$$V_\mu(x) = \epsilon_\mu(k) e^{\pm ik \cdot x}. \quad (5.1)$$

These are solutions of the Klein-Gordon equation with in addition a condition,

$$(\square + M^2)V_\mu = 0, \quad \text{with} \quad \partial_\mu V^\mu = 0 \text{ (Lorentz condition)}. \quad (5.2)$$

The Lorentz condition implies that $k^\mu \epsilon_\mu(k) = 0$. In the restframe, where $k = k_0 = (M, \mathbf{0})$, there are then indeed three independent possibilities, namely $\epsilon^\mu(k_0) = (0, \epsilon_\lambda)$. The vectorfield is therefore suited to describe spin 1. In an arbitrary frame $\epsilon_\mu^{(\lambda)}(k)$ with $\lambda = 0, \pm(1)$ are obtained by boosting the restframe vectors. A real vector field $V_\mu(x)$ then can be expanded in modes as

$$V_\mu(x) = \sum_{\lambda=0,\pm} \int \frac{d^3k}{(2\pi)^3 2E} \left(\epsilon_\mu^{(\lambda)}(k) e^{-ik \cdot x} c(\mathbf{k}, \lambda) + \epsilon_\mu^{(\lambda)*}(k) e^{ik \cdot x} c^*(\mathbf{k}, \lambda) \right). \quad (5.3)$$

Since the vectors $\epsilon_\mu^{(\lambda)}(k)$ together with the momentum k^μ form a complete set of four-vectors, one has

$$\sum_{\lambda=0,\pm} \epsilon_\mu^{(\lambda)}(k) \epsilon_\nu^{(\lambda)*}(k) = -g_{\mu\nu} + \frac{k_\mu k_\nu}{M^2}. \quad (5.4)$$

5.2 The electromagnetic field

With the electromagnetic field tensor

$$F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu \quad (5.5)$$

$$= \begin{pmatrix} 0 & -E^1 & -E^2 & -E^3 \\ E^1 & 0 & -B^3 & B^2 \\ E^2 & B^3 & 0 & -B^1 \\ E^3 & -B^2 & B^1 & 0 \end{pmatrix}, \quad (5.6)$$

where A^μ is the four-vector potential

$$A^\mu = (\phi, \mathbf{A}), \quad (5.7)$$

and the (conserved) current $j^\mu = (\rho, \mathbf{j})$ the Maxwell equations read

$$\partial_\mu F^{\mu\nu} = j^\nu. \quad (5.8)$$

Furthermore, the antisymmetry of $F^{\mu\nu}$ implies that

$$\partial_\mu \tilde{F}^{\mu\nu} = 0 \quad \text{or} \quad \partial_\lambda F_{\mu\nu} + \partial_\mu F_{\nu\lambda} + \partial_\nu F_{\lambda\mu} = 0, \quad (5.9)$$

where $\tilde{F}^{\mu\nu} \equiv -\frac{1}{2}\epsilon^{\mu\nu\rho\sigma}F_{\rho\sigma}$ is the tensor where $\mathbf{E} \rightarrow \mathbf{B}$ and $\mathbf{B} \rightarrow -\mathbf{E}$. It states the absence of magnetic charges and Faraday's law.

For electrodynamics one has furthermore the freedom of gauge transformations. Under a gauge transformation

$$A_\mu \longrightarrow A_\mu + \partial_\mu \chi, \quad (5.10)$$

the electric and magnetic fields are unchanged,

$$F_{\mu\nu} \longrightarrow F_{\mu\nu} + (\partial_\mu \partial_\nu \chi - \partial_\nu \partial_\mu \chi) = F_{\mu\nu}. \quad (5.11)$$

The equations of motion for the fields A_μ become

$$\square A_\mu - \partial_\mu (\partial_\nu A^\nu) = j_\mu. \quad (5.12)$$

This equation is not affected by a gauge transformation. The gauge freedom can first of all be used to impose the Lorentz condition for the electromagnetic field,

$$\partial_\mu A^\mu = 0, \quad (5.13)$$

in which case one has

$$\square A_\mu = j_\mu, \quad (5.14)$$

of which the solutions give the Liénard-Wiechert potentials. The equation in vacuum, $\square A_\mu = 0$, moreover, shows that the electromagnetic fields correspond to the case of a massless spin 1 field. Although the Lorenz condition is a constraint, it does not eliminate the freedom of gauge transformations but they are now restricted to $A_\mu \rightarrow A_\mu + \partial_\mu \chi$ with $\square \chi = 0$.

The Lorenz condition can actually be incorporated by changing the equations of motion to

$$\square A_\mu + (\lambda - 1)\partial_\mu (\partial_\nu A^\nu) = j_\mu. \quad (5.15)$$

Taking the divergence of this equation, one finds for a conserved current and $\lambda \neq 0$

$$\square \partial_\mu A^\mu = 0. \quad (5.16)$$

Hence, if $\partial_\mu A^\mu = 0$ for large times $|t|$, it will vanish at all times. Note that the incorporation of the Lorenz condition can also be done for the case of the massive vector field, i.e. if in Eq. 5.15 the d'Alembertian is replaced by $\square + M^2$.

As remarked above, for $M = 0$, there is still a gauge freedom left ($A_\mu \rightarrow A_\mu + \partial_\mu \chi$ with $\square \chi = 0$). One possible choice is the gauge


$$A_0 = 0 \quad (5.17)$$

(radiation, transverse or Coulomb gauge). In that case one also has $\nabla \cdot \mathbf{A} = 0$ or $k_i \epsilon_i(k) = 0$. Therefore if $k^\mu = (|\mathbf{k}|, 0, 0, |\mathbf{k}|)$, only two independent vectors remain

$$\epsilon^{(\pm)\mu}(k) = (0, \mp 1, -i, 0), \quad (5.18)$$

corresponding to the helicity states $\lambda = \pm 1$ of the massless photon field, in accordance with what we found in section 3.5.

Note that via parity transformation one obtains another solution, $A_\mu^P(x) = \tilde{A}_\mu(\tilde{x})$, while the fact that the Minkowski space is real implies that $A_\mu^C(x) = A_\mu^*(x) = A_\mu(x)$.


 Figure 5.1: *The Aharonov-Bohm experiment*

5.3 The electromagnetic field and topology*

The electric and magnetic fields \mathbf{E} and \mathbf{B} as appearing for instance in the Lorentz force on a moving charge,

$$\mathbf{F} = e\mathbf{E} + e\mathbf{v} \times \mathbf{B} \quad (5.19)$$

are gauge invariant, while the electromagnetic field A_μ is not. Nevertheless the significance of A_μ is shown in the Aharonov-Bohm experiment. In this experiment an observable phase difference is measured for electrons moving through field-free space ($\mathbf{E} = \mathbf{B} = 0$ but $A_\mu \neq 0$), illustrated in fig. 5.1. The phase difference between the electrons travelling two different paths is given by

$$\delta = 2\pi \frac{a}{\lambda} = \frac{a}{\lambda} \approx \frac{\Delta x}{L} \frac{d}{\lambda},$$

or $\Delta x = (L\lambda/d)\delta$. This causes an interference pattern as a function of Δx . In the presence of an electromagnetic field A_μ an additional phase difference is observed. This occurs in the region outside the solenoid, where $A_\mu \neq 0$, but where $\mathbf{E} = \mathbf{B} = 0$. To be precise,

$$\begin{aligned} \text{Inside solenoid:} \quad & \mathbf{B} = B\hat{z}, \quad \mathbf{A} = \frac{Br}{2}\hat{\varphi} = \left(-\frac{By}{2}, \frac{Bx}{2}, 0\right), \\ \text{Outside solenoid:} \quad & \mathbf{B} = 0, \quad \mathbf{A} = \frac{BR^2}{2r}\hat{\varphi} = \left(-\frac{BR^2y}{2r^2}, \frac{BR^2x}{2r^2}, 0\right), \end{aligned}$$

where $\hat{r} = (\cos \varphi, \sin \varphi, 0)$ and $\hat{\varphi} = (-\sin \varphi, \cos \varphi, 0)$. The explanation of the significance of the vector potential \mathbf{A} for the phase of the electron wave function is found in minimal substitution

$$\begin{aligned} \psi \propto \exp\left(\frac{i}{\hbar}\mathbf{p} \cdot \mathbf{r}\right) & \longrightarrow \exp\left(\frac{i}{\hbar}(\mathbf{p} - e\mathbf{A}) \cdot \mathbf{r}\right) \\ & = \exp\left(-i\frac{e}{\hbar}\mathbf{A} \cdot \mathbf{r}\right) \exp\left(\frac{i}{\hbar}\mathbf{p} \cdot \mathbf{r}\right). \end{aligned}$$

The phase difference between the two paths of the electron then are given by

$$\begin{aligned} \Delta\delta &= -\frac{e}{\hbar} \int_1 \mathbf{A} \cdot d\mathbf{r} + \frac{e}{\hbar} \int_2 \mathbf{A} \cdot d\mathbf{r} = \frac{e}{\hbar} \oint \mathbf{A} \cdot d\mathbf{r} \\ &= \frac{e}{\hbar} \int (\nabla \times \mathbf{A}) \cdot d\mathbf{s} = \frac{e}{\hbar} \int \mathbf{B} \cdot d\mathbf{s} = \frac{e}{\hbar} \Phi, \end{aligned} \quad (5.20)$$

where Φ is the flux through the solenoid. This phase difference is actually observed.

The situation, nevertheless, may look a bit awkward. It is however, nothing more than a manifestation of a nontrivial vacuum (a la the Dirac sea for fermions). The energy density $\frac{1}{2}(\mathbf{E}^2 + \mathbf{B}^2) = 0$ (hence a vacuum), but $\mathbf{A} \neq 0$ (hence there is structure in the vacuum).




Figure 5.2: *The topology of the space in the Aharonov-Bohm experiment, illustrating also that the group structure of the mapping of $U(1)$ into this space is that of the group of integers*

Indeed when one considers the \mathbf{A} -field in the example of the Aharonov-Bohm effect outside the solenoid, it can be obtained from the situation $\mathbf{A} = 0$ by a gauge transformation,

$$\mathbf{A} = \nabla\chi \quad \text{with} \quad \chi = \frac{BR^2}{2}\varphi \quad (5.21)$$

(φ being the azimuthal angle). This situation that \mathbf{A} can be written as $\nabla\chi$ is always true when $\nabla \times \mathbf{A} = 0$. The observable phase going around in general is

$$\Delta\delta = \frac{e}{\hbar} \oint \mathbf{A} \cdot d\mathbf{r} = \frac{e}{\hbar} \oint \nabla\chi \cdot d\mathbf{r} = \frac{e}{\hbar} \chi|_{\varphi=0}^{\varphi=2\pi}.$$

The function χ in Eq. 5.21, however, is multivalued as $\varphi = 0$ and $\varphi = 2\pi$ are the same point in space. If this would be happening in empty space one would be in trouble. By going around an arbitrary loop a different number of times the electron wave function would acquire different phases or it would acquire an arbitrary phase in a point by contracting a loop continuously into that point. Therefore the gauge transformation must be uniquely defined in the space one is working in, implying that it must be single-valued¹.

Now back to the Aharonov-Bohm experiment. The difference here is that we are working in a space with a 'defect' (the infinitely long solenoid). In such a space loops with different winding number (i.e. the number of times they go around the defect) cannot be continuously deformed into one another. Therefore space outside the solenoid doesn't care that χ is multi-valued. The Aharonov-Bohm experiment shows a realization of this possibility.

Another situation in which the topology of space can be used is in the case of superconductors. Consider a superconductor, forming a simple (connected) space. Below the critical temperature the magnetic field is squeezed out of the superconducting material, organizing itself in tiny flux tubes (Abrikosov strings), minimizing the space occupied by \mathbf{B} fields. The only way for flux tubes to be formed and move around without global consequences is when each flux tube contains a flux Φ such that $\Delta\delta = (e/\hbar)\Phi = n \cdot 2\pi$, giving no observable phase².

¹The occurrence of nontrivial possibilities, i.e. nonobservable phases $\varphi = 2\pi n$, has been employed by Dirac in constructing magnetic monopoles in electrodynamics

²Actually, the relevant charge turns out to be $2e$ because the electrons appear in Cooper pairs

Exercises

Exercise 5.1

Show that Eqs. 5.8 and 5.9 explicitly give the equations

$$\begin{aligned}\nabla \cdot \mathbf{E} &= \rho, \\ \nabla \times \mathbf{B} &= \mathbf{j} + \frac{\partial \mathbf{E}}{\partial t}, \\ \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} &= 0, \\ \nabla \cdot \mathbf{B} &= 0.\end{aligned}$$

Chapter 6

Classical lagrangian field theory

6.1 Euler-Lagrange equations

In the previous chapter we have seen the equations for scalar fields (Klein-Gordon equation), Dirac fields (Dirac equation) and massless vector fields (Maxwell equations) and corresponding to these fields conserved currents describing the probability and probability current. These equations can be obtained from a lagrangian using the action principle.

As an example, recall classical mechanics with the action

$$S = \int_{t_1}^{t_2} dt L(x, \dot{x}) \quad (6.1)$$

and as an example the lagrangian

$$L(x, \dot{x}) = K - V = \frac{1}{2}m\dot{x}^2 - V(x). \quad (6.2)$$

The principle of minimal action looks for a stationary action under variations in the coordinates and time, thus

$$t' = t + \delta\tau, \quad (6.3)$$

$$x'(t) = x(t) + \delta x(t), \quad (6.4)$$

and the total change

$$x'(t') = x(t) + \Delta x(t), \quad (6.5)$$

with $\Delta x(t) = \delta x(t) + \dot{x}(t) \delta\tau$. The requirement $\delta S = 0$ with fixed boundaries $x(t_1) = x_1$ and $x(t_2) = x_2$ leads to

$$\begin{aligned} \delta S &= \int_{t_1}^{t_2} dt \left\{ \frac{\partial L}{\partial x} \delta x + \frac{\partial L}{\partial \dot{x}} \delta \dot{x} \right\} + L \delta\tau \Big|_{t_1}^{t_2} \\ &= \int_{t_1}^{t_2} dt \left\{ \frac{\partial L}{\partial x} - \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{x}} \right) \right\} \delta x + \left(\frac{\partial L}{\partial \dot{x}} \delta x + L \delta\tau \right) \Big|_{t_1}^{t_2}. \end{aligned} \quad (6.6)$$

The first term leads to the Euler-Lagrange equations,

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{x}} \right) = \frac{\partial L}{\partial x}. \quad (6.7)$$

The quantity $\partial L / \partial \dot{x}$ plays a special role and is known as the *canonical momentum*,

$$p = \frac{\partial L}{\partial \dot{x}}. \quad (6.8)$$

For the lagrangian specified above this leads directly to Newton's law $\dot{p} = m\ddot{x} = -\partial V/\partial x$.

The second term in Eq. 6.6 can be rewritten as

$$\delta S = \dots + \left(\frac{\partial L}{\partial \dot{x}} \Delta x - H \delta \tau \right) \Big|_{t_1}^{t_2}, \quad (6.9)$$

which is done because the first term (multiplying Δx , which in classical mechanics vanishes at the boundary) does not play a role. The *hamiltonian* H is defined by

$$H(p, x) \equiv p \dot{x} - L. \quad (6.10)$$

One sees that invariance under time translations requires that $H(t_1) = H(t_2)$, i.e. H is a conserved quantity. In general any continuous transformation, $\Delta x = (dx/d\lambda)_{\lambda=0} \delta\lambda$ and $\delta t = (dt/d\lambda)_{\lambda=0} \delta\lambda$, gives rise to a conserved quantity (at least if H is invariant!)

$$Q = p \left(\frac{dx}{d\lambda} \right)_{\lambda=0} - H \left(\frac{dt}{d\lambda} \right)_{\lambda=0}, \quad (6.11)$$

of which p and H are the simplest examples related to space and time translations, but which in 3 dimensions includes conserved quantities related to rotations and boosts.

In classical field theory one proceeds in complete analogy but using functions depending on space and time (classical fields, think for instance of a temperature or density distribution or of an electromagnetic field). Consider a lagrangian density \mathcal{L} which depends on these functions, their derivatives and possibly on the position, $\mathcal{L}(\phi(x), \partial_\mu \phi(x), x)$ and an action

$$S[\phi] = \int_{t_1}^{t_2} dt L = \int dt d^3x \mathcal{L}(\phi(x), \partial_\mu \phi(x)) = \int_R d^4x \mathcal{L}(\phi(x), \partial_\mu \phi(x)). \quad (6.12)$$

Here R indicates a space-time volume bounded by (R^3, t_1) and (R^3, t_2) , also indicated by ∂R (a more general volume in four-dimensional space-time with some boundary ∂R can also be considered). Variations in the action can come from the coordinates or the fields, indicated as

$$x'^\mu = x^\mu + \delta x^\mu, \quad (6.13)$$

$$\phi'(x) = \phi(x) + \delta \phi(x) \quad (6.14)$$

or combined

$$\phi'(x') = \phi(x) + \Delta \phi(x), \quad (6.15)$$

with $\Delta \phi(x) = \delta \phi(x) + (\partial_\mu \phi) \delta x^\mu$. The resulting variation of the action is

$$\delta S = \int_R d^4x' \mathcal{L}(\phi', \partial_\mu \phi', x') - \int_R d^4x \mathcal{L}(\phi, \partial_\mu \phi, x). \quad (6.16)$$

The change in variables $x \rightarrow x'$ in the integration volume involves a surface variation of the form

$$\int_{\partial R} d\sigma_\mu \mathcal{L} \delta x^\mu.$$

Note for the specific choice of the surface for constant times t_1 and t_2 ,

$$\int_{\partial R} d\sigma_\mu \dots = \int_{(R^3, t_2)} d^3x \dots - \int_{(R^3, t_1)} d^3x \dots \quad (6.17)$$

Furthermore the variations $\delta \phi$ and $\delta \partial_\mu \phi$ contribute to δS , giving¹

$$\begin{aligned} \delta S &= \int_R d^4x \left[\frac{\delta \mathcal{L}}{\delta(\partial_\mu \phi)} \delta(\partial_\mu \phi) + \frac{\delta \mathcal{L}}{\delta \phi} \delta \phi \right] + \int_{\partial R} d\sigma_\mu \mathcal{L} \delta x^\mu \\ &= \int_R d^4x \left[\frac{\delta \mathcal{L}}{\delta \phi} - \partial_\mu \left(\frac{\delta \mathcal{L}}{\delta(\partial_\mu \phi)} \right) \right] \delta \phi + \int_{\partial R} d\sigma_\mu \left[\frac{\delta \mathcal{L}}{\delta(\partial_\mu \phi)} \delta \phi + \mathcal{L} \delta x^\mu \right]. \end{aligned} \quad (6.18)$$

¹Taking a functional derivative, indicated with $\delta F[\phi]/\delta \phi$ should pose no problems. We will come back to it in a bit more formal way in section 9.2.

With for the situation of classical fields all variations of the fields and coordinates at the surface vanishing, the second term is irrelevant. The integrand of the first term must vanish, leading to the Euler-Lagrange equations,

$$\partial_\mu \left(\frac{\delta \mathcal{L}}{\delta(\partial_\mu \phi)} \right) = \frac{\delta \mathcal{L}}{\delta \phi}. \quad (6.19)$$

6.2 Lagrangians for spin 0, 1/2 and 1 fields

By an appropriate choice of lagrangian density the equations of motion discussed in previous chapters for the scalar field (spin 0), the Dirac field (spin 1/2) and the vector field (spin 1) can be found.

The scalar field

It is straightforward to derive the equations of motion for a real scalar field ϕ from the lagrangian densities,

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} M^2 \phi^2 \quad (6.20)$$

$$= -\frac{1}{2} \phi (\partial_\mu \partial^\mu + M^2) \phi, \quad (6.21)$$

which differ only by surface terms, leading to

$$(\square + M^2)\phi(x) = 0. \quad (6.22)$$

For the complex scalar field one conventionally uses

$$\mathcal{L} = \partial_\mu \phi^* \partial^\mu \phi - M^2 \phi^* \phi \quad (6.23)$$

$$= -\phi^* (\partial_\mu \partial^\mu + M^2) \phi, \quad (6.24)$$

which can be considered as the sum of the lagrangian densities for two real scalar fields ϕ_1 and ϕ_2 with $\phi = (\phi_1 + i\phi_2)/\sqrt{2}$. One easily obtains

$$(\square + M^2)\phi(x) = 0, \quad (6.25)$$

$$(\square + M^2)\phi^*(x) = 0. \quad (6.26)$$

The Dirac field

The appropriate lagrangian from which to derive the equations of motion is

$$\mathcal{L} = \frac{i}{2} \bar{\psi} \overleftrightarrow{\not{\partial}} \psi - M \bar{\psi} \psi = \frac{i}{2} \bar{\psi} \overrightarrow{\not{\partial}} \psi - \frac{i}{2} \bar{\psi} \overleftarrow{\not{\partial}} \psi - M \bar{\psi} \psi \quad (6.27)$$

$$= \bar{\psi} (i\overrightarrow{\not{\partial}} - M) \psi, \quad (6.28)$$

where the second line is not symmetric but in the action only differs from the symmetric version by a surface term (partial integration). Using the variations in $\bar{\psi}$ (in the symmetric form),

$$\begin{aligned} \frac{\delta \mathcal{L}}{\delta(\partial_\mu \bar{\psi})} &= -\frac{i}{2} \gamma^\mu \psi \\ \frac{\delta \mathcal{L}}{\delta \bar{\psi}} &= \frac{i}{2} \overrightarrow{\not{\partial}} \psi - M \psi, \end{aligned}$$

one obtains immediately

$$\left(i \overrightarrow{\not{\partial}} - M \right) \psi = 0, \quad (6.29)$$

and similarly from the variation with respect to ψ

$$\bar{\psi} \left(i \overleftarrow{\not{\partial}} + M \right) = 0. \quad (6.30)$$

It is often useful to link to the two-component spinors ξ and η which we started with in chapter 4, or equivalently separate the field into right- and lefthanded ones. In that case one finds trivially

$$\mathcal{L} = \frac{1}{2} \overline{\psi_R} i \overleftrightarrow{\not{\partial}} \psi_R + \frac{1}{2} \overline{\psi_L} i \overleftrightarrow{\not{\partial}} \psi_L - M (\overline{\psi_R} \psi_L + \overline{\psi_L} \psi_R), \quad (6.31)$$

showing e.g. that the lagrangian separates into two independent parts for $M = 0$.

The mass term in the Dirac lagrangian 6.31 rewritten in terms of ξ and η is

$$\mathcal{L}_M(\text{Dirac}) = -M (\xi^\dagger \eta + \eta^\dagger \xi). \quad (6.32)$$

There exists a different possibility to write down a mass term with only one kind of fields, namely²

$$\mathcal{L}_M(\text{Majorana}) = +\frac{1}{2} (M \eta^\dagger \epsilon \eta^* - M^* \eta^T \epsilon \eta). \quad (6.33)$$

Since the kinetic term in \mathcal{L} separates naturally in ξ and η , it is possible to introduce 'real' spinors or *Majorana spinors* that satisfies $\Upsilon^c = \Upsilon$. The spinor to be used satisfies

$$\Upsilon^c = \Upsilon \quad \text{and} \quad \Upsilon_L = \psi_L = \begin{pmatrix} 0 \\ \eta \end{pmatrix} \Rightarrow \Upsilon \equiv \begin{pmatrix} -\epsilon \eta^* \\ \eta \end{pmatrix}, \quad (6.34)$$

for which $\eta_0 = \epsilon \eta_0^*$. We note that

$$\psi_L^c \equiv (\psi_L)^c = C \overline{\psi_L}^T = \begin{pmatrix} -\epsilon \eta^* \\ 0 \end{pmatrix} = \Upsilon_R. \quad (6.35)$$

Rewriting the Majorana mass term in Eq. 6.33 in terms of ψ_L and ψ_L^c one finds

$$\mathcal{L} = \frac{1}{2} \overline{\psi_L} i \overleftrightarrow{\not{\partial}} \psi_L - \frac{1}{2} (M \overline{\psi_L^c} \psi_L + M^* \overline{\psi_L} \psi_L^c). \quad (6.36)$$

Note actually that using for Majorana spinors the relations $\Upsilon_R^c = \Upsilon_L$ and $\Upsilon_L^c = \Upsilon_R$, the Majorana lagrangian is actually almost of the same form as the Dirac mass term,

$$\mathcal{L} = \frac{1}{4} \overline{\Upsilon} i \overleftrightarrow{\not{\partial}} \Upsilon - \frac{1}{2} (M \overline{\Upsilon_R} \Upsilon_L + M^* \overline{\Upsilon_L} \Upsilon_R). \quad (6.37)$$

Note the factor 1/2 as compared to the Dirac lagrangian, which comes because we in essence use 'real' spinors. The Majorana case is actually more general, since a lagrangian with both Dirac and Majorana mass terms can be rewritten as the sum of two Majorana lagrangians after redefining the fields (See e.g. Peshkin and Schroeder).

Vector field

From the lagrangian density

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} M^2 V_\mu V^\mu - \frac{\lambda}{2} (\partial_\mu V^\mu)^2 \quad (6.38)$$

$$= \frac{1}{2} V^\mu [(\partial^2 + M^2) g_{\mu\nu} - (1 - \lambda) \partial_\mu \partial_\nu] V^\nu, \quad (6.39)$$

²This term is written down with η being an anticommuting Grassmann number for which

$$\begin{aligned} \alpha\beta &= -\beta\alpha, \\ (\alpha\beta)^* &= \beta^* \alpha^* = -\alpha^* \beta^*, \end{aligned}$$

and thus $(\beta^* \alpha)^* = \alpha^* \beta = -\beta \alpha^*$. The reasons for Grassmann variables will become clear in the next chapter.

where $F_{\mu\nu} = \partial_\mu V_\nu - \partial_\nu V_\mu$, one immediately obtains

$$\frac{\delta\mathcal{L}}{\delta(\partial_\mu V_\nu)} = -\partial^\mu V^\nu + \partial^\nu V^\mu - \lambda g^{\mu\nu} (\partial_\rho V^\rho) = -F^{\mu\nu} - \lambda g^{\mu\nu} (\partial_\rho V^\rho)$$

leading to the equations of motion

$$\partial_\mu F^{\mu\nu} + M^2 V^\nu + \lambda \partial^\nu (\partial \cdot V) = (\square + M^2) V^\nu - (1 - \lambda) \partial^\nu (\partial \cdot V) = 0, \quad (6.40)$$

implying

$$\partial_\mu F^{\mu\nu} + M^2 V^\nu = 0 \quad \text{and} \quad \partial_\mu V^\mu = 0. \quad (6.41)$$

6.3 Symmetries and conserved (Noether) currents

Next, we consider the second term in Eq. 6.18. In particular when the lagrangian is invariant under symmetries, one can consider the variations at the initial or final surface. These do not affect the dynamics and will give rise to conserved currents. Returning to δS the surface term is rewritten to

$$\begin{aligned} \delta S &= \int_R d^4x \dots + \int_{\partial R} d\sigma_\mu \left\{ \frac{\delta\mathcal{L}}{\delta(\partial_\mu\phi)} \Delta\phi - \left[\frac{\delta\mathcal{L}}{\delta(\partial_\mu\phi)} \partial^\nu\phi - \mathcal{L}g^{\mu\nu} \right] \delta x_\nu \right\} \\ &\equiv \int_R d^4x \dots + \int_{\partial R} d\sigma_\mu \left\{ \frac{\delta\mathcal{L}}{\delta(\partial_\mu\phi)} \Delta\phi - \Theta^{\mu\nu}(x) \delta x_\nu \right\}, \end{aligned} \quad (6.42)$$

where

$$\Theta^{\mu\nu} = \frac{\delta\mathcal{L}}{\delta(\partial_\mu\phi)} \partial^\nu\phi - \mathcal{L}g^{\mu\nu}. \quad (6.43)$$

The variation δS thus can be expressed as

$$\delta S = F(\sigma_1) - F(\sigma_2) = \int_{\partial R} d\sigma_\mu J^\mu(x) = \int_R d^4x \partial_\mu J^\mu(x), \quad (6.44)$$

with

$$J^\mu(x) = -\frac{\delta\mathcal{L}}{\delta(\partial_\mu\phi)} \Delta\phi + \Theta^{\mu\nu}(x) \delta x_\nu.$$

For continuous transformations, $\Delta\Phi = (d\phi/d\lambda)_{\lambda=0} \delta\lambda$ and $\delta x^\mu = (dx^\mu/d\lambda)_{\lambda=0} \delta\lambda$, we get

$$J^\mu(x) = -\frac{\delta\mathcal{L}}{\delta(\partial_\mu\phi)} \left(\frac{d\phi}{d\lambda} \right)_{\lambda=0} + \Theta^{\mu\nu}(x) \left(\frac{dx_\nu}{d\lambda} \right)_{\lambda=0}. \quad (6.45)$$

Thus considering $\partial R = \sigma_2 - \sigma_1$ with $\sigma = (R^3, t)$, the presence of a symmetry that leaves the lagrangian invariant, requires the presence of a conserved quantity, $Q(t_1) = Q(t_2)$, which is the space-integral over the zero-component of a conserved current, $\partial_\mu J^\mu(x) = 0$,

$$Q = \int d^3x J^0(\mathbf{x}, t), \quad (6.46)$$

For quantum fields (ϕ operator!) things become more subtle as it is not possible to specify for instance ϕ and $\dot{\phi}$ on the surface σ_1 . In this case, discussed in the next chapter, these conserved quantities become operators, which in a consistent picture precisely generate the symmetries.

As an example for the classical field, consider $U(1)$ transformations of the Dirac field proportional to the charge e ,

$$\psi(x) \rightarrow e^{ie\Lambda} \psi(x) \quad \text{or} \quad \Delta\psi(x) = ie\Lambda \psi(x). \quad (6.47)$$

From the lagrangian for the Dirac field, we obtain (since $\delta x^\mu = 0$), omitting the parameter Λ

$$\begin{aligned} j^\mu &= -\frac{\delta\mathcal{L}}{\delta(\partial_\mu\bar{\psi}^i)} \cdot (-ie\bar{\psi}^i) - \frac{\delta\mathcal{L}}{\delta(\partial_\mu\psi^i)} \cdot (ie\psi^i) \\ &= e\bar{\psi}\gamma^\mu\psi. \end{aligned} \quad (6.48)$$

For the complex scalar field the $U(1)$ transformations leave the lagrangian invariant and lead to the conserved current (see Exercise 6.3)

$$j^\mu = ie\phi^* \overleftrightarrow{\partial}_\mu \phi. \quad (6.49)$$

These currents are conserved as discussed already in chapter 2. The integral over the zero-component, $Q = \int d^3x j^0(\mathbf{x}, t)$ is the conserved charge ($\dot{Q} = 0$). In the next section we will see the quantity show up as the charge operator.

6.4 Space-time symmetries

One kind of symmetries that leave the lagrangian invariant are the Poincaré transformations, including the space-time translations and the Lorentz transformations.

Translations

Under translations (generated by P_{op}^μ) we have

$$\delta x^\mu = \epsilon^\mu, \quad (6.50)$$

$$\Delta\phi(x) = 0, \quad (6.51)$$

$$\delta\phi(x) = -(\partial_\mu\phi)\delta x^\mu = -\epsilon^\mu\partial_\mu\phi(x). \quad (6.52)$$

The behavior of the field under translations ($\delta\phi$) is governed by the translational behavior of the argument in such a way that $\Delta\phi = 0$. From Noether's theorem one sees that the current $\Theta^{\mu\nu}\epsilon_\nu$ is conserved. Therefore there are four conserved currents $\Theta^{\mu\nu}$ (ν labeling the currents!) and four conserved quantities

$$P^\nu = \int d\sigma_\mu \Theta^{\mu\nu} = \int d^3x \Theta^{0\nu}. \quad (6.53)$$

These are the energy and momentum. For quantized fields this will become the expressions of the hamiltonian and the momentum operators in terms of the fields, e.g. $H = \int d^3x \Theta^{00}(x)$.

Lorentz transformations

In this case the transformation of the coordinates and fields are written as

$$\delta x^\mu = \omega^{\mu\nu}x_\nu, \quad (\omega^{\mu\nu} \text{ antisymmetric}) \quad (6.54)$$

$$\Delta\phi^i(x) = \frac{1}{2}\omega_{\rho\sigma}(-iS^{\rho\sigma})^i_j\phi^j(x). \quad (6.55)$$

Note that the coordinate transformations can be written in a form similar to that for the fields,

$$\delta x^\mu = \frac{1}{2}\omega_{\rho\sigma}(a^{\rho\sigma})^\mu_\nu x^\nu \quad (6.56)$$

with

$$(a^{\rho\sigma})^\mu_\nu = g^{\rho\mu}g^\sigma_\nu - g^{\sigma\mu}g^\rho_\nu. \quad (6.57)$$

For the fields the behavior under Lorentz transformations has been the subject of previous chapters. Summarizing,

- **Scalar field** ϕ : $-i S_{\rho\sigma} = 0$.
- **Dirac field** ψ : $-i S_{\rho\sigma} = -(1/4)[\gamma_\rho, \gamma_\sigma]$
This result for the Dirac field has been discussed in chapter 3 (see also Exercise 4.3).
- **Vector field** A_μ : $-i S_{\rho\sigma} = a_{\rho\sigma}$.

The current deduced from Noether's theorem is

$$\begin{aligned} J^\mu &= -\frac{\delta\mathcal{L}}{\delta(\partial_\mu\phi^i)}\Delta\phi^i + \Theta^{\mu\nu}\delta x_\nu \\ &= \frac{1}{2}\omega_{\rho\sigma}\left\{-\frac{\delta\mathcal{L}}{\delta(\partial_\mu\phi^i)}(-iS^{\rho\sigma})^i_j\phi^j + \Theta^{\mu\rho}x^\sigma - \Theta^{\mu\sigma}x^\rho\right\} \end{aligned} \quad (6.58)$$

$$= \frac{1}{2}\omega_{\rho\sigma}\{H^{\mu\rho\sigma} + \Theta^{\mu\rho}x^\sigma - \Theta^{\mu\sigma}x^\rho\} \quad (6.59)$$

$$= \frac{1}{2}\omega_{\rho\sigma}M^{\mu\rho\sigma}. \quad (6.60)$$

Therefore there are six conserved currents $M^{\mu\rho\sigma}$ labeled by ρ and σ (antisymmetric) and corresponding to it there exist conserved quantities

$$M^{\rho\sigma} = \int d^3x M^{0\rho\sigma}. \quad (6.61)$$

A final note concerns the symmetry properties of $\Theta^{\mu\nu}$. In general this tensor is not symmetric. In some applications (specifically coupling to gravity) it is advantageous to have an equivalent current that is symmetric. Defining

$$T^{\mu\nu} = \Theta^{\mu\nu} - \partial_\rho G^{\rho\mu\nu}, \quad (6.62)$$

with $G^{\rho\mu\nu} = (H^{\rho\mu\nu} + H^{\mu\nu\rho} + H^{\nu\rho\mu})/2$, where $H^{\rho\mu\nu}$ is the quantity appearing in $M^{\rho\mu\nu}$, one has a tensor that satisfies (exercise 6.6)

$$T^{\mu\nu} = T^{\nu\mu}, \quad (6.63)$$

$$\partial_\mu T^{\mu\nu} = \partial_\mu \Theta^{\mu\nu}, \quad (6.64)$$

$$M^{\mu\rho\sigma} = T^{\mu\rho}x^\sigma - T^{\mu\sigma}x^\rho. \quad (6.65)$$

6.5 (Abelian) gauge theories

In section 6.3 we have seen *global transformations* or gauge transformations of the first kind, e.g.

$$\phi(x) \rightarrow e^{i e \Lambda} \phi(x), \quad (6.66)$$

in which the $U(1)$ transformation involves an angle $e \Lambda$, independent of x . Gauge transformations of the second kind or *local gauge transformations* are transformations of the type

$$\phi(x) \rightarrow e^{i e \Lambda(x)} \phi(x), \quad (6.67)$$

i.e. the angle of the transformation depends on the space-time point x . The lagrangians which we have considered sofar are invariant under global gauge transformations and corresponding to this there exist a conserved Noether current. Any lagrangian containing derivatives, however, is not invariant under local gauge transformations,

$$\phi(x) \rightarrow e^{i e \Lambda(x)} \phi(x), \quad (6.68)$$

$$\phi^*(x) \rightarrow e^{-i e \Lambda(x)} \phi^*(x), \quad (6.69)$$

$$\partial_\mu \phi(x) \rightarrow e^{i e \Lambda(x)} \partial_\mu \phi(x) + i e \partial_\mu \Lambda(x) e^{i e \Lambda(x)} \phi(x), \quad (6.70)$$

where the last term spoils gauge invariance.

A solution is the one known as *minimal substitution* in which the derivative ∂_μ is replaced by a *covariant derivative* D_μ which satisfies

$$D_\mu \phi(x) \rightarrow e^{ie\Lambda(x)} D_\mu \phi(x). \quad (6.71)$$

For this purpose it is necessary to introduce a vector field A_μ ,

$$D_\mu \phi(x) \equiv (\partial_\mu + ieA_\mu(x))\phi(x), \quad (6.72)$$

The required transformation for D_μ then demands

$$\begin{aligned} D_\mu \phi(x) &= (\partial_\mu + ieA_\mu(x))\phi(x) \\ &\rightarrow e^{ie\Lambda} \partial_\mu \phi + ie(\partial_\mu \Lambda) e^{ie\Lambda} \phi + ieA'_\mu e^{ie\Lambda} \phi \\ &= e^{ie\Lambda} (\partial_\mu + ie(A'_\mu + \partial_\mu \Lambda)) \phi \\ &\equiv e^{ie\Lambda} (\partial_\mu + ieA_\mu) \phi. \end{aligned} \quad (6.73)$$

Thus the covariant derivative has the correct transformation behavior provided

$$A_\mu \rightarrow A_\mu - \partial_\mu \Lambda, \quad (6.74)$$

the behavior which we have encountered before as a gauge freedom for massless vector fields with the (free) lagrangian density $\mathcal{L} = -(1/4)F_{\mu\nu}F^{\mu\nu}$. Replacing derivatives by covariant derivatives and adding the (free) part for the massless vector fields to the original lagrangian therefore produces a gauge invariant lagrangian,

$$\mathcal{L}(\phi, \partial_\mu \phi) \implies \mathcal{L}(\phi, D_\mu \phi) - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}. \quad (6.75)$$

The field ϕ is used here in a general sense standing for any possible field. As an example consider the Dirac lagrangian,

$$\mathcal{L} = \frac{i}{2} (\bar{\psi} \gamma^\mu \partial_\mu \psi - (\partial_\mu \bar{\psi}) \gamma^\mu \psi) - M \bar{\psi} \psi.$$

Minimal substitution $\partial_\mu \psi \rightarrow (\partial_\mu + ieA_\mu)\psi$ leads to the gauge invariant lagrangian

$$\mathcal{L} = \frac{i}{2} \bar{\psi} \overleftrightarrow{\partial} \psi - M \bar{\psi} \psi - e \bar{\psi} \gamma^\mu \psi A_\mu - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}. \quad (6.76)$$

We note first of all that the coupling of the Dirac field (electron) to the vector field (photon) can be written in the familiar interaction form

$$\mathcal{L}_{int} = -e \bar{\psi} \gamma^\mu \psi A_\mu = -e j^\mu A_\mu, \quad (6.77)$$

involving the interaction of the charge ($\rho = j^0$) and three-current density (\mathbf{j}) with the electric potential ($\phi = A^0$) and the vector potential (\mathbf{A}), $-e j^\mu A_\mu = -e \rho \phi + e \mathbf{j} \cdot \mathbf{A}$. The equation of motion for the fermion follow from

$$\begin{aligned} \frac{\delta \mathcal{L}}{\delta(\partial_\mu \bar{\psi})} &= -\frac{i}{2} \gamma^\mu \psi, \\ \frac{\delta \mathcal{L}}{\delta \bar{\psi}} &= \frac{i}{2} \overleftrightarrow{\partial} \psi - M \psi - e \mathbf{A} \psi \end{aligned}$$

giving the Dirac equation in an electromagnetic field,

$$(iD - M) \psi = (i\overleftrightarrow{\partial} - e\mathbf{A} - M) \psi = 0. \quad (6.78)$$

For the photon the equations of motion follow from

$$\begin{aligned}\frac{\delta\mathcal{L}}{\delta(\partial_\mu A_\nu)} &= -F^{\mu\nu}, \\ \frac{\delta\mathcal{L}}{\delta A_\nu} &= -e\bar{\psi}\gamma^\nu\psi,\end{aligned}$$

giving the Maxwell equation coupling to the electromagnetic current,

$$\partial_\mu F^{\mu\nu} = j^\nu, \quad (6.79)$$

where $j^\mu = e\bar{\psi}\gamma^\mu\psi$.

Exercises

Exercise 6.1

- (a) Show that the Klein-Gordon equation for the real scalar field can be derived from the lagrangian density

$$\mathcal{L} = \frac{1}{2} \partial_\mu\phi\partial^\mu\phi - \frac{1}{2} M^2 \phi^2.$$

- (b) Show that the Klein-Gordon equation for the complex scalar field (considering ϕ and ϕ^* as independent fields can be derived from the lagrangian density

$$\mathcal{L} = \partial_\mu\phi^*\partial^\mu\phi - M^2\phi^*\phi.$$

Exercise 6.2

- (a) Show that the homogeneous Maxwell equations can be derived from the lagrangian density

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu}F^{\mu\nu}.$$

- (b) What is the form of the interaction term involving a current j_μ and the field A^μ that will give the inhomogeneous Maxwell equations, $\partial_\mu F^{\mu\nu} = j^\nu$.
- (c) Show that the interaction term is invariant under gauge transformations only if the current j_μ is conserved, i.e. $\partial_\mu j^\mu = 0$ (Note that the addition of a total derivative to the lagrangian density does not modify the equations of motion).

Exercise 6.3

Show that the current $j_\mu = i\phi^* \overleftrightarrow{\partial}_\mu \phi$ for a complex scalar field is connected to a $U(1)$ transformation on the fields, $\phi \rightarrow e^{i\Lambda}\phi$.

Exercise 6.4

Given the Dirac equation for a (negatively charged) spin-1/2 particle in an external electromagnetic field, $[i\gamma^\mu(\partial_\mu - ieA_\mu) - M]\psi = 0$, give the equation which is satisfied by $\psi^c = C\bar{\psi}^T$, and deduce what is the charge of the antiparticle as compared to a particle.

Exercise 6.5

Show that the Dirac equation for an electron with charge $-e$ in an external electromagnetic field $A^\mu = (A^0, \mathbf{A})$ (see also exercise 6.4) for a stationary solution in the non-relativistic ($|E_{\text{nr}}| \ll M$) and weak-field ($|A^0| \ll M$) limits yields in the standard representation for the 'upper (two) components' ψ_u ,

$$\left(\frac{1}{2M} (-i\nabla + e\mathbf{A})^2 + \frac{e}{2M} \boldsymbol{\sigma} \cdot \mathbf{B} - eA^0 \right) \psi_u(\mathbf{r}, t) = E_{\text{nr}} \psi_u(\mathbf{r}, t),$$

with $E_{\text{nr}} \equiv E - M \ll M$. This relation is known as the Pauli equation.

Comment: this relation is known as the Pauli equation. It shows that for a spin 1/2 particle the g -factor in the coupling of spin to a magnetic field takes the value $g_s = 2$,

$$H_{\text{int}} \equiv -g_s \boldsymbol{\mu}_e \cdot \mathbf{B} = -g_s \frac{Q_e}{2m_e} \mathbf{s} \cdot \mathbf{B}.$$

Exercise 6.6 (optional)

Prove the properties of the tensor $T^{\mu\nu}$ in section 6.4.

Hints: it may be useful to realize that $G^{\mu\rho\sigma}$ is antisymmetric in first two indices; use $\partial_\mu T^{\mu\nu} = 0$ and $\partial_\mu M^{\mu\rho\sigma} = 0$, in order to show that $\Theta^{\rho\sigma} - \Theta^{\sigma\rho} = \partial_\mu H^{\mu\rho\sigma}$; finally note that it is sufficient to show the last equation for $\int d\sigma_\mu M^{\mu\rho\sigma}$.

Chapter 7

Quantization of fields

7.1 Canonical quantization

We will first recall the example of classical mechanics for one coordinate $q(t)$, starting from the lagrangian $L(q, \dot{q})$ also considered in the previous chapter,

$$L(q, \dot{q}) = \frac{1}{2} m \dot{q}^2 - V(q). \quad (7.1)$$

The Hamiltonian (also corresponding to a conserved quantity because of time translation invariance) is given by

$$H(p, q) = p \dot{q} - L(q, \dot{q}(q, p)) = \frac{p^2}{2m} + V(q), \quad (7.2)$$

where the (canonical) momentum $p = \partial L / \partial \dot{q}$, in our example $p = m \dot{q}$. Further, there are conserved quantities Q (Eq. 6.11) found through Noether's theorem. Actually the conserved quantities satisfy a similar algebra as the Lie algebras for continuous symmetries that they are obtained from. The bilinear product for the conserved quantities, however, is the Poisson bracket

$$[A, B]_{\text{P}} = \frac{dA}{dq} \frac{dB}{dp} - \frac{dB}{dq} \frac{dA}{dp}. \quad (7.3)$$

For variations $\delta O(p, q) \approx (dO/d\lambda)_{\lambda=0} \delta \lambda$ under symmetry transformations one has

$$\frac{dO}{d\lambda} = [O, Q]_{\text{P}}.$$

Examples are

$$\frac{dO}{dt} = [O, H]_{\text{P}} \quad \text{and} \quad \frac{dO}{dq} = [O, p]_{\text{P}}.$$

The basic Poisson bracket is actually the one between q and its canonical momentum p , namely $[q, p]_{\text{P}} = 1$. Quantization of the theory is achieved by imposing canonical commutation relations between q and p ,

$$[q, p] = i, \quad (7.4)$$

with a possible realization as operators in the Hilbert space of (coordinate space) wave functions through $q_{op}\psi(q) = q\psi(q)$ and $p_{op}\psi(q) = -i d\psi/dq$. This is nothing else as representing the classical Poisson algebra as operators in the Hilbert space.

An immediate generalization for fields can be obtained by considering them as coordinates, labeled by the position,

$$q_{\mathbf{x}}(t) = \frac{1}{\Delta^3 x} \int_{\Delta^3 x} d^3 x \phi(\mathbf{x}, t), \quad (7.5)$$

etc. The lagrangian is given by

$$\begin{aligned} L &= \int d^3x \mathcal{L}(\phi(x), \partial_\mu \phi(x)) = \int d^3x \mathcal{L}(\phi, \dot{\phi}, \nabla \phi) \\ &= \sum_{\mathbf{x}} \Delta^3 x \mathcal{L}_{\mathbf{x}}(q_{\mathbf{x}}, \dot{q}_{\mathbf{x}}, q_{\mathbf{x}+\Delta \mathbf{x}}), \end{aligned} \quad (7.6)$$

In order to construct the hamiltonian it is necessary to find the canonical momenta,

$$p_{\mathbf{x}}(t) = \frac{\partial L}{\partial \dot{q}_{\mathbf{x}}} = \Delta^3 x \frac{\partial \mathcal{L}_{\mathbf{x}}}{\partial \dot{q}_{\mathbf{x}}} = \Delta^3 x \Pi_{\mathbf{x}}(t), \quad (7.7)$$

where $\Pi_{\mathbf{x}}(t)$ is obtained from the continuous field $\Pi(x) \equiv \delta \mathcal{L} / \delta(\partial_0 \phi)$. The hamiltonian then is

$$H = \sum_{\mathbf{x}} p_{\mathbf{x}} \dot{q}_{\mathbf{x}} - L = \sum_{\mathbf{x}} \Delta^3 x [\Pi_{\mathbf{x}} \dot{q}_{\mathbf{x}} - \mathcal{L}_{\mathbf{x}}] \quad (7.8)$$

$$= \sum_{\mathbf{x}} \Delta^3 x \mathcal{H}_{\mathbf{x}} \quad (7.9)$$

where

$$\mathcal{H}(x) = \frac{\delta \mathcal{L}}{\delta \dot{\phi}(x)} \dot{\phi}(x) - \mathcal{L}(x) = \Theta^{00}(x). \quad (7.10)$$

Note that this indeed corresponds to the zero-zero component (Θ^{00}) of the conserved energy-momentum stress tensor $\Theta^{\mu\nu}$, discussed in the section 6.2.

As an example, for the scalar field theory, we have

$$\begin{aligned} \mathcal{L}(x) &= \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} M^2 \phi^2 \\ &= \frac{1}{2} (\partial_0 \phi)^2 - \frac{1}{2} (\nabla \phi)^2 - \frac{1}{2} M^2 \phi^2 \end{aligned} \quad (7.11)$$

$$\Pi(x) = \frac{\delta \mathcal{L}}{\delta \partial_0 \phi} = \partial_0 \phi, \quad (7.12)$$

$$\mathcal{H}(x) = \Theta^{00}(x) = \frac{1}{2} (\partial_0 \phi)^2 + \frac{1}{2} (\nabla \phi)^2 + \frac{1}{2} M^2 \phi^2. \quad (7.13)$$

For the quantization procedure we can formulate a number of basic axioms of quantum field theory. Sometimes it is useful to keep in mind that, formally, the fields $\phi(x)$ can be considered as regular operators in the Hilbert space after smearing with a test function f ,

$$\Phi(f) = \int d^4x \phi(x) f(x). \quad (7.14)$$

In fact, the discretization procedure above is an explicit example, albeit with 'sharp' functions.

The essential conditions to consistently quantize a theory then are the following.

- **Canonical commutation relations**

The quantization of the theory is achieved by imposing the canonical quantization condition $[q_{\mathbf{x}}(t), p_{\mathbf{y}}(t)] = i \delta_{\mathbf{x}\mathbf{y}}$ or for the fields $\phi(x)$ and $\Pi(x)$ the so-called *equal time commutation relations*

$$[\phi(\mathbf{x}, t), \Pi(\mathbf{y}, t)] = i \delta^3(\mathbf{x} - \mathbf{y}), \quad (7.15)$$

with furthermore the relations $[\phi(\mathbf{x}, t), \phi(\mathbf{y}, t)] = [\Pi(\mathbf{x}, t), \Pi(\mathbf{y}, t)] = 0$.

- **Poincaré invariance**

The full variation of the fields, $\Delta \phi$, for symmetry transformations. This implies specific transformation properties in the Hilbert space for the quantum fields ϕ ,

$$U^\dagger(\Lambda, a) \phi^i(x) U(\Lambda, a) = R_j^i(\Lambda^{-1}) \phi^j(\Lambda^{-1}x - a), \quad (7.16)$$

or if

$$\begin{aligned} U(\Lambda, a) &= 1 + i\epsilon_\mu P^\mu - \frac{i}{2}\omega_{\mu\nu}M^{\mu\nu}, \\ R_j^i(\Lambda^{-1}) &= 1 - \frac{i}{2}\omega_{\mu\nu}(S^{\mu\nu})_j^i, \end{aligned}$$

one has

$$[\phi^i(x), P_\mu] = i\partial_\mu\phi^i(x), \quad (7.17)$$

$$[\phi^i(x), M_{\mu\nu}] = i(x_\mu\partial_\nu - x_\nu\partial_\mu)\phi^i(x) + (S_{\mu\nu})_j^i\phi^j(x), \quad (7.18)$$

with $S^{\mu\nu}$ given in 6.4. This must be valid for the – now operators – generators P^μ and $M^{\mu\nu}$ that in the previous chapter have been found in terms of the fields through Noether's theorem.

- **Causality**

Operators $\Phi(f)$ and $\Phi(g)$ for which the test functions are space-like separated can be measured simultaneously (*macroscopic causality*). The measurements cannot influence each other or $[\Phi(f), \Phi(g)] = 0$. *Microscopic causality* implies local commutativity,

$$[\phi(x), \phi(y)] = 0 \quad \text{if} \quad (x - y)^2 < 0. \quad (7.19)$$

7.2 Creation and annihilation operators

Before discussing (real and complex) scalar fields and Dirac fields we recall the analogy with the well-known harmonic oscillator as an example of quantization using creation and annihilation operators, sometimes referred to as second quantization. In simplified form the hamiltonian is given by

$$H = \frac{1}{2}P^2 + \frac{1}{2}\omega^2 Q^2, \quad (7.20)$$

where the coordinate Q and the momentum P satisfy the canonical commutation relations

$$[Q, P] = i, \quad [Q, Q] = [P, P] = 0 \quad (7.21)$$

Writing Q and P in terms of creation (a^\dagger) and annihilation (a) operators,

$$Q = \frac{1}{\sqrt{2\omega}}(a + a^\dagger) \quad \text{and} \quad P = -i\sqrt{\frac{\omega}{2}}(a - a^\dagger) \quad (7.22)$$

it is straightforward to check that the commutation relations between Q and P are equivalent with the commutation relations

$$[a, a^\dagger] = 1, \quad [a, a] = [a^\dagger, a^\dagger] = 0. \quad (7.23)$$

The hamiltonian in this case can be expressed in the number operator $N = a^\dagger a$,

$$\begin{aligned} H &= \omega \left\{ \left(\sqrt{\frac{\omega}{2}}Q - i\frac{P}{\sqrt{2\omega}} \right) \left(\sqrt{\frac{\omega}{2}}Q + i\frac{P}{\sqrt{2\omega}} \right) - \frac{i}{2}[Q, P] \right\} \\ &= \omega \left\{ a^\dagger a + \frac{1}{2} \right\} = \omega \left\{ N + \frac{1}{2} \right\}. \end{aligned} \quad (7.24)$$

It is straightforward to find the commutation relations between N and a and a^\dagger ,

$$[N, a^\dagger] = a^\dagger, \quad \text{and} \quad [N, a] = -a. \quad (7.25)$$

Defining states $|n\rangle$ as eigenstates of N with eigenvalue n , $N|n\rangle = n|n\rangle$ one finds

$$\begin{aligned} N a^\dagger |n\rangle &= (n+1) a^\dagger |n\rangle, \\ N a |n\rangle &= (n-1) a |n\rangle \end{aligned}$$

i.e. a^\dagger and a act as raising and lowering operators. From the normalizations one obtains $a^\dagger |n\rangle = \sqrt{n+1} |n+1\rangle$ and $a |n\rangle = \sqrt{n} |n-1\rangle$, and we see that a state $|0\rangle$ must exist for which $N|0\rangle = a|0\rangle = 0$. In this way one has found for the harmonic oscillator the spectrum of eigenstates $|n\rangle$ (with n a non-negative integer) with $E_n = (n+1/2)\omega$.

7.3 The real scalar field

We have expanded the (classical) field in plane wave solutions, which we have split into positive and negative energy pieces with (complex) coefficients $a(\mathbf{k})$ and $a^*(\mathbf{k})$ multiplying them. The quantization of the field is achieved by quantizing the coefficients in the Fourier expansion, e.g. the real scalar field $\phi(x)$ becomes

$$\phi(x) = \int \frac{d^3k}{(2\pi)^3 2E_k} [a(\mathbf{k}) e^{-ik \cdot x} + a^\dagger(\mathbf{k}) e^{ik \cdot x}], \quad (7.26)$$

where the Fourier coefficients $a(\mathbf{k})$ and $a^\dagger(\mathbf{k})$ are now operators. Note that we will often write $a(k)$ or $a^\dagger(k)$, but one needs to realize that in that case $k^0 = E_k = \sqrt{\mathbf{k}^2 + M^2}$. The canonical momentum becomes

$$\Pi(x) = \dot{\phi}(x) = \frac{-i}{2} \int \frac{d^3k}{(2\pi)^3} [a(k) e^{-ik \cdot x} - a^\dagger(k) e^{ik \cdot x}]. \quad (7.27)$$

It is easy to check that these equations can be inverted (see Exercise 2.4 for the classical field)

$$a(k) = \int d^3x e^{ik \cdot x} i \overleftrightarrow{\partial}_0 \phi(x), \quad (7.28)$$

$$a^\dagger(k') = \int d^3x \phi(x) i \overleftrightarrow{\partial}_0 e^{-ik' \cdot x} \quad (7.29)$$

It is straightforward to prove that the equal time commutation relations between $\phi(x)$ and $\Pi(x')$ are equivalent with 'harmonic oscillator - like' commutation relations between $a(k)$ and $a^\dagger(k')$, i.e.

$$\begin{aligned} [\phi(x), \Pi(x')]_{x^0=x'^0} &= i \delta^3(\mathbf{x} - \mathbf{x}') \quad \text{and} \\ [\phi(x), \phi(x')]_{x^0=x'^0} &= [\Pi(x), \Pi(x')]_{x^0=x'^0} = 0, \end{aligned} \quad (7.30)$$

is equivalent with

$$\begin{aligned} [a(k), a^\dagger(k')] &= (2\pi)^3 2E_k \delta^3(\mathbf{k} - \mathbf{k}') \quad \text{and} \\ [a(k), a(k')] &= [a^\dagger(k), a^\dagger(k')] = 0. \end{aligned} \quad (7.31)$$

The hamiltonian can be rewritten in terms of a number operator $N(k) = N(\mathbf{k}) = a^\dagger(k)a(k)$, which represents the 'number of particles' with momentum \mathbf{k} .

$$\begin{aligned} H &= \int d^3x \left[\frac{1}{2} (\partial_0 \phi)^2 + \frac{1}{2} (\nabla \phi)^2 + \frac{1}{2} M^2 \phi^2 \right] \\ &= \int \frac{d^3k}{(2\pi)^3 2E_k} \frac{E_k}{2} (a^\dagger(k)a(k) + a(k)a^\dagger(k)) \end{aligned} \quad (7.32)$$

$$= \int \frac{d^3k}{(2\pi)^3 2E_k} E_k N(k) + E_{\text{vac}}, \quad (7.33)$$

where the necessity to commute $a(k)a^\dagger(k)$ (as in the case of the quantum mechanics case) leads to a zero-point energy, in field theory also referred to as vacuum energy

$$E_{\text{vac}} = \frac{1}{2} V \int \frac{d^3 k}{(2\pi)^3} E_k, \quad (7.34)$$

where $V = (2\pi)^3 \delta^3(\mathbf{0})$ is the space-volume. This term will be addressed below. For the momentum operator one has

$$P^i = \int d^3 x \Theta^{0i}(x) = \frac{1}{2} \int d^3 x \partial^{\{0} \phi \partial^{i\}} \phi = \int \frac{d^3 k}{(2\pi)^3} \frac{1}{2E_k} k^i N(k), \quad (7.35)$$

where the vacuum contribution disappears because of rotational symmetry. Just as in the case of the harmonic oscillator it is essential (axiom) that there exists a ground state $|0\rangle$ that is annihilated by $a(k)$, $a(k)|0\rangle = 0$. The rest of the states are then obtained from the groundstate via the creation operator, defining particle states $|k\rangle = |\mathbf{k}\rangle$ (with positive energy, $k^0 = E_k = \sqrt{\mathbf{k}^2 + M^2}$),

$$|k\rangle = a^\dagger(k)|0\rangle, \quad (7.36)$$

and multiparticle states

$$|(k_1)^{n_1} (k_2)^{n_2} \dots\rangle = \frac{(a^\dagger(k_1))^{n_1}}{\sqrt{n_1!}} \frac{(a^\dagger(k_2))^{n_2}}{\sqrt{n_2!}} \dots |0\rangle, \quad (7.37)$$

normalized as

$$\langle k|k'\rangle = (2\pi)^3 2E_k \delta^3(\mathbf{k} - \mathbf{k}'), \quad (7.38)$$

and satisfying the completeness condition

$$1 = \int \frac{d^4 k}{(2\pi)^4} 2\pi \delta(k^2 - M^2) \theta(k^0) |k\rangle \langle k| = \int \frac{d^3 k}{(2\pi)^3} \frac{1}{2E_k} |k\rangle \langle k|. \quad (7.39)$$

The problem with the vacuum or zero-point energy, which now contains an infinite number of oscillators, is solved by subtracting it as an (infinite) constant, which amounts to redefining H as

$$H = \int d^3 x : \mathcal{H}(x) := \int \frac{d^3 k}{(2\pi)^3} \frac{1}{2E_k} E_k N(k). \quad (7.40)$$

This procedure is known as *normal ordering*, i.e. writing all annihilation operators to the right of the creation operators, assuring that the vacuum (by definition) has eigenvalue 0!

For the purpose of normal ordering it is convenient to decompose the field in positive and negative frequency parts,

$$\phi(x) = \phi_+(x) + \phi_-(x), \quad (7.41)$$

$$\phi_+(x) = \int \frac{d^3 k}{(2\pi)^3} \frac{1}{2E_k} a(k) e^{-ik \cdot x}, \quad (7.42)$$

$$\phi_-(x) = \int \frac{d^3 k}{(2\pi)^3} \frac{1}{2E_k} a^\dagger(k) e^{ik \cdot x}. \quad (7.43)$$

The normal ordered product can be expressed as

$$: \phi(x)\phi(y) := \phi_+(x)\phi_+(y) + \phi_-(x)\phi_+(y) + \phi_-(y)\phi_+(x) + \phi_-(x)\phi_-(y). \quad (7.44)$$

The 1-particle wave function $e^{-ip \cdot x}$ is obtained via

$$\langle 0|\phi(x)|p\rangle = \langle 0|\phi_+(x)|p\rangle = \int \frac{d^3 k}{(2\pi)^3} \frac{1}{2E_k} \langle 0|a(k)a^\dagger(p)|0\rangle e^{-ik \cdot x} = e^{-ip \cdot x} \quad (7.45)$$

$$\langle p|\phi(x)|0\rangle = \langle p|\phi_-(x)|0\rangle = e^{ip \cdot x}. \quad (7.46)$$

In order to ensure the consistency of the theory it is necessary to check that the operators P^ν and $M^{\mu\nu}$ obtained from the conserved currents $\Theta^{\rho\sigma}$ and $M^{\mu\rho\sigma}$ are indeed the generators of the Poincaré group, i.e. that they satisfy the required commutation relations in Eqs (7.17) and (7.18).

The last item to be checked for the scalar field are the causality condition. In order to calculate $[\phi(x), \phi(y)]$ consider

$$[\phi_+(x), \phi_-(y)] \equiv i\Delta_+(x-y), \quad (7.47)$$

$$\begin{aligned} &= \int \frac{d^3k}{(2\pi)^3 2E_k} \int \frac{d^3k'}{(2\pi)^3 2E_{k'}} e^{-ik \cdot x + ik' \cdot y} [a(k), a^\dagger(k')] \\ &= \int \frac{d^3k}{(2\pi)^3 2E_k} e^{-ik \cdot (x-y)}, \end{aligned} \quad (7.48)$$

$$[\phi_-(x), \phi_+(y)] \equiv i\Delta_-(x-y), \quad (7.49)$$

$$= -\int \frac{d^3k}{(2\pi)^3 2E_k} e^{ik \cdot (x-y)} = -i\Delta_+(y-x), \quad (7.50)$$

or as integrals over d^4k ,

$$i\Delta_+(x) = \int \frac{d^4k}{(2\pi)^4} \theta(k^0) 2\pi \delta(k^2 - M^2) e^{-ik \cdot x}, \quad (7.51)$$

$$\begin{aligned} i\Delta_-(x) &= -\int \frac{d^4k}{(2\pi)^4} \theta(k^0) 2\pi \delta(k^2 - M^2) e^{ik \cdot x} \\ &= -\int \frac{d^4k}{(2\pi)^4} \theta(-k^0) 2\pi \delta(k^2 - M^2) e^{-ik \cdot x} = -i\Delta_+(-x) \end{aligned} \quad (7.52)$$

The result for the *invariant commutator function* is

$$[\phi(x), \phi(y)] = i\Delta(x-y) = i(\Delta_+(x-y) + \Delta_-(x-y)), \quad (7.53)$$

which has the properties

(i) $i\Delta(x) = i\Delta_+(x) + i\Delta_-(x)$ can be expressed as

$$i\Delta(x) = \int \frac{d^4k}{(2\pi)^4} \epsilon(k^0) 2\pi \delta(k^2 - M^2) e^{-ik \cdot x}, \quad (7.54)$$

where $\epsilon(k^0) = \theta(k^0) - \theta(-k^0)$.

(ii) $\Delta(x)$ is a solution of the homogeneous Klein-Gordon equation.

(iii) $\Delta(0, \mathbf{x}) = 0$ and hence $\Delta(x) = 0$ for $x^2 < 0$.

(iv) The equal time commutation relations follow from

$$\left. \frac{\partial}{\partial t} \Delta(x) \right|_{t=0} = -\delta^3(x). \quad (7.55)$$

(v) For $M = 0$,

$$\Delta(x) = -\frac{\epsilon(x^0)}{2\pi} \delta(x^2). \quad (7.56)$$

7.4 The complex scalar field

In spite of the similarity with the case of the real field, we will consider it as a repetition of the quantization procedure, extending it with the charge operator and the introduction of particle and antiparticle operators. The field satisfies the Klein-Gordon equation and the density current ($U(1)$ transformations) and the energy-momentum tensor are

$$j_\mu = i \phi^* \overleftrightarrow{\partial}_\mu \phi, \quad (7.57)$$

$$\Theta_{\mu\nu} = \partial_{\{\mu} \phi^* \partial_{\nu\}} \phi - \mathcal{L} g_{\mu\nu}. \quad (7.58)$$

The quantized fields are written as

$$\phi(x) = \int \frac{d^3k}{(2\pi)^3 2E_k} [a(\mathbf{k}) e^{-ik \cdot x} + b^\dagger(\mathbf{k}) e^{ik \cdot x}], \quad (7.59)$$

$$\phi^\dagger(x) = \int \frac{d^3k}{(2\pi)^3 2E_k} [b(\mathbf{k}) e^{-ik \cdot x} + a^\dagger(\mathbf{k}) e^{ik \cdot x}], \quad (7.60)$$

and satisfy the equal time commutation relation (only nonzero ones)

$$[\phi(x), \partial_0 \phi^\dagger(y)]_{x^0=y^0} = i \delta^3(x-y), \quad (7.61)$$

which is equivalent to the relations (only nonzero ones)

$$[a(k), a^\dagger(k')] = [b(k), b^\dagger(k')] = (2\pi)^3 2E_k \delta^3(\mathbf{k} - \mathbf{k}'). \quad (7.62)$$

The hamiltonian is as before given by the normal ordered expression

$$\begin{aligned} H &= \int d^3x : \Theta^{00}(x) : \\ &= \int \frac{d^3k}{(2\pi)^3 2E_k} E_k : [a^\dagger(k)a(k) + b(k)b^\dagger(k)] : \\ &= \int \frac{d^3k}{(2\pi)^3 2E_k} E_k [a^\dagger(k)a(k) + b^\dagger(k)b(k)], \end{aligned} \quad (7.63)$$

i.e. particles (created by a^\dagger) and antiparticles (created by b^\dagger) with the same momentum contribute equally to the energy. Also the charge operator requires normal ordering (in order to give the vacuum eigenvalue zero),

$$\begin{aligned} Q &= i \int d^3x : [\phi^\dagger \partial_0 \phi - \partial_0 \phi^\dagger \phi(x)] : \\ &= \int \frac{d^3k}{(2\pi)^3 2E_k} : [a^\dagger(k)a(k) - b(k)b^\dagger(k)] : \\ &= \int \frac{d^3k}{(2\pi)^3 2E_k} [a^\dagger(k)a(k) - b^\dagger(k)b(k)]. \end{aligned} \quad (7.64)$$

The commutator of ϕ and ϕ^\dagger is as for the real field given by

$$[\phi(x), \phi^\dagger(y)] = i\Delta(x-y). \quad (7.65)$$

7.5 The Dirac field

From the lagrangian density

$$\mathcal{L} = \frac{i}{2} \bar{\psi} \gamma^\mu \overleftrightarrow{\partial}_\mu \psi - M \bar{\psi} \psi, \quad (7.66)$$

the conserved density and energy-momentum currents are easily obtained,

$$j_\mu = \bar{\psi} \gamma_\mu \psi, \quad (7.67)$$

$$\Theta_{\mu\nu} = \frac{i}{2} \bar{\psi} \gamma_\mu \overleftrightarrow{\partial}_\nu \psi - \left(\frac{i}{2} \bar{\psi} \overleftrightarrow{\not{\partial}} \psi - M \bar{\psi} \psi \right) g_{\mu\nu}. \quad (7.68)$$

The canonical momentum and the hamiltonian are given by

$$\Pi(x) = \frac{\delta \mathcal{L}}{\delta \dot{\psi}(x)} = i \psi^\dagger(x), \quad (7.69)$$

$$\begin{aligned} \mathcal{H}(x) &= \Theta^{00}(x) = -\frac{i}{2} \bar{\psi} \gamma^i \overleftrightarrow{\partial}_i \psi + M \bar{\psi} \psi \\ &= i \bar{\psi} \gamma^0 \partial_0 \psi = i \psi^\dagger \partial_0 \psi, \end{aligned} \quad (7.70)$$

where the last line is obtained by using the Dirac equation.

The quantized fields are written

$$\psi(x) = \sum_s \int \frac{d^3 k}{(2\pi)^3 2E_k} [b(\mathbf{k}, s) u(k, s) e^{-i k \cdot x} + d^\dagger(\mathbf{k}, s) v(k, s) e^{i k \cdot x}], \quad (7.71)$$

$$\bar{\psi}(x) = \sum_s \int \frac{d^3 k}{(2\pi)^3 2E_k} [b^\dagger(\mathbf{k}, s) \bar{u}(k, s) e^{i k \cdot x} + d(\mathbf{k}, s) \bar{v}(k, s) e^{-i k \cdot x}]. \quad (7.72)$$

In terms of the operators for the b and d quanta the hamiltonian and charge operators are (omitting mostly the spin summation in the rest of this section)

$$H = \int d^3 x : \psi^\dagger(x) i \partial_0 \psi(x) : \quad (7.73)$$

$$= \int \frac{d^3 k}{(2\pi)^3 2E_k} E_k : [b^\dagger(k) b(k) - d(k) d^\dagger(k)] : \quad (7.74)$$

$$Q = \int d^3 x : \psi^\dagger \psi : \quad (7.75)$$

$$= \int \frac{d^3 k}{(2\pi)^3 2E_k} : [b^\dagger(k) b(k) + d(k) d^\dagger(k)] : , \quad (7.76)$$

which seems to cause problems as the antiparticles (d-quanta) contribute negatively to the energy and the charges of particles (b-quanta) and antiparticles (d-quanta) are the same.

The solution is the introduction of anticommutation relations,

$$\{b(k, s), b^\dagger(k', s')\} = \{d(k, s), d^\dagger(k', s')\} = (2\pi)^3 2E_k \delta^3(\mathbf{k} - \mathbf{k}') \delta_{ss'}. \quad (7.77)$$

Note that achieving normal ordering, i.e. interchanging creation and annihilation operators, then leads to additional minus signs and

$$H = \int \frac{d^3 k}{(2\pi)^3 2E_k} E_k [b^\dagger(k) b(k) + d^\dagger(k) d(k)] \quad (7.78)$$

$$Q = \int \frac{d^3 k}{(2\pi)^3 2E_k} [b^\dagger(k) b(k) - d^\dagger(k) d(k)]. \quad (7.79)$$

Also for the field and the canonical conjugate momentum anticommutation relations are considered,

$$\begin{aligned} \{\psi_i(x), \psi_j^\dagger(y)\}_{x^0=y^0} &= \int \frac{d^3 k}{(2\pi)^3 2E_k} \left[\sum_s u_i(k, s) u_j^\dagger(k, s) e^{-i k \cdot (x-y)} \right. \\ &\quad \left. + v_i(k, s) v_j^\dagger(k, s) e^{i k \cdot (x-y)} \right]_{x^0=y^0}. \end{aligned}$$

Using the positive and negative energy projection operators discussed in section 4, one has

$$\begin{aligned}
\{\psi_i(x), \psi_j^\dagger(y)\}_{x^0=y^0} &= \int \frac{d^3k}{(2\pi)^3 2E_k} \left[(\not{k} + M)_{il} (\gamma_0)_{lj} e^{-ik \cdot (x-y)} \right. \\
&\quad \left. + (\not{k} - M)_{il} (\gamma_0)_{lj} e^{ik \cdot (x-y)} \right]_{x^0=y^0} \\
&= \int \frac{d^3k}{(2\pi)^3 2E_k} 2E_k (\gamma_0 \gamma_0)_{ij} e^{i\mathbf{k} \cdot (\mathbf{x}-\mathbf{y})} \\
&= \delta^3(\mathbf{x} - \mathbf{y}) \delta_{ij}.
\end{aligned} \tag{7.80}$$

For the scalar combination $\{\psi_i(x), \bar{\psi}_j(y)\}$ at arbitrary times one has

$$\begin{aligned}
\{\psi_i(x), \bar{\psi}_j(y)\} &= \int \frac{d^3k}{(2\pi)^3 2E_k} \left[(\not{k} + M)_{ij} e^{-ik \cdot (x-y)} + (\not{k} - M)_{ij} e^{ik \cdot (x-y)} \right] \\
&= (i\partial_x + M)_{ij} \int \frac{d^3k}{(2\pi)^3 2E_k} \left[e^{-ik \cdot (x-y)} + -e^{ik \cdot (x-y)} \right] \\
&= (i\partial_x + M)_{ij} i\Delta(x-y),
\end{aligned} \tag{7.81}$$

where $i\Delta = i\Delta_+ + i\Delta_-$ is the same invariant commutator function as encountered before. When we would have started with commutation relations for the field ψ and the canonical momentum, we would have obtained

$$[\psi_i(x), \bar{\psi}_j(y)] = (i\partial_x + M)_{ij} i\Delta^1(x-y), \tag{7.82}$$

where $i\Delta^1 = i\Delta_+ - i\Delta_-$, which however has wrong causality properties! Therefore the relation between spin and statistics is required to get micro-causality (which is also known as the *spin statistics theorem*).

7.6 The electromagnetic field

From the lagrangian density

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu}, \tag{7.83}$$

the canonical momenta are

$$\Pi^0 = \frac{\delta\mathcal{L}}{\delta\dot{A}_0} = 0, \tag{7.84}$$

$$\Pi^i = \frac{\delta\mathcal{L}}{\delta\dot{A}_i} = F^{i0} = E^i, \tag{7.85}$$

which reflects the gauge freedom, but has the problem of being noncovariant, as the vanishing of Π^0 induces a constraint. It is possible to continue in a covariant way with the lagrangian

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{\lambda}{2} (\partial_\mu A^\mu)^2,$$

This gives the equations of motion discussed before, it implies the Lorentz constraint and leads to the canonical momenta

$$\Pi^0 = -\lambda(\partial_\rho A^\rho), \tag{7.86}$$

$$\Pi^i = E^i. \tag{7.87}$$

If one wants to impose canonical commutation relations $\partial_\mu A^\mu = 0$ cannot hold as an operator identity, but we must restrict ourselves to the weaker condition

$$\langle B | \partial_\mu A^\mu | A \rangle = 0, \tag{7.88}$$

for physical states $|A\rangle$ and $|B\rangle$.

The quantized field is expanded as

$$A_\mu(x) = \int \frac{d^3k}{(2\pi)^3 2E_k} \sum_{\lambda=0}^3 \epsilon_\mu^{(\lambda)}(k) [c(\mathbf{k}, \lambda) e^{-i\mathbf{k}\cdot\mathbf{x}} + c^\dagger(\mathbf{k}, \lambda) e^{i\mathbf{k}\cdot\mathbf{x}}], \quad (7.89)$$

with four independent vectors $\epsilon_\mu^{(\lambda)}$, containing a time-like photon, a longitudinal photon and two transverse photons. The canonical equal time commutation relations are

$$[A_\mu(x), \Pi_\nu(y)]_{x^0=y^0} = i g_{\mu\nu} \delta^3(\mathbf{x} - \mathbf{y}), \quad (7.90)$$

where $\Pi^\nu = F^{\nu 0} - \lambda g^{\nu 0} (\partial_\rho A^\rho)$ and we have furthermore $[A_\mu(x), A_\nu(y)] = [\Pi_\mu(x), \Pi_\nu(y)] = 0$. In fact the commutation relations imply

$$[\dot{A}_\mu(x), A_\nu(y)]_{x^0=y^0} = i g_{\mu\nu} \delta^3(\mathbf{x} - \mathbf{y}), \quad (7.91)$$

and are equivalent with

$$[c(k, \lambda), c^\dagger(k', \lambda')] = -g^{\lambda\lambda'} 2E_k (2\pi)^3 \delta^3(\mathbf{k} - \mathbf{k}'). \quad (7.92)$$

Note that for the transverse states there are no problems with the normalization and the statistics of the states as $-g^{ij} = \delta^{ij}$ for $i, j = 1, 2$. The hamiltonian in terms of the creation and annihilation operators is (after normal ordering) given by

$$H = \int \frac{d^3k}{(2\pi)^3 2E_k} E_k \left[\sum_{\lambda=1}^3 c^\dagger(k, \lambda) c(k, \lambda) - c^\dagger(k, 0) c(k, 0) \right], \quad (7.93)$$

which does exhibit problems with the time-like photon. These problems are solved by the Lorentz constraint between physical states given above, for which it is sufficient that $\partial_\mu A_+^\mu |A\rangle = 0$, where A_+^μ is the part of the vector field containing the annihilation operators. It gives

$$\sum_{\lambda=0}^3 k^\mu \epsilon_\mu^{(\lambda)}(k) c(k, \lambda) |A\rangle = 0. \quad (7.94)$$

Choosing $k^\mu = (|k^3|, 0, 0, k^3)$ this reads

$$\begin{aligned} (|k^3| c^{(0)}(k) - k^3 c^{(3)}(k)) |A\rangle &= 0 \\ (c^{(0)}(k) \mp c^{(3)}(k)) |A\rangle &= 0, \end{aligned} \quad (7.95)$$

i.e. *one* time-like photon by itself is not allowed! This solves the problems with the normalization and the negative energies.

Exercises

Exercise 7.1 (optional)

If you haven't seen Poisson brackets in classical mechanics you should

- Proof that Poisson brackets (Eq. 7.3) satisfy the conditions of a Lie algebra.
- Proof for a quantity $O(p, q)$ that $dO/d\lambda = [O, Q]_{\text{P}}$, where Q is the conserved quantity in Eq. 6.11.
- Calculate the Poisson brackets for the coordinates \mathbf{x} and the standard conserved quantities \mathbf{p} and $\boldsymbol{\ell}$, i.e. what are $[r_i, p_j]_{\text{P}}$, $[\ell_i, r_j]_{\text{P}}$, and $[\ell_i, p_j]_{\text{P}}$.
- Use Poisson brackets to calculate \dot{q} and \dot{p} , which reproduce the Hamilton equations.

Exercise 7.2

Prove that the equal time commutation relations between $\phi(x)$ and $\Pi(x')$ are equivalent with the commutation relations between $a(k)$ and $a^\dagger(k')$

Exercise 7.3

(a) Show that

$$e^{-i a^\mu P_\mu} \phi(x) e^{i a^\mu P_\mu} = \phi(x - a)$$

requires $[\phi(x), P_\mu] = i \partial_\mu \phi(x)$.

(b) Check the above commutation relation $[\phi(x), P_\mu]$ for the (real) scalar field using the expressions for the fields and momentum operator in terms of creation and annihilation operators. Is the subtraction of 'zero point' contributions essential in this check?

Exercise 7.4 (Green's functions)

(a) Show that

$$\Delta^{\text{hom}}(x) = \frac{1}{(2\pi)^4} \int d^4k e^{-ik \cdot x} \delta(k^2 - M^2) f(k)$$

is a solution of the homogeneous Klein-Gordon equation, $(\square + M^2) \Delta^{\text{hom}}(x) = 0$. It is invariant, $\Delta^{\text{hom}}(\Lambda x) = \Delta^{\text{hom}}(x)$ if f is an invariant function, $f(\Lambda k) = f(k)$ for Lorentz transformations Λ .


(b) Show that

$$\Delta^{\text{inhom}}(x) = \frac{1}{(2\pi)^4} \int d^4k e^{-ikx} \frac{1}{k^2 - M^2}$$

is a solution of the inhomogeneous Klein-Gordon equation, $(\square + M^2) \Delta^{\text{inhom}}(x) = -\delta^4(x)$.

(c) What are the poles in the integral under (b) in the (complex) k_0 plane?

(d) Depending on the paths in the k_0 plane going from $k_0 = -\infty$ to $k_0 = +\infty$ one can distinguish four different Green's functions (solutions to the above inhomogeneous Klein-Gordon equation):



Show that by appropriate shifts of the poles into the complex plane, we can write for Δ_R an integral where the k^0 integration remains along the real axis,

$$\Delta_R = \frac{1}{(2\pi)^4} \int d^4k e^{-ikx} \frac{1}{k^2 - M^2 + i\epsilon k^0}$$


(with the implicit prescription to take $\epsilon \rightarrow 0$ after integration). Give also the expression for Δ_C .

(e) Show that $\Delta_R(x) = 0$ if $x^0 < 0$. (Similarly one can show that for the advanced Green's function $\Delta_A(x) = 0$ if $x^0 > 0$).

(f) By performing the k_0 integration over a closed path C going around the poles show that

$$\Delta(x) = -\frac{1}{(2\pi)^4} \int_C d^4k e^{-ikx} \frac{1}{k^2 - M^2}$$

are homogeneous solutions for the contours:



One way of showing this is to write these contours as integrals of the form under (a). In that case one actually needs discontinuous functions $f(k)$ such as the step function, $f(k) \propto \theta(k^0)$, or $\epsilon(k^0) = \theta(k^0) - \theta(-k^0)$.

- (g) Show that the homogeneous solution Δ in (f) satisfies $\Delta(0, \mathbf{x}) = 0$ and argue that one can use Lorentz invariance to show that $\Delta(x) = 0$ for $x^2 < 0$.
- (h) Show that the causal Green's function

$$\Delta_C(x) = \theta(x^0) \Delta_+(x) - \theta(-x^0) \Delta_-(x).$$

- (i) (Optional) Give similar expressions for the other Green's functions under (d) in terms of the homogeneous solutions under (f).

Chapter 8

Discrete symmetries

In this chapter we discuss the discrete symmetries, parity (P), time reversal (T) and charge conjugation (C). The consequences of P, T and C for classical quantities is shown in the table 1.

8.1 Parity

The parity operator transforms

$$x^\mu = (t, \mathbf{r}) \longrightarrow \tilde{x}^\mu \equiv x_\mu = (t, -\mathbf{r}). \quad (8.1)$$

We will consider the transformation properties for a fermion field $\psi(x)$, writing

$$\psi(x) \longrightarrow P_{op}\psi(x)P_{op}^{-1} = \eta_P A\psi(\tilde{x}) \equiv \psi^P(x), \quad (8.2)$$

where η_P is the intrinsic parity of the field and A is a 4×4 matrix acting in the spinor space. Both ψ^P and ψ satisfy the Dirac equation. We can determine A , starting with the Dirac equation for $\psi(x)$,

$$(i\gamma^\mu \partial_\mu - M)\psi(x) = 0.$$

Table 8.1: *The behavior of classical quantities under P, T, and C*

quantity	P	T	C
t	t	-t	t
\mathbf{r}	$-\mathbf{r}$	\mathbf{r}	\mathbf{r}
x^μ	$\tilde{x}^\mu \equiv x_\mu$	$-\tilde{x}^\mu$	x^μ
E	E	E	E
\mathbf{p}	$-\mathbf{p}$	$-\mathbf{p}$	\mathbf{p}
p^μ	\tilde{p}^μ	\tilde{p}^μ	p^μ
\mathbf{L}	\mathbf{L}	$-\mathbf{L}$	\mathbf{L}
\mathbf{s}	\mathbf{s}	$-\mathbf{s}$	\mathbf{s}
$\lambda = \mathbf{s} \cdot \hat{\mathbf{p}}$	$-\lambda$	λ	λ

After parity transforming x to \tilde{x} the Dirac equation becomes after some manipulations

$$\begin{aligned} (i\gamma^\mu \tilde{\partial}_\mu - M) \psi(\tilde{x}) &= 0, \\ (i\tilde{\gamma}^\mu \partial_\mu - M) \psi(\tilde{x}) &= 0, \\ (i\gamma^{\mu\dagger} \partial_\mu - M) \psi(\tilde{x}) &= 0, \\ (i\gamma^\mu \partial_\mu - M) \gamma_0 \psi(\tilde{x}) &= 0. \end{aligned} \quad (8.3)$$

Therefore $\gamma_0 \psi(\tilde{x})$ is again a solution of the Dirac equation and we have

$$\psi^P(x) = \gamma_0 \psi(\tilde{x}). \quad (8.4)$$

It is straightforward to apply this to the explicit field operator $\psi(x)$ using

$$\gamma_0 u(k, m) = u(\tilde{k}, m), \quad (8.5)$$

$$\gamma_0 v(k, m) = -v(\tilde{k}, m), \quad (8.6)$$

(check this for the standard representation; if helicity λ is used instead of the z-component of the spin m , the above operation reverses the sign of λ). The result is

$$\begin{aligned} \psi^P(x) &= P_{op} \psi(x) P_{op}^{-1} = \eta_P \gamma_0 \psi(\tilde{x}) \quad (8.7) \\ &= \sum_\lambda \int \frac{d^3 k}{(2\pi)^3 2E_k} \eta_P [b(k, \lambda) \gamma_0 u(k, \lambda) e^{-ik \cdot \tilde{x}} + d^\dagger(k, \lambda) \gamma_0 v(k, \lambda) e^{ik \cdot \tilde{x}}] \\ &= \sum_\lambda \int \frac{d^3 k}{(2\pi)^3 2E_k} \eta_P [b(k, \lambda) u(\tilde{k}, -\lambda) e^{-i\tilde{k} \cdot x} - d^\dagger(k, \lambda) v(\tilde{k}, -\lambda) e^{i\tilde{k} \cdot x}] \\ &= \sum_\lambda \int \frac{d^3 \tilde{k}}{(2\pi)^3 2E_{\tilde{k}}} \eta_P [b(k, \lambda) u(\tilde{k}, -\lambda) e^{-i\tilde{k} \cdot x} - d^\dagger(k, \lambda) v(\tilde{k}, -\lambda) e^{i\tilde{k} \cdot x}] \\ &= \sum_\lambda \int \frac{d^3 k}{(2\pi)^3 2E_k} \eta_P [b(\tilde{k}, -\lambda) u(k, \lambda) e^{-ik \cdot x} - d^\dagger(\tilde{k}, -\lambda) v(k, \lambda) e^{ik \cdot x}]. \end{aligned} \quad (8.8)$$

From this one sees immediately that

$$P_{op} b(k, \lambda) P_{op}^{-1} = \eta_P b(\tilde{k}, -\lambda), \quad (8.9)$$

$$P_{op} d(k, \lambda) P_{op}^{-1} = -\eta_P^* d(\tilde{k}, -\lambda), \quad (8.10)$$

i.e. choosing η_P is real ($\eta_P = \pm 1$) particle and antiparticle have opposite parity.

In the same way as the Fermion field, one can also consider the scalar field and vector fields. For the scalar field we have seen

$$\phi(x) \longrightarrow P_{op} \phi(x) P_{op}^{-1} = \eta_P \phi(\tilde{x}), \quad (8.11)$$

and for the vector field

$$A^\mu(x) \longrightarrow P_{op} A^\mu(x) P_{op}^{-1} = -A_\mu(\tilde{x}). \quad (8.12)$$

The latter behavior of the vector field will be discussed further below.

8.2 Charge conjugation

We have already seen the particle-antiparticle symmetry with under what we will call charge conjugation the behavior

$$\psi(x) \longrightarrow \psi^c(x) = \eta_C C \bar{\psi}^T(x), \quad (8.13)$$

the latter being also a solution of the Dirac equation. The action on the spinors (using $C = i\gamma^2\gamma^0 = -i\rho^1\sigma^2$ in standard representation) gives

$$C \bar{u}^T(k, m) = v(k, m), \quad (8.14)$$

$$C \bar{v}^T(k, m) = u(k, m), \quad (8.15)$$

(where one must be aware of the choice of spinors made in the expansion, as discussed in section 4). The same relations hold for helicity states. Therefore

$$\psi^c(x) = C_{op} \psi(x) C_{op}^{-1} = \eta_C C \bar{\psi}^T(x) \quad (8.16)$$

$$\begin{aligned} &= \sum_{\lambda} \int \frac{d^3k}{(2\pi)^3 2E_k} \eta_C [d(k, \lambda) C \bar{v}^T(k, \lambda) e^{-ik \cdot x} + b^\dagger(k, \lambda) C \bar{u}^T(k, \lambda) e^{ik \cdot x}] \\ &= \sum_{\lambda} \int \frac{d^3k}{(2\pi)^3 2E_k} \eta_C [d(k, \lambda) u(k, \lambda) e^{-ik \cdot x} + b^\dagger(k, \lambda) v(k, \lambda) e^{ik \cdot x}]. \end{aligned} \quad (8.17)$$

This shows that

$$C_{op} b(k, \lambda) C_{op}^{-1} = \eta_C d(k, \lambda), \quad (8.18)$$

$$C_{op} d(k, \lambda) C_{op}^{-1} = \eta_C^* b(k, \lambda). \quad (8.19)$$

8.3 Time reversal

The time reversal operator transforms

$$x^\mu = (t, \mathbf{r}) \longrightarrow -\tilde{x}^\mu \equiv -x_\mu = (-t, \mathbf{r}). \quad (8.20)$$

We will again consider the transformation properties for a fermion field $\psi(x)$, writing

$$\psi(x) \longrightarrow T_{op} \psi(x) T_{op}^{-1} = \eta_T A \psi(-\tilde{x}) \equiv \psi^t(-\tilde{x}), \quad (8.21)$$

where A is a 4×4 matrix acting in the spinor space. As time reversal will transform 'bra' into 'ket', $T_{op}|\phi\rangle = \langle\phi^t| = (|\phi^t\rangle)^*$, it is antilinear¹. Norm conservation requires T_{op} to be anti-unitary². For a quantized field one has

$$T_{op} f_k(x) b_k T_{op}^{-1} = f_k^*(x) T_{op} b_k T_{op}^{-1},$$

i.e. to find $\psi^t(-\tilde{x})$ that is a solution of the Dirac equation, we start with the complex conjugated Dirac equation for ψ ,

$$((i\gamma^\mu)^* \partial_\mu - M) \psi(x) = 0.$$

The (time-reversed) Dirac equation becomes,

$$\begin{aligned} &(-i\gamma^\mu)^* \tilde{\partial}_\mu - M) \psi(-\tilde{x}) = 0, \\ &(i\tilde{\gamma}^{\mu*} \partial_\mu - M) \psi(-\tilde{x}) = 0, \\ &(i\gamma^{\mu T} \partial_\mu - M) \psi(-\tilde{x}) = 0, \\ &(-iC^{-1} \gamma^\mu C \partial_\mu - M) \psi(-\tilde{x}) = 0. \\ &(i(\gamma_5 C)^{-1} \gamma^\mu \gamma_5 C \partial_\mu - M) \psi(-\tilde{x}) = 0. \\ &(i\gamma^\mu \partial_\mu - M) \gamma_5 C \psi(-\tilde{x}) = 0. \end{aligned} \quad (8.22)$$

¹A is antilinear if $A(\lambda|\phi\rangle + \mu|\psi\rangle) = \lambda^* A|\phi\rangle + \mu^* A|\psi\rangle$.

²An antilinear operator is anti-unitary if $A^\dagger = A^{-1}$. One has $\langle A\phi|A\psi\rangle = \langle\phi|\psi\rangle^* = \langle A\psi|A\phi\rangle^* = \langle\psi|A^\dagger A\phi\rangle = \langle\psi|\phi\rangle$.

Table 8.2: *The transformation properties of physical states for particles (a) and antiparticles (\bar{a}).*

state	P	T	C
$ a; \mathbf{p}, \lambda\rangle$	$ a; -\mathbf{p}, -\lambda\rangle$	$\langle a; -\mathbf{p}, \lambda $	$ \bar{a}; \mathbf{p}, \lambda\rangle$
$ \bar{a}; \mathbf{p}, \lambda\rangle$	$ \bar{a}; -\mathbf{p}, -\lambda\rangle$	$\langle \bar{a}; -\mathbf{p}, \lambda $	$ a; \mathbf{p}, \lambda\rangle$

Therefore $\gamma_5 C \psi(-\tilde{x})$ is again a solution of the (ordinary) Dirac equation and we can choose (phase is convention)

$$\psi^t(x) = i \gamma_5 C \psi(-\tilde{x}). \quad (8.23)$$

In the standard representation $i \gamma_5 C = \sigma_2$ and it is straightforward to apply this to the explicit field operator $\psi(x)$ using

$$i \gamma_5 C u(k, \lambda) = u^*(\tilde{k}, \lambda), \quad (8.24)$$

$$i \gamma_5 C v(k, \lambda) = v^*(\tilde{k}, \lambda), \quad (8.25)$$

(check this for the standard representation). The result is

$$\begin{aligned} \psi^t(x) &= T_{op} \psi(x) T_{op}^{-1} = i \eta_T \gamma_5 C \psi(-\tilde{x}) \quad (8.26) \\ &= \sum_{\lambda} \int \frac{d^3 k}{(2\pi)^3 2E_k} \eta_T [b(k, \lambda) i \gamma_5 C u(k, \lambda) e^{ik \cdot \tilde{x}} + d^\dagger(k, \lambda) i \gamma_5 C v(k, \lambda) e^{-ik \cdot \tilde{x}}] \\ &= \sum_{\lambda} \int \frac{d^3 k}{(2\pi)^3 2E_k} \eta_T [b(k, \lambda) u^*(\tilde{k}, \lambda) e^{i\tilde{k} \cdot x} + d^\dagger(k, \lambda) v^*(\tilde{k}, \lambda) e^{-i\tilde{k} \cdot x}] \\ &= \sum_{\lambda} \int \frac{d^3 \tilde{k}}{(2\pi)^3 2E_k} \eta_T [b(k, \lambda) u^*(\tilde{k}, \lambda) e^{i\tilde{k} \cdot x} + d^\dagger(k, \lambda) v^*(\tilde{k}, \lambda) e^{-i\tilde{k} \cdot x}] \\ &= \sum_{\lambda} \int \frac{d^3 k}{(2\pi)^3 2E_k} \eta_T [b(\tilde{k}, \lambda) u^*(k, \lambda) e^{ik \cdot x} + d^\dagger(\tilde{k}, \lambda) v^*(k, \lambda) e^{-ik \cdot x}] \quad (8.27) \end{aligned}$$

From this one obtains

$$T_{op} b(k, \lambda) T_{op}^{-1} = \eta_T b(\tilde{k}, \lambda), \quad (8.28)$$

$$T_{op} d(k, \lambda) T_{op}^{-1} = \eta_T^* d(\tilde{k}, \lambda). \quad (8.29)$$

In table 2 the behavior of particle states under the various transformations has been summarized. Note that applying an anti-unitary transformation such as T_{op} one must take for the matrix element the complex conjugate. Therefore one has $\langle A|k\rangle = \langle A|b^\dagger(k)|0\rangle = \langle A|T^\dagger T b^\dagger(k) T^\dagger T|0\rangle^* = \langle A^t|b^\dagger(\tilde{k})|0\rangle^* = \langle 0|b(\tilde{k})|A^t\rangle = \langle \tilde{k}|A^t\rangle$.

8.4 Bi-linear combinations

In quantities such as currents and lagrangians often bilinear combinations of spinor fields are encountered. Since there are 16 independent 4×4 matrices, there are 16 independent of these bilinear combinations. They are the following

$$S(x) = \bar{\psi}(x)\psi(x) \quad (\text{scalar}) \quad (8.30)$$

$$V^\mu(x) = \bar{\psi}(x)\gamma^\mu\psi(x) \quad (\text{vector}) \quad (8.31)$$

$$T^{\mu\nu}(x) = \bar{\psi}(x)\sigma^{\mu\nu}\psi(x) \quad (\text{tensor}) \quad (8.32)$$

$$A^\mu(x) = \bar{\psi}(x)\gamma_5\gamma^\mu\psi(x) \quad (\text{axial vector}) \quad (8.33)$$

$$P(x) = i\bar{\psi}(x)\gamma_5\psi(x) \quad (\text{pseudoscalar}) \quad (8.34)$$

Table 8.3: The behavior of the independent bilinear combinations of fermi fields under P, C, and T

	P	C	T	$\Theta = \text{PCT}$
$S(x)$	$S(\tilde{x})$	$S(x)$	$S(-\tilde{x})$	$S(-x)$
$V^\mu(x)$	$V_\mu(\tilde{x})$	$-V^\mu(x)$	$V_\mu(-\tilde{x})$	$-V^\mu(-x)$
$T^{\mu\nu}(x)$	$T_{\mu\nu}(\tilde{x})$	$-T^{\mu\nu}(x)$	$-T_{\mu\nu}(-\tilde{x})$	$T^{\mu\nu}(-x)$
$A^\mu(x)$	$-A_\mu(\tilde{x})$	$A^\mu(x)$	$A_\mu(-\tilde{x})$	$-A^\mu(-x)$
$P(x)$	$-P(\tilde{x})$	$P(x)$	$-P(-\tilde{x})$	$P(-x)$

The matrix $\sigma^{\mu\nu} \equiv (i/2)[\gamma^\mu, \gamma^\nu]$. The 16 combinations of Dirac matrices appearing above are linearly independent. Applying the results from the previous sections it is straightforward to determine the behavior of the combinations under P, C, and T, as well as under the combined operation $\Theta = \text{PCT}$ (see Table 3). As the coupling of the photon field to fermions is given by an interaction term in the lagrangian of the form $:\bar{\psi}(x)\gamma^\mu\psi(x): A_\mu(x)$ and behaves as a scalar one sees immediately that the photon field $A^\mu(x)$ behaves in the same way as the vector combination $\bar{\psi}(x)\gamma^\mu\psi(x)$. Note that the lagrangian density $\mathcal{L}(x) \rightarrow \mathcal{L}(-x)$ under Θ .

8.5 Form factors

Currents play an important role in field theory. In many applications the expectation values of currents are needed, e.g. for the vector current $V^\mu(x)$,

$$\langle p', s' | V^\mu(x) | p, s \rangle. \quad (8.35)$$

The x -dependence can be accounted for straightforwardly using translation invariance, $V^\mu(x) = e^{i P_{op} \cdot x} V^\mu(0) e^{-i P_{op} \cdot x}$. This implies

$$\langle p' | V^\mu(x) | p \rangle = e^{-i(p-p') \cdot x} \langle p' | V^\mu(0) | p \rangle. \quad (8.36)$$

As an example consider the vector current for a point fermion,

$$V^\mu(x) =: \bar{\psi}(x)\gamma^\mu\psi(x) :, \quad (8.37)$$

of which the expectation value between momentum eigenstates can be simply found,

$$\langle p' | V^\mu(0) | p \rangle = \bar{u}(p')\gamma^\mu u(p). \quad (8.38)$$

In general the expectation value between momentum states can be more complicated,

$$\langle p' | V^\mu(x) | p \rangle = \bar{u}(p')\Gamma^\mu(p', p)u(p) e^{-i(p-p') \cdot x}, \quad (8.39)$$

where $\Gamma^\mu(p', p)$ can be built from any combination of Dirac matrices ($1, \gamma^\mu, \sigma^{\mu\nu}, \gamma_5\gamma^\mu$ or γ_5), momenta (p^μ or p'^μ) or constant tensors ($\epsilon_{\mu\nu\rho\sigma}$ or $g_{\mu\nu}$). For instance for nucleons one has

$$\Gamma_\mu(p', p) = \gamma_\mu F_1(q^2) + \frac{i\sigma_{\mu\nu}q^\nu}{2M} F_2(q^2), \quad (8.40)$$

where the coefficients are functions of invariants constructed out of the momenta, in this case only $q^2 = (p' - p)^2$, called *form factors*. There are several other terms that also have the correct tensorial behavior such as

$$q^\mu F_3(q^2), \quad \frac{(p^\mu + p'^\mu)}{2M} F_4(q^2), \quad \frac{\gamma_5 \sigma^{\mu\nu} q_\nu}{2M} F_5(q^2), \quad (8.41)$$

but that are eliminated because of relations between Dirac matrices, e.g. $\gamma_5 \sigma_{\mu\nu} = (1/2)\epsilon_{\mu\nu\rho\sigma}\gamma^\rho\gamma^\sigma$, or relations that follow from the equations of motion, e.g. the Gordon decomposition

$$\bar{u}(p')\gamma^\mu u(p) = \frac{1}{2M}\bar{u}(p')[(p' + p)^\mu + i\sigma^{\mu\nu}q_\nu]u(p), \quad (8.42)$$

where $q = p' - p$, or relations based on hermiticity of the operator or P , C and T invariance.

- **Hermiticity**

$$\begin{aligned} \langle p'|V^\mu(0)|p\rangle &= \bar{u}(p')\Gamma^\mu(p', p)u(p) \\ &= \langle p'|V^{\mu\dagger}(0)|p\rangle = \langle p|V^\mu(0)|p'\rangle^* \\ &= (\bar{u}(p)\Gamma^\mu(p, p')u(p'))^* \\ &= (u^\dagger(p)\gamma_0\Gamma^\mu(p, p')u(p'))^\dagger \\ &= u^\dagger(p')\Gamma^{\mu\dagger}(p, p')\gamma_0u(p). \end{aligned}$$

Therefore hermiticity implies for the structure of $\Gamma^\mu(p', p)$ that

$$\gamma_0\Gamma^{\mu\dagger}(p, p')\gamma_0 = \Gamma^\mu(p', p). \quad (8.43)$$

- **Parity**

$$\begin{aligned} \langle p'|V^\mu(x)|p\rangle &= e^{i q \cdot x} \bar{u}(p')\Gamma^\mu(p', p)u(p) \\ &= \langle p'|P_{op}^\dagger P_{op} V^\mu(x) P_{op}^\dagger P_{op}|p\rangle \\ &= \langle \tilde{p}'|V_\mu(\tilde{x})|\tilde{p}\rangle = e^{i \tilde{q} \cdot \tilde{x}} \bar{u}(\tilde{p}')\Gamma_\mu(\tilde{p}', \tilde{p})u(\tilde{p}) \\ &= e^{i q \cdot x} \bar{u}(p')\gamma_0\Gamma_\mu(p', \tilde{p})\gamma_0u(p). \end{aligned}$$

Therefore parity invariance implies for the structure of $\Gamma^\mu(p', p)$ that

$$\Gamma^\mu(p', p) = \gamma_0\Gamma_\mu(\tilde{p}', \tilde{p})\gamma_0, \quad (8.44)$$

- **Time reversal**

$$\begin{aligned} \langle p'|V^\mu(x)|p\rangle &= e^{i q \cdot x} \bar{u}(p')\Gamma^\mu(p', p)u(p) \\ &= \langle p'|T_{op}^\dagger T_{op} V^\mu(x) T_{op}^\dagger T_{op}|p\rangle^* \\ &= \langle \tilde{p}'|V_\mu(-\tilde{x})|\tilde{p}\rangle^* = e^{i \tilde{q} \cdot \tilde{x}} [\bar{u}(\tilde{p}')]^* \Gamma_\mu^*(\tilde{p}', \tilde{p}) u^*(\tilde{p}) \\ &= e^{i q \cdot x} \bar{u}(p') (i\gamma_5 C) \Gamma_\mu^*(\tilde{p}', \tilde{p}) (i\gamma_5 C) u(p). \end{aligned}$$

Therefore time reversal invariance implies for the structure of $\Gamma^\mu(p', p)$ that

$$\Gamma^\mu(p', p) = (i\gamma_5 C) \Gamma_\mu^*(\tilde{p}', \tilde{p}) (i\gamma_5 C). \quad (8.45)$$

Exercises

Exercise 8.1

The matrix element of the electromagnetic current between nucleon states is written as

$$\langle p'|J_\mu(x)|p\rangle = e^{-i(p-p') \cdot x} \bar{u}(p')\Gamma_\mu u(p)$$

Here

$$\begin{aligned} \Gamma_\mu &= \gamma_\mu F_1(q^2) + \frac{i\sigma_{\mu\nu}q^\nu}{2M} F_2(q^2) \\ &= \gamma_\mu H_1(q^2) - \frac{p_\mu + p'_\mu}{2M} H_2(q^2) \end{aligned}$$

- (a) Use the Dirac equation to prove the Gordon-decomposition

$$\bar{u}(p')\gamma^\mu u(p) = \frac{1}{2M}\bar{u}(p')[(p' + p)^\mu + i\sigma^{\mu\nu}(p' - p)_\nu]u(p)$$

- (b) Give the relation between H_i en F_i .
 (c) Show that hermiticity of the current requires that F_1 en F_2 are real.

Exercise 8.2

- (a) Calculate the current matrix element (using the explicit fermion spinors) in the Breit-frame in which $q = (0, 0, 0, |\mathbf{q}|)$, $p = (E, 0, 0, -|\mathbf{q}|/2)$, $p' = (E, 0, 0, |\mathbf{q}|/2)$ with $E^2 = M^2 + |\mathbf{q}|^2/4$ and express them in terms of the Sachs form factors depending on $Q^2 = -q^2$,

$$\begin{aligned} G_E &= F_1 - \frac{Q^2}{4M^2}F_2, \\ G_M &= F_1 + F_2. \end{aligned}$$

- (b) Show that in the interaction with the electromagnetic field,

$$\mathcal{L}_{int} = e j_\mu A^\mu$$

the charge is given by $e G_E(0)$ and the magnetic moment by $e G_M(0)/2M$.

Exercise 8.3

- (a) Show that because of parity conservation no term of the form

$$\gamma^5 \frac{\sigma_{\mu\nu} q^\nu}{2M} F_3(q^2)$$

can appear in the current matrix element.

- (b) Show that such a term is also not allowed by time-reversal symmetry.
 (c) Show that if a term of this form would exist, it would correspond to an electric dipole moment $d = -e F_3(0)/2M$ (Interaction term $d \sigma \cdot \mathbf{E}$).

Chapter 9

Path integrals and quantum mechanics

9.1 Time evolution as path integral

The time evolution from $t_0 \rightarrow t$ of a quantum mechanical system is generated by the Hamiltonian,

$$U(t, t_0) = e^{-i(t-t_0)H} \quad (9.1)$$

or

$$i\frac{\partial}{\partial t}U(t, t_0) = HU(t, t_0) \quad (9.2)$$

Two situations can be distinguished:

- (i) *Schrödinger picture*, in which the operators are time-independent, $A_S(t) = A_S$ and the states are time dependent, $|\psi_S(t)\rangle = U(t, t_0)|\psi_S(t_0)\rangle$,

$$i\frac{\partial}{\partial t}|\psi_S\rangle = H|\psi_S\rangle, \quad (9.3)$$

$$i\frac{\partial}{\partial t}A_S \equiv 0. \quad (9.4)$$

- (ii) *Heisenberg picture*, in which the states are time-independent, $|\psi_H(t)\rangle = |\psi_H\rangle$, and the operators are time-dependent, $A_H(t) = U^{-1}(t, t_0)A_H(t_0)U(t, t_0)$,

$$i\frac{\partial}{\partial t}|\psi_H\rangle \equiv 0, \quad (9.5)$$

$$i\frac{\partial}{\partial t}A_H = [A_H, H]. \quad (9.6)$$

Of these the Heisenberg picture is most appropriate for quantum field theory since the field operators do depend on the position and one would like to have position and time on the same footing.

Consider the two (time-independent) Heisenberg states:

$$\begin{aligned} |q, t\rangle & \quad Q_H(t)|q, t\rangle \equiv q|q, t\rangle, \\ |q', t'\rangle & \quad Q_H(t')|q', t'\rangle \equiv q'|q', t'\rangle, \end{aligned} \quad (9.7)$$

and Schrödinger states

$$\begin{aligned} |q\rangle & Q_S |q\rangle \equiv q|q\rangle, \\ |q'\rangle & Q_S |q'\rangle \equiv q'|q'\rangle. \end{aligned} \quad (9.8)$$

Choose t as the starting point with $|q\rangle = |q, t\rangle$ and $Q_H(t) = Q_S$ and study the evolution of the system by calculating the quantum mechanical overlap amplitude

$$\langle q', t' | q, t \rangle = \langle q' | e^{-iH(t'-t)} | q \rangle. \quad (9.9)$$

Dividing the interval from $t \equiv t_0$ to $t' \equiv t_n$ into n pieces of length $\Delta\tau$ and using completeness (at each time t_i) one writes

$$\begin{aligned} \langle q', t' | q, t \rangle &= \langle q' | e^{-inH\Delta\tau} | q \rangle = \langle q' | (e^{-iH\Delta\tau})^n | q \rangle \\ &= \int dq_1 \int \dots \int dq_{n-1} \langle q' | e^{-iH\Delta\tau} | q_{n-1} \rangle \langle q_{n-1} | \dots | q_1 \rangle \langle q_1 | e^{-iH\Delta\tau} | q \rangle. \end{aligned} \quad (9.10)$$

The purpose of this is to calculate the evolution for an infinitesimal time interval. The hamiltonian is an operator $H = H(P, Q)$ expressed in terms of the operators P and Q . These can be written in coordinate or momentum representation as

$$Q = \int dq |q\rangle q \langle q| \quad (9.11)$$

$$P = \int dq |q\rangle \left(-i\frac{\partial}{\partial q}\right) \langle q| = \int \frac{dp}{2\pi} |p\rangle p \langle p|, \quad (9.12)$$

where the transformation between coordinate and momentum space involves

$$\langle q | p \rangle = e^{ip \cdot q}. \quad (9.13)$$

At least for a simple hamiltonian such as consisting of a kinetic energy term and a local potential, $H(P, Q) = K(P) + V(Q) = (P^2/2M) + V(Q)$ one can split $e^{-iH\Delta\tau} \approx e^{-iK\Delta\tau} e^{-iV\Delta\tau}$, with the correction¹ being of order $(\Delta\tau)^2$. Then

$$\begin{aligned} \langle q_{j+1}, t_{j+1} | q_j, t_j \rangle &\approx \langle q_{j+1} | e^{-iK\Delta\tau} e^{-iV\Delta\tau} | q_j \rangle \\ &= \int \frac{dp_j}{2\pi} \langle q_{j+1} | p_j \rangle \langle p_j | e^{-iK\Delta\tau} e^{-iV\Delta\tau} | q_j \rangle. \end{aligned} \quad (9.15)$$

By letting the kinetic and potential parts act to left and right respectively one can express the expectation values in terms of integrals containing $H(p_j, q_j)$, which is a hamiltonian function in which the operators are replaced by real-numbered variables. Combining the two terms gives

$$\begin{aligned} \langle q_{j+1}, t_{j+1} | q_j, t_j \rangle &= \int \frac{dp_j}{2\pi} e^{ip_j(q_{j+1}-q_j)} e^{-iH(p_j, q_j)\Delta\tau} \\ &= \int \frac{dp_j}{2\pi} e^{ip_j(q_{j+1}-q_j) - i\Delta\tau H(p_j, q_j)} \\ &= \int \frac{dp_j}{2\pi} \exp(i\Delta\tau [p_j \dot{q}_j - H(p_j, q_j)]), \end{aligned} \quad (9.16)$$

¹For this, use the Campbell-Baker-Hausdorff formula,

$$e^A e^B = e^C \quad \text{with} \quad C = A + B + \frac{1}{2}[A, B] + \frac{1}{12}[A, [A, B]] + \frac{1}{12}[[A, B], B] + \dots \quad (9.14)$$

or for the full interval

$$\begin{aligned}\langle q', t' | q, t \rangle &= \int \prod_{\tau} \frac{dq(\tau) dp(\tau)}{2\pi} \exp \left(i \sum_t^{t'} \Delta\tau [p\dot{q} - H(p, q)] \right) \\ &= \int \mathcal{D}q \mathcal{D}p \exp \left(i \int_t^{t'} d\tau [p\dot{q} - H(p, q)] \right),\end{aligned}\quad (9.17)$$

where $\mathcal{D}q$ and $\mathcal{D}p$ indicate functional integrals. The importance of this expression is that it expresses a quantum mechanical amplitude as a *path integral* with in the integrand a classical hamiltonian function.

Before proceeding we also give the straightforward extension to more than one degree of freedom,

$$\begin{aligned}\langle q'_1, \dots, q'_N, t' | q_1, \dots, q_N, t \rangle & \\ = \int \left(\prod_{n=1}^N \mathcal{D}q_n \mathcal{D}p_n \right) \exp \left(i \int_t^{t'} d\tau \left[\sum_n p_n \dot{q}_n - H(p_1, q_1, \dots, p_N, q_N) \right] \right).\end{aligned}\quad (9.18)$$

9.2 Functional integrals

In this section I want to give a fairly heuristic discussion of functional integrals. What one is after is the meaning of

$$\int \mathcal{D}\alpha F[\alpha], \quad (9.19)$$

where $F[\alpha]$ is a functional that represents a mapping from a function space \mathcal{F} of (real) functions α ($R \rightarrow R$) into the real numbers R , i.e. $F[\alpha] \in R$. Examples are

$$\begin{aligned}F_1[\alpha] &= \alpha \equiv \int dx \alpha(x), \\ F_2[\alpha] &= \alpha^2 \equiv \int dx \alpha^2(x), \\ F_3[\alpha, \beta] &= \alpha K \beta \equiv \int dx dy \alpha(x) K(x, y) \beta(y), \\ F_4[\alpha] &= \exp \left(-\frac{1}{2} \alpha^2 \right) \equiv \exp \left(-\frac{1}{2} \int dx \alpha^2(x) \right), \\ F_5[\alpha, \alpha^*] &= \exp(-\alpha^* K \alpha) \equiv \exp \left(-\int dx dy \alpha^*(x) K(x, y) \alpha(y) \right).\end{aligned}$$

The kernels $K(x, y)$ in the above examples can be (hermitean) operators acting on the functions, including differential operators, etc.

Two working approaches for functional integration are the following:

- (i) the heuristic division of the space on which α acts into cells, i.e. $\alpha(x) \rightarrow \alpha_x$ and $F[\alpha] = F(\alpha_x)$ changes into a multivariable function, while

$$\int \mathcal{D}\alpha F[\alpha] = \int \left(\prod_x \frac{d\alpha_x}{N} \right) F(\alpha_x) \quad (9.20)$$

becomes a multidimensional integral.

- (ii) Write α in terms of a sum of orthonormal basis functions, $\alpha(x) = \sum_n \alpha_n f_n$ and $K(x, y) = \sum_{m,n} f_m(x) K_{mn} f_n^*(y)$ and consider $F[\alpha] = F(\alpha_n)$ as a multivariable function, with

$$\int \mathcal{D}\alpha F[\alpha] = \int \left(\prod_n \frac{d\alpha_n}{N} \right) F(\alpha_n) \quad (9.21)$$

again a multidimensional integral. In both cases N is an appropriately chosen normalization constant in such a way that the integral is finite. Note that procedure (i) is an example of the more general procedure under (ii).

Consider the gaussian functional as an example. Using method 1 one writes

$$\begin{aligned} \int \mathcal{D}\alpha \exp\left(-\frac{1}{2}\alpha^2\right) &= \int \left(\prod_x \frac{d\alpha_x}{N} \right) \exp\left(-\frac{1}{2} \sum_x \Delta x \alpha_x^2\right) \\ &= \prod_x \int \frac{d\alpha_x}{N} \exp\left(-\frac{1}{2} \Delta x \alpha_x^2\right) \\ &= \prod_x \left(\frac{1}{N} \sqrt{\frac{2\pi}{\Delta x}} \right) \equiv 1. \end{aligned} \quad (9.22)$$

The last equality is obtained by defining the right measure (normalization N) in the integration. Physical answers will usually come out as the ratio of two functional integrals and are thus independent of the chosen measure. Using the second method and expanding in a basis set of functions one obtains for the (real) gaussian functional

$$\begin{aligned} \int \mathcal{D}\alpha \exp\left(-\frac{1}{2}\alpha^2\right) &= \int \left(\prod_n \frac{d\alpha_n}{N} \right) \exp\left(-\frac{1}{2} \sum_{m,n} \alpha_m \alpha_n \int dx f_m(x) f_n(x)\right) \\ &= \prod_n \int \frac{d\alpha_n}{N} \exp\left(-\frac{1}{2}\alpha_n^2\right) \\ &= \prod_n \left(\frac{\sqrt{2\pi}}{N} \right) \equiv 1. \end{aligned} \quad (9.23)$$

Having defined the Gaussian integral, the following integrals can be derived for a real symmetric or complex hermitean kernel K ,

$$\int \mathcal{D}\alpha \exp\left(-\frac{1}{2} \alpha K \alpha\right) = \frac{1}{\sqrt{\det K}} \quad (\alpha \text{ real-valued}) \quad (9.24)$$

$$\int \mathcal{D}\alpha \mathcal{D}\alpha^* \exp(-\alpha^* K \alpha) = \frac{1}{\det K} \quad (\alpha \text{ complex-valued}) \quad (9.25)$$

(see Exercises). A useful property of functional integration is the translation invariance,

$$\int \mathcal{D}\alpha F[\alpha] = \int \mathcal{D}\alpha F[\alpha + \beta]. \quad (9.26)$$

As an important application of translation invariance we mention the identities

$$\int \mathcal{D}\alpha \exp\left(-\frac{1}{2}\alpha^2 - \omega\alpha\right) = \exp\left(\frac{1}{2}\omega^2\right), \quad (9.27)$$

$$\int \mathcal{D}\alpha \mathcal{D}\alpha^* \exp(-\alpha^* \alpha - \alpha^* \omega - \omega^* \alpha) = \exp(\omega^* \omega), \quad (9.28)$$

Functional differentiation is defined as

$$\left[\frac{\delta}{\delta\alpha(x)}, \alpha(y) \right] = \delta(x-y), \quad (9.29)$$

or in discretized form

$$\frac{\delta}{\delta\alpha(x)} \longrightarrow \frac{1}{\Delta x} \frac{\partial}{\partial\alpha_x} \quad \text{with} \quad \left[\frac{\partial}{\partial\alpha_x}, \alpha_y \right] = \delta_{xy}, \quad (9.30)$$

$$\frac{\delta}{\delta\alpha(x)} \longrightarrow \sum_n f_n^*(x) \frac{\partial}{\partial\alpha_n} \quad \text{with} \quad \left[\frac{\partial}{\partial\alpha_m}, \alpha_n \right] = \delta_{mn}. \quad (9.31)$$

Note that $\delta(x-y) = \sum_n f_n(x) f_n^*(y)$. Examples of functional differentiation are

$$\frac{\delta}{\delta\alpha(x)} \alpha = \frac{\delta}{\delta\alpha(x)} \int dy \alpha(y) = \int dy \delta(x-y) = 1, \quad (9.32)$$

$$\frac{\delta}{\delta\alpha(x)} \alpha K \beta = \int dy K(x,y) \beta(y) \quad (9.33)$$

$$\frac{\delta}{\delta\alpha(x)} \exp\left(-\frac{1}{2}\alpha^2\right) = -\alpha(x) \exp\left(-\frac{1}{2}\alpha^2\right), \quad (9.34)$$

$$\frac{\delta}{\delta\alpha(x)} \exp(-\alpha\beta) = -\beta(x) \exp(-\alpha\beta). \quad (9.35)$$

For applications to fermion fields, we need to consider Grassmann-valued functions involving anticommuting Grassmann variables, i.e. $\theta\eta = -\eta\theta$, $\theta^2 = 0$. In principle the definitions of functionals is the same, e.g. the Gaussian-type functionals,

$$F[\theta, \theta^*] = \exp(-\theta^* \theta) \equiv \exp\left(-\int dx \theta^*(x) \theta(x)\right),$$

$$F[\theta, \theta^*] = \exp(-\theta^* K \theta) \equiv \exp\left(-\int dx dy \theta^*(x) K(x,y) \theta(y)\right).$$

Integration for Grassmann variables is defined as

$$\int d\theta \, 1 = 0 \quad \text{and} \quad \int d\theta \, \theta = 1. \quad (9.36)$$

This gives for the Gaussian functional integral after expanding $\theta(x) = \sum_n \theta_n f_n(x)$ where the coefficients θ_n are Grassmann variables the result

$$\begin{aligned} \int \mathcal{D}\theta^* \mathcal{D}\theta \exp(-\theta^* \theta) &= \int \left(\prod_n d\theta_n^* d\theta_n \right) \exp\left(-\sum_n \theta_n^* \theta_n\right) \\ &= \prod_n \int d\theta_n^* d\theta_n \exp(-\theta_n^* \theta_n) \\ &= \prod_n \int d\theta_n^* d\theta_n (1 - \theta_n^* \theta_n) = 1. \end{aligned} \quad (9.37)$$

We note that from the first to the second line one needs to realize that a pair of different Grassmann variables behaves as ordinary complex variables. In the expansion of the exponential (from second to third line), however, products of the same pair appear, which vanish. It is now easy to check that

$$\int \mathcal{D}\theta^* \mathcal{D}\theta \exp(-\theta^* K \theta) = \det K. \quad (9.38)$$

Functional differentiation of Grassmann-valued functions is given by

$$\left\{ \frac{\delta}{\delta\theta(x)}, \theta(y) \right\} = \delta(x-y), \quad (9.39)$$

leading for two Grassmann variables $\bar{\theta}$ and η to e.g.

$$\frac{\delta}{\delta\eta(x)} e^{-\theta^*\eta} = \theta^*(x) = \theta^*(x) e^{-\theta^*\eta}, \quad (9.40)$$

$$\frac{\delta}{\delta\eta^*(x)} e^{-\eta^*\theta} = -\theta(x) = -\theta(x) e^{-\eta^*\theta}. \quad (9.41)$$

The translation invariance property and the 'Fourier transform' identity of Gaussian integrals, remains in essence the same, e.g.

$$\int \mathcal{D}\theta^* \mathcal{D}\theta \exp(-\theta^*\theta - \theta^*\eta - \eta^*\theta) = \exp(\eta^*\eta). \quad (9.42)$$

9.3 Time ordered products of operators and path integrals

As an example of working with functional integrals consider the expression for $K(q', t'; q, t)$ for a lagrangian $L(q, \dot{q}) = \frac{1}{2}\dot{q}^2 - V(q)$ and the corresponding hamiltonian $H(p, q) = \frac{1}{2}p^2 + V(q)$. The expression

$$\langle q', t' | q, t \rangle = \int \mathcal{D}q \mathcal{D}\frac{p}{2\pi} \exp\left(i \int_t^{t'} d\tau \left[p\dot{q} - \frac{1}{2}p^2 - V(q) \right]\right), \quad (9.43)$$

can be rewritten after rewriting the integrand as $-\frac{1}{2}(p - \dot{q})^2 + \frac{1}{2}\dot{q}^2 - V(q) = -\frac{1}{2}(p - \dot{q})^2 + L(q, \dot{q})$. The result is

$$\langle q', t' | q, t \rangle = \int \mathcal{D}q \exp\left(i \int_t^{t'} d\tau L(q, \dot{q})\right), \quad (9.44)$$

which was considered the starting point for path integral quantization by Feynman.

Note, however, that not always the $\mathcal{D}p$ integration can be removed that easily. A counter example is the lagrangian $L(q, \dot{q}) = \frac{1}{2}\dot{q}^2 f(q)$ for which $H(p, q) = p^2/[2f(q)]$. As discussed for instance in Ryder the $\mathcal{D}p$ integration can still be removed but one ends up with an effective lagrangian in the path integral

$$\langle q', t' | q, t \rangle = \int \mathcal{D}q \exp\left(i \int_t^{t'} d\tau L_{\text{eff}}(q, \dot{q})\right), \quad (9.45)$$

which is of the form $L_{\text{eff}}(q, \dot{q}) = L(q, \dot{q}) - \frac{i}{2}\delta(0) \ln f(q)$.

Making use of path integrals it is straightforward to calculate the expectation value $\langle q', t' | Q(s) | q, t \rangle$ of an operator $Q(s)$ if $t \leq s \leq t'$. By sandwiching the time s in one of the infinitesimal intervals, $t_j \leq s \leq t_{j+1}$, we have

$$\begin{aligned} \langle q', t' | Q(s) | q, t \rangle &= \int \prod_i dq_i \langle q', t' | q_n, t_n \rangle \dots \\ &\quad \times \langle q_{j+1}, t_{j+1} | Q(s) | q_j, t_j \rangle \dots \langle q_1, t_1 | q, t \rangle. \end{aligned} \quad (9.46)$$

Using $Q(s) | q_j, t_j \rangle = q(s) | q_j, t_j \rangle$, one gets

$$\langle q', t' | Q(s) | q, t \rangle = \int \mathcal{D}q \mathcal{D}\frac{p}{2\pi} q(s) \exp\left(i \int_t^{t'} d\tau [p\dot{q} - H(p, q)]\right) \quad (9.47)$$

$$= \int \mathcal{D}q q(s) \exp\left(i \int_t^{t'} d\tau L(q, \dot{q})\right). \quad (9.48)$$

Defining the time ordered product of operators

$$\mathcal{T} Q(t_1) \dots Q(t_n) \equiv Q(t_{i_1}) \dots Q(t_{i_n}), \quad (9.49)$$

where $t_{i_1} \geq \dots \geq t_{i_n}$ is a permutation of $\{t_1, \dots, t_n\}$, one has

$$\begin{aligned} \langle q', t' | \mathcal{T} Q(t_1) \dots Q(t_n) | q, t \rangle \\ = \int \mathcal{D}q \ q(t_{i_1}) \dots q(t_{i_n}) \exp \left(i \int_t^{t'} d\tau L(q, \dot{q}) \right). \end{aligned} \quad (9.50)$$

Thus not only the quantum mechanical overlap of states can be calculated via a *classical* path integral, but also expectation of operators, at least if they appear time-ordered.

9.4 An application: time-dependent perturbation theory

From quantum mechanics one should be familiar with the procedure of time-dependent perturbation theory, the lowest order leading to Fermi's golden rule. One works in the so-called interaction picture, in which a separation is made of the hamiltonian $H = H_0 + H_I$. The fast time evolution is described in H_0 while H_I is considered as a perturbation. The interaction picture is defined as

$$U_I(t', t) \equiv e^{-i H_0(t'-t)}, \quad (9.51)$$

$$\psi_I(t) \equiv e^{i H_0 t} \psi_S(t) = e^{i H_0 t} e^{-i H t} \psi_S(0) = e^{-i H_I t} \psi_S(0), \quad (9.52)$$

$$A_I(t) \equiv e^{i H_0 t} A_S(t) e^{-i H_0 t} = e^{i H_0 t} A_S e^{-i H_0 t}, \quad (9.53)$$

i.e. if $H_I = 0$ it is the Heisenberg picture and the evolution $\langle q', t' | q, t \rangle$ is described through the operators by H_0 . The evolution of the (interaction) states is described only by H_I ,

$$i \frac{\partial}{\partial t} \psi_I = H_I(t) \psi_I(t), \quad (9.54)$$

$$i \frac{\partial}{\partial t} A_I = [A_I, H_0], \quad (9.55)$$

and

$$\psi_I(t') = U(t', t) \psi_I(t), \quad (9.56)$$

$$i \frac{\partial}{\partial t} U(t', t) = H_I U(t', t) \quad (9.57)$$

with $U(t, t) = 1$ or

$$U(t', t) = 1 - i \int_t^{t'} d\tau H_I(\tau) U(\tau, t), \quad (9.58)$$

which can be solved by iteration, i.e. writing

$$U(t', t) = \sum_{n=0}^{\infty} U^{(n)}(t', t), \quad (9.59)$$

one has

$$\begin{aligned} U^{(0)}(t', t) &= 1 \\ U^{(1)}(t', t) &= 1 - i \int_t^{t'} d\tau H_I(\tau) U^{(0)}(\tau, t) - U^{(0)}(t', t) \\ &= (-i) \int_t^{t'} d\tau H_I(\tau) \end{aligned}$$

$$\begin{aligned}
U^{(2)}(t', t) &= 1 - i \int_t^{t'} d\tau H_I(\tau) \left[U^{(0)}(\tau, t) + U^{(1)}(\tau, t) \right] - \left[U^{(0)}(t', t) + U^{(1)}(t', t) \right] \\
&= (-i) \int_t^{t'} d\tau H_I(\tau) U^{(1)}(\tau, t) \\
&\quad \vdots \\
U^{(n)}(t', t) &= (-i) \int_t^{t'} d\tau H_I(\tau) U^{(n-1)}(\tau, t) \\
&= (-i)^n \int_t^{t'} d\tau_1 \int_t^{\tau_1} d\tau_2 \dots \int_t^{\tau_{n-1}} d\tau_n H_I(\tau_1) \dots H_I(\tau_n) \\
&= \frac{(-i)^n}{n!} \int_t^{t'} d\tau_1 \int_t^{\tau_1} d\tau_2 \dots \int_t^{\tau_{n-1}} d\tau_n \mathcal{T} H_I(\tau_1) \dots H_I(\tau_n). \tag{9.60}
\end{aligned}$$

The last equality is illustrated for the second term $U^{(2)}$ in the following. The integration

$$\int_t^{t'} d\tau_1 \int_t^{\tau_1} d\tau_2 H_I(\tau_1) H_I(\tau_2)$$

can also be performed by first integrating over τ_2 but changing the integration limits (check!). Thus we can write the integration as the sum of the two expressions (multiplying with 1/2),

$$\begin{aligned}
&= \frac{1}{2} \int_t^{t'} d\tau_1 \int_t^{\tau_1} d\tau_2 H_I(\tau_1) H_I(\tau_2) \\
&\quad + \frac{1}{2} \int_t^{t'} d\tau_2 \int_{\tau_2}^{t'} d\tau_1 H_I(\tau_1) H_I(\tau_2),
\end{aligned}$$

Now the integration can be extended by adding theta functions,

$$\begin{aligned}
&= \frac{1}{2} \int_t^{t'} d\tau_1 \int_t^{\tau_1} d\tau_2 H_I(\tau_1) H_I(\tau_2) \theta(\tau_1 - \tau_2) \\
&\quad + \frac{1}{2} \int_t^{t'} d\tau_2 \int_t^{\tau_2} d\tau_1 H_I(\tau_1) H_I(\tau_2) \theta(\tau_1 - \tau_2),
\end{aligned}$$

which can be rewritten (by interchanging in the second term the names of the integration variables)

$$= \frac{1}{2} \int_t^{t'} d\tau_1 \int_t^{\tau_1} d\tau_2 \left[H_I(\tau_1) H_I(\tau_2) \theta(\tau_1 - \tau_2) + H_I(\tau_2) H_I(\tau_1) \theta(\tau_2 - \tau_1) \right],$$

the desired result.

The time evolution operator, therefore, can be written as

$$\begin{aligned}
U(t', t) &= \sum_{n=0}^{\infty} \frac{(-i)^n}{n!} \int_t^{t'} d\tau_1 \int_t^{\tau_1} d\tau_2 \dots \int_t^{\tau_{n-1}} d\tau_n \mathcal{T} H_I(\tau_1) \dots H_I(\tau_n) \\
&\equiv \mathcal{T} \exp \left(-i \int_t^{t'} d\tau H_I(\tau) \right). \tag{9.61}
\end{aligned}$$




Figure 9.1: Physical picture of vacuum to vacuum amplitude in presence of a source term

This is not a surprising result. We can again rewrite the time-ordered products as functional integrals as derived in the previous section,

$$\begin{aligned}
 \langle q', t' | q, t \rangle^V &= \langle q', t' | \mathcal{T} \exp \left(-i \int_t^{t'} d\tau H_I(Q(\tau)) \right) | q, t \rangle^{V=0} \\
 &= \int \mathcal{D}q \mathcal{D}\frac{p}{2\pi} \exp \left(-i \int_t^{t'} d\tau H_I(q) \right) \exp \left(i \int_t^{t'} d\tau [p\dot{q} - H_0(p, q)] \right) \\
 &= \int \mathcal{D}q \mathcal{D}\frac{p}{2\pi} \exp \left(i \int_t^{t'} d\tau [p\dot{q} - H(p, q)] \right). \tag{9.62}
 \end{aligned}$$

The combination of exponentials in the last step is allowed because we are simply dealing with classical quantities (not operators!).

9.5 The generating functional for time ordered products

By introducing a source-term, $L(q, \dot{q}) \rightarrow L(q, \dot{q}) + J(t) \cdot q$, it is possible to switch on an interaction, physically pictured as, say, the creation of an electron (think of a radio-tube, making the electron) and the absorption of an electron (think of a detector). Before and after these processes there is only the vacuum or ground-state $|0\rangle$. Consider, furthermore, a set $|n\rangle$ of physical eigenstates. The Heisenberg state $|q, t\rangle$ is related to the Schrödinger state $|q\rangle$ by $|q, t\rangle = e^{iHt}|q\rangle$, i.e. for the physical states

$$\langle q, t | n \rangle = \langle q | n \rangle e^{-iE_n t} \tag{9.63}$$

with $\langle q | n \rangle$ the time-independent wave function, for example

$$\begin{aligned}
 \langle x | n \rangle &= \phi_n(x) = e^{i k_n x} \quad \text{for plane waves,} \\
 \langle x | n \rangle &= e^{-\omega^2 x^2 / 2} \quad \text{for g.s. harmonic oscillator.}
 \end{aligned}$$


Considering the source to be present between times t and t' , which in turn are embedded between an early time T and a future time T' , i.e. $T < t < t' < T'$, one has

$$\langle q', t' | q, t \rangle^J = \int \mathcal{D}q \exp \left(i \int_t^{t'} d\tau [L(q, \dot{q}) + J(\tau)q] \right), \tag{9.64}$$

$$\langle Q', T' | Q, T \rangle^J = \int dq' dq \langle Q', T' | q', t' \rangle \langle q', t' | q, t \rangle^J \langle q, t | Q, T \rangle, \tag{9.65}$$

with

$$\langle q, t | Q, T \rangle = \sum_n \langle q, t | n \rangle \langle n | Q, T \rangle = \sum_n \phi_n(q, t) \phi_n^*(Q) e^{+iE_n T}. \tag{9.66}$$


 Figure 9.2: Analytic continuation of boundaries T and T'

We can project out the groundstate by an analytic continuation in the time, $T \rightarrow i\infty$ and $T' \rightarrow -i\infty$, in which case $e^{iE_n T} \rightarrow e^{-E_n \cdot \infty}$ and the term $e^{iE_0 T}$ is dominant, thus

$$\lim_{T \rightarrow i\infty} \frac{\langle q, t | Q, T \rangle}{e^{iE_0 T} \phi_0^*(Q)} = \phi_0(q, t).$$

We define the generating functional $Z[J]$ as

$$Z[J] = \lim_{\substack{T' \rightarrow -i\infty \\ T \rightarrow i\infty}} \frac{\langle Q', T' | Q, T \rangle^J}{e^{-iE_0(T'-T)} \phi_0(Q') \phi_0^*(Q)} \quad (9.67)$$

$$= \int dq' dq \phi_0^*(q', t') \langle q', t' | q, t \rangle^J \phi_0(q, t) \quad (9.68)$$

$$= \langle 0_{out} | 0_{in} \rangle^J. \quad (9.69)$$

The factor that has been divided out in the first line of this equation is a numerical factor depending on the boundaries of the space-time volume (T and T'). The generating functional precisely represents the vacuum to vacuum amplitude from initial ('in') to final ('out') situation in the presence of a source. The importance of $Z[J]$ is that the time ordered product of operators can be obtained from it. Since we have (neglecting multiplicative factors),

$$Z[J] = \lim_{\substack{T' \rightarrow -i\infty \\ T \rightarrow i\infty}} \int \mathcal{D}q \exp \left(i \int_T^{T'} d\tau \left[L \left(q, \frac{dq}{d\tau} \right) + J(\tau)q(\tau) \right] \right), \quad (9.70)$$

one immediately finds

$$\frac{\delta^n Z[J]}{\delta J(t_1) \dots \delta J(t_n)} \Big|_{J=0} = (i)^n \langle 0 | \mathcal{T} Q(t_1) \dots Q(t_n) | 0 \rangle, \quad (9.71)$$

hence the name generating functional.

9.6 Euclidean formulation

The above expression (Eq. 9.70) for the generating functional is in fact ill-defined. Better is the use of imaginary time $t = -i\tilde{t}$, such that

$$\begin{aligned} t \rightarrow i\infty &\leftrightarrow \tilde{t} \rightarrow -\infty, \\ t' \rightarrow -i\infty &\leftrightarrow \tilde{t}' \rightarrow \infty. \end{aligned}$$

In terms of the imaginary time we can write the Euclidean generating functional,

$$Z_E[J] = \int \mathcal{D}q \exp \left(\int_{-\infty}^{\infty} d\tilde{\tau} \left[L \left(q, i \frac{dq}{d\tilde{\tau}} \right) + J(\tilde{\tau})q(\tilde{\tau}) \right] \right) \quad (9.72)$$

$$= \int \mathcal{D}q \exp \left(- \int_{-\infty}^{\infty} d\tilde{\tau} \left[L_E \left(q, \frac{dq}{d\tilde{\tau}} \right) - J(\tilde{\tau})q(\tilde{\tau}) \right] \right), \quad (9.73)$$

where

$$L_E \left(q, \frac{dq}{d\tilde{\tau}} \right) \equiv -L \left(q, i \frac{dq}{d\tilde{\tau}} \right), \quad (9.74)$$

e.g. when

$$L \left(q, \frac{dq}{dt} \right) = \frac{1}{2} \left(\frac{dq}{dt} \right)^2 - V(q) \quad (9.75)$$

$$= -\frac{1}{2} \left(\frac{dq}{d\tilde{t}} \right)^2 - V(q),$$

$$L_E \left(q, \frac{dq}{d\tilde{t}} \right) = \frac{1}{2} \left(\frac{dq}{d\tilde{t}} \right)^2 + V(q), \quad (9.76)$$

which is a positive definite quantity, ensuring convergence for the functional integral $Z_E[J]$. As discussed in the previous section the interesting quantities are obtained from functional differentiation with respect to the sources. The differentiations in $Z[J]$ and $Z_E[J]$ are related,

$$\frac{1}{Z[J]} \frac{\delta^n Z[J]}{\delta J(t_1) \dots \delta J(t_n)} \Big|_{J=0} = (i)^n \frac{1}{Z_E[J]} \frac{\delta^n Z_E[J]}{\delta J(\tilde{t}_1) \dots \delta J(\tilde{t}_n)} \Big|_{\substack{J=0 \\ \tilde{t} = it}}, \quad (9.77)$$

where the expressions have been divided by $Z[J]$ and $Z_E[J]$ in order to get rid of dependence on multiplicative factors.

Exercises

Exercise 9.1

Convince yourself by choosing an appropriate (orthonormal) set of functions that

$$\int \mathcal{D}\alpha \exp \left(-\frac{1}{2} \alpha K \alpha \right) = \frac{1}{\sqrt{\det K}},$$

for real-valued functions,

$$\int \mathcal{D}\alpha \mathcal{D}\alpha^* \exp(-\alpha^* K \alpha) = \frac{1}{\det K}.$$

for complex-valued functions and

$$\int \mathcal{D}\theta^* \mathcal{D}\theta \exp(-\theta^* K \theta) = \det K.$$

for Grassmann valued functions. In all of these cases the determinant of the (hermitean) operator K is defined as $\det K \equiv \prod_p K_p$, where K_p are the eigenvalues of the matrix K_{mn} in $K(x, y) = \sum_{m,n} f_m(x) K_{mn} f_n^*(y)$.

Exercise 9.2

Check the examples of functional derivation for real-, complex- and Grassmann-valued functions.

Exercise 9.3

Prove the following relations

$$\int \mathcal{D}\alpha \exp\left(-\frac{1}{2} \alpha K \alpha - \alpha \omega\right) = \frac{1}{\sqrt{\det K}} \exp\left(\frac{1}{2} \omega K^{-1} \omega\right), \quad (9.78)$$

$$\int \mathcal{D}\alpha^* \mathcal{D}\alpha \exp(-\alpha^* K \alpha - \alpha^* \omega - \omega^* \alpha) = \frac{1}{\det K} \exp(\omega^* K^{-1} \omega), \quad (9.79)$$

$$\int \mathcal{D}\theta^* \mathcal{D}\theta \exp(-\theta^* K \theta - \theta^* \eta - \eta^* \theta) = \det K \exp(\eta^* K^{-1} \eta), \quad (9.80)$$

where $\int dy K(x, y) K^{-1}(y, z) = \delta(x - z)$.

Chapter 10

Feynman diagrams for scattering amplitudes

10.1 Generating functionals for free scalar fields

The generating functional for quantum fields is a generalization of the results in the previous section to a system with more degrees of freedom, i.e. $\phi(\mathbf{x}, t)$ is considered as a set of quantum operators $\phi_{\mathbf{x}}(t)$ in the Heisenberg picture and

$$\begin{aligned} \langle \phi'; x', t' | \phi; x, t \rangle &= \int \mathcal{D}\phi \mathcal{D}\Pi \exp \left(i \int_t^{t'} d\tau d^3x \left[\Pi(x) \dot{\phi}(x) - \mathcal{H}(\phi, \Pi) \right] \right) \\ &= \int \mathcal{D}\phi \exp \left(i \int_{\sigma}^{\sigma'} d^4x \mathcal{L}(x) \right) \end{aligned} \quad (10.1)$$

and

$$Z[J] = \lim_{\substack{T \rightarrow i\infty \\ T' \rightarrow -i\infty}} \int \mathcal{D}\phi \exp \left(i \int_T^{T'} d^4x [\mathcal{L}(\phi, \partial_{\mu}\phi) + J\phi] \right) \quad (10.2)$$

$$= \int \mathcal{D}\phi \exp \left(- \int_{-\infty}^{\infty} d^4x_E [\mathcal{L}_E(\phi, \partial_{\mu}\phi) - J\phi] \right) \quad (10.3)$$

The Euclidean formulation is as before, implying at the level of four vectors in coordinate and momentum space (when $k \cdot x \equiv k_E \cdot x_E$) for instance

$$\begin{aligned} x^4 &= i x^0 = i t & k^4 &= -i k^0 \\ x \cdot x &= (x^0)^2 - \sum_{i=1}^3 (x^i)^2 = - \sum_{\mu=1}^4 (x^{\mu})^2 & k \cdot k &= (k^0)^2 - \sum_{i=1}^3 (k^i)^2 = - \sum_{\mu=1}^4 (k^{\mu})^2 \\ d^4x_E &= i d^4x & d^4k_E &= -i d^4k \\ \mathcal{L} &= \frac{1}{2} \partial_{\mu}\phi \partial^{\mu}\phi - \frac{1}{2} M^2 \phi^2 & \mathcal{L}_E &= \frac{1}{2} (\partial_E^{\mu}\phi)^2 + \frac{1}{2} M^2 \phi^2 \end{aligned}$$

Furthermore we find the n -point Green functions

$$\langle 0 | \mathcal{T} \phi(x_1) \dots \phi(x_n) | 0 \rangle = (-i)^n \frac{\delta^n Z[J]}{\delta J(x_1) \dots \delta J(x_n)} \Big|_{J=0} \equiv G^{(n)}(x_1, \dots, x_n), \quad (10.4)$$

for which we introduce the picture

$$G^{(n)}(x_1, \dots, x_n) = \begin{array}{c} x_1 \\ \times \\ \diagdown \\ \text{---} \circ \text{---} \\ \diagup \\ x_2 \\ \times \\ \text{---} \circ \text{---} \\ \diagup \\ x \\ \times \\ \text{---} \circ \text{---} \\ \vdots \\ \times \\ \text{---} \circ \text{---} \\ \vdots \\ x_n \\ \times \\ \text{---} \circ \text{---} \end{array} .$$

These Green functions appear in the expansion

$$Z[J] = \sum_n \frac{i^n}{n!} \int d^4x_1 \dots d^4x_n G^{(n)}(x_1, \dots, x_n) J(x_1) \dots J(x_n). \quad (10.5)$$

$Z_0[J]$ for the free scalar field

For the (free) scalar field lagrangian

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} M^2 \phi^2, \quad (10.6)$$

the generating functional is (using $\overleftarrow{\partial}_\mu = -\overrightarrow{\partial}_\mu$) given by

$$Z_0[J] = \int \mathcal{D}\phi \exp \left(-i \int d^4x \left[\frac{1}{2} \phi \underbrace{(\partial_\mu \partial^\mu + M^2)}_K \phi - J\phi \right] \right) \quad (10.7)$$

where $K_{xy} = (\partial_\mu \partial^\mu + M^2) \delta^4(x - y)$. As discussed in the previous section (exercise 9.3) one finds

$$Z_0[J] = \exp \left(-\frac{1}{2} i \int d^4x d^4y J(x) \Delta_F(x - y) J(y) \right) \equiv \exp(i W_0[J]), \quad (10.8)$$

where we made the choice $Z_0[0] = 1$ and

$$(\partial_\mu \partial^\mu + M^2) \Delta_F(x - y) = -\delta^4(x - y), \quad (10.9)$$

i.e. Δ_F (the so-called Feynman propagator) is the Green's function of the Klein-Gordon equation. Furthermore, with $Z_0[0] = 1$, one immediately sees that

$$i\Delta_F(x_1 - x_2) = \frac{(-i)^2}{Z_0[0]} \left. \frac{\delta^2 Z_0[J]}{\delta J(x_1) \delta J(x_2)} \right|_{J=0} = G_0^{(2)}(x_1 - x_2) \equiv x_1 \times \longrightarrow \times x_2 \quad (10.10)$$

In order to determine Δ_F , consider the general solution of

$$(\partial_\mu \partial^\mu + M^2) \Delta(x) = -\delta^4(x),$$

which can be written as

$$\Delta(x) = \int \frac{d^4k}{(2\pi)^4} e^{-ik \cdot x} \tilde{\Delta}(k) \quad (10.11)$$

with $(k^2 - M^2) \tilde{\Delta}(k) = 1$ or

$$\tilde{\Delta}(k) = \frac{1}{k^2 - M^2} = \frac{1}{(k^0)^2 - \mathbf{k}^2 - M^2} = \frac{1}{(k^0)^2 - E^2}. \quad (10.12)$$

Depending on the path chosen in the complex k^0 plane (see exercise 7.4) one distinguishes the *retarded* Green's function

$$\Delta_R(x) = \int \frac{d^4k}{(2\pi)^4} \frac{e^{-ik \cdot x}}{(k^0 + i\epsilon)^2 - E^2}, \quad (10.13)$$

satisfying $\Delta_R(\mathbf{x}, t) = 0$ for $t < 0$,
the *advanced* Green's function

$$\Delta_A(x) = \int \frac{d^4 k}{(2\pi)^4} \frac{e^{-i k \cdot x}}{(k^0 - i\epsilon)^2 - E^2}, \quad (10.14)$$

satisfying $\Delta_A(\mathbf{x}, t) = 0$ for $t > 0$,
the *causal* Green's function

$$\Delta_C(x) = \int \frac{d^4 k}{(2\pi)^4} \frac{e^{-i k \cdot x}}{(k^0)^2 - E^2 + i\epsilon} = \int \frac{d^4 k}{(2\pi)^4} \frac{e^{-i k \cdot x}}{k^2 - M^2 + i\epsilon}, \quad (10.15)$$

and finally the Green's function

$$\bar{\Delta}(x) = \text{PV} \int \frac{d^4 k}{(2\pi)^4} \frac{e^{-i k \cdot x}}{k^2 - M^2}. \quad (10.16)$$

They are solutions of the inhomogeneous equation. The solution of the homogeneous equation $(\partial_\mu \partial^\mu + M^2) \Delta(x) = 0$ can also be written as an integral in k -space,

$$\Delta(x) = - \int_C \frac{d^4 k}{(2\pi)^4} \frac{e^{-i k \cdot x}}{k^2 - M^2}, \quad (10.17)$$

where C is a closed contour in the complex k^0 -plane. The contours for

$$\begin{aligned} i \Delta(x) &= [\phi(x), \phi(0)] = i \Delta_+(x) + i \Delta_-(x) \\ i \Delta_+(x) &= [\phi_+(x), \phi_-(0)] = \int \frac{d^3 k}{(2\pi)^3 2E} e^{-i k \cdot x} \\ i \Delta_-(x) &= [\phi_-(x), \phi_+(0)] = \int \frac{d^3 k}{(2\pi)^3 2E} e^{-i k \cdot x} \end{aligned}$$

are also shown in exercise (7.4). It is straightforward (closing contours in the appropriate half of the complex plane) to prove for instance that

$$i \Delta_C(x) = \theta(x^0) i \Delta_+(x) - \theta(-x^0) i \Delta_-(x). \quad (10.18)$$

Thus we see various solutions. In order to see which is the appropriate Green's function to be used in the generating functional $Z_0[J]$ we will take two routes. The first possibility is to consider the well-defined Euclidean formulation, the second is to explicitly consider $\langle 0 | \mathcal{T} \phi(x_1) \phi(x_2) | 0 \rangle$.

- Firstly, in the (well-defined) Euclidean formulation starting with $\mathcal{L}_E = \frac{1}{2} (\partial_E^\mu \phi)^2 + \frac{1}{2} M^2 \phi^2 - J\phi$ with the equation of motion $((\partial_E^\mu)^2 - M^2) \phi(x) = -J(x)$, the generating functional can be written

$$Z_0[J] = \int \mathcal{D}\phi \exp \left(\int d^4 x_E \left[\frac{1}{2} \phi (\partial_E^\mu \partial_E^\mu - M^2) \phi + J\phi \right] \right) \quad (10.19)$$

$$= \exp \left(-\frac{1}{2} \int d^4 x_E d^4 y_E J(x) (-i \Delta_F(x-y)) J(y) \right), \quad (10.20)$$

where


$$((\partial_E^\mu)^2 - M^2) (i \Delta_F(x-y)) = -\delta^4(x-y) \quad (10.21)$$

or (see fig. 10.1 for contours)

$$\Delta_F(x) = -i \int_{k_E^0 = -\infty}^{k_E^0 = \infty} \frac{d^4 k_E}{(2\pi)^4} \frac{e^{-i k_E \cdot x}}{k_E^2 + M^2} \quad (10.22)$$

$$= \int_{k^0 = -i\infty}^{k^0 = i\infty} \frac{d^4 k}{(2\pi)^4} \frac{e^{-i k \cdot x}}{k^2 - M^2}, \quad (10.23)$$

where the latter contour can be deformed to the contour for Δ_C , i.e. $\Delta_F = \Delta_C$.


 Figure 10.1: Contour in the k^0 plane for the Feynman propagator in Euclidean and Minkowski space

- As the second possibility, we use the fact that $i\Delta_F(x-y) = G^{(2)}(x-y) = \langle 0|\mathcal{T}\phi(x)\phi(y)|0\rangle$ to calculate Δ_F via the time-ordered product. We find (see also section 7.3)


$$\begin{aligned}
 i\Delta_F(x-y) &= \langle 0|\mathcal{T}\phi(x)\phi(y)|0\rangle \\
 &= \theta(x^0 - y^0)\langle 0|\phi(x)\phi(y)|0\rangle + \theta(y^0 - x^0)\langle 0|\phi(y)\phi(x)|0\rangle \\
 &= \theta(x^0 - y^0)\langle 0|\phi_+(x)\phi_-(y)|0\rangle + \theta(y^0 - x^0)\langle 0|\phi_+(y)\phi_-(x)|0\rangle \\
 &= \theta(x^0 - y^0)\langle 0|[\phi_+(x), \phi_-(y)]|0\rangle - \theta(y^0 - x^0)\langle 0|[\phi_-(x), \phi_+(y)]|0\rangle \\
 &= \theta(x^0 - y^0)i\Delta_+(x-y) - \theta(y^0 - x^0)i\Delta_-(x-y) \\
 &= i\Delta_C(x-y),
 \end{aligned} \tag{10.24}$$

the same result as above.

Knowing the explicit form of $Z_0[J]$ it is straightforward to calculate the 4-points Green's function. It should be clear that $G_0^{(4)}$ can be expressed in terms of $G_0^{(2)} = i\Delta_F$, because this is the only quantity entering $Z_0[J]$. Neglecting multiplicative factors or equivalently assuming that $Z_0[0] = 1$, we have

$$\begin{aligned}
 G_0^{(4)}(x_1, x_2, x_3, x_4) &= \langle 0|\mathcal{T}\phi(x_1)\phi(x_2)\phi(x_3)\phi(x_4)|0\rangle \\
 &= (-i)^4 \frac{\delta^4}{\delta J(x_1)\dots\delta J(x_4)} \exp\left(-\frac{1}{2}i \int J \Delta_F J\right) \Big|_{J=0} \\
 &= -[\Delta_F(x_1 - x_2)\Delta_F(x_3 - x_4) + \Delta_F(x_1 - x_3)\Delta_F(x_2 - x_4) + \Delta_F(x_1 - x_4)\Delta_F(x_2 - x_3)],
 \end{aligned} \tag{10.25}$$

or diagrammatically



10.2 Generating functionals for interacting scalar fields

When interactions are present, i.e. $\mathcal{L}(\phi) = \mathcal{L}_0(\phi) + \mathcal{L}_I(\phi)$, the generating functional can be written,

$$Z[J] = \int \mathcal{D}\phi \exp\left(i \int d^4x [\mathcal{L}_0(\phi) + \mathcal{L}_I(\phi) + J\phi]\right) \quad (10.26)$$

$$= \exp\left(i \int d^4z \mathcal{L}_I\left(\frac{1}{i} \frac{\delta}{\delta J(z)}\right)\right) \int \mathcal{D}\phi \exp\left(i \int d^4x [\mathcal{L}_0(\phi) + J\phi]\right) \quad (10.27)$$

$$= \exp\left(i \int d^4z \mathcal{L}_I\left(\frac{1}{i} \frac{\delta}{\delta J(z)}\right)\right) \exp\left(-\frac{1}{2} i \int d^4x d^4y J(x) \Delta_F(x-y) J(y)\right) \quad (10.28)$$

$$= \exp\left(\frac{1}{2} \int d^4x d^4y \frac{\delta}{\delta\phi(x)} i \Delta_F(x-y) \frac{\delta}{\delta\phi(y)}\right) \exp\left(i \int d^4z [\mathcal{L}_I(\phi) + J(z)\phi(z)]\right) \Big|_{\phi=0}. \quad (10.29)$$

This expression will be the one from which Feynman rules will be derived, with propagators ($i \Delta_F$) being connected to vertices ($i \mathcal{L}_I$) according to the above expression for the functional $Z[J]$.

Consider as an example the interaction

$$\mathcal{L}_I(\phi) = -\frac{g}{4!} \phi^4$$

in the scalar field theory discussed sofar. To zeroth order in the coupling one has

$$Z^{(g^0)}[J] = Z_0[J] = \exp\left(-\frac{1}{2} i \int J \Delta_F J\right), \quad (10.30)$$

and in zeroth order

$$G^{(0,g^0)} = G_0^{(0)} = 1, \quad (10.31)$$

$$G^{(2,g^0)}(x_1, x_2) = G_0^{(2)}(x_1, x_2) = i \Delta_F(x_1 - x_2) = \text{---} \times \text{---} \quad (10.32)$$

$$G^{(4,g^0)}(x_1, x_2, x_3, x_4) = G_0^{(4)}(x_1, x_2, x_3, x_4) = \text{---} \times \text{---} + \text{---} \text{---} + \text{---} \text{---} + \text{---} \times \text{---} \quad (10.33)$$

To first order in g one has

$$\begin{aligned} Z^{(g^1)}[J] &= i \int d^4z \mathcal{L}_I\left(\frac{1}{i} \frac{\delta}{\delta J(z)}\right) \exp\left(-\frac{1}{2} i \int J \Delta_F J\right) \\ &= -i \frac{g}{4!} \int d^4z \left\{ -3 \Delta_F^2(0) + 6i \Delta_F(0) \left[\int d^4x \Delta_F(z-x) J(x) \right]^2 \right. \\ &\quad \left. + \left[\int d^4x \Delta_F(z-x) J(x) \right]^4 \right\} \exp\left(-\frac{1}{2} i \int J \Delta_F J\right). \end{aligned} \quad (10.34)$$

Introducing a vertex point to which four propagators are connected.

$$\text{---} \times \text{---} \equiv -i g \int d^4z \quad (10.35)$$

one has

$$Z^{(g^1)}[J] = \left(\frac{1}{8} \text{diagram} - \frac{1}{4} \text{diagram} + \frac{1}{24} \text{diagram} \right) Z_0[J], \quad (10.36)$$

from which one obtains

$$G^{(0,g^1)} = \frac{1}{8} \text{diagram}, \quad (10.37)$$

$$G^{(2,g^1)}(x_1, x_2) = \frac{1}{8} \text{diagram} + \frac{1}{2} \text{diagram}, \quad (10.38)$$

The result for the 4-points Green's function is left as an exercise.

Connected Green's functions

We have now seen the Green's functions as the quantities appearing in the expansion of the generating functional $Z[J]$,

$$Z[J] = \sum_n \frac{i^n}{n!} \int d^4x_1 \dots d^4x_n G^{(n)}(x_1, \dots, x_n) J(x_1) \dots J(x_n). \quad (10.39)$$

We have calculated the function up to first order in the coupling constant for ϕ^4 -theory. In the free case we have seen that the exponent of $Z[J]$ contained all essential information. Diagrammatically this exponent only contained the two-point function, which also was the only Green's function for which the diagram was connected, i.e. did not contain parts that could be written as products of simpler Green's functions. This remains also true for the interacting case. To see this write

$$Z[J] = \exp(iW[J]) \quad \text{or} \quad iW[J] = \ln Z[J], \quad (10.40)$$

with (by definition) the expansion

$$W[J] = \sum_n \frac{i^{n-1}}{n!} \int d^4x_1 \dots d^4x_n G_c^{(n)}(x_1, \dots, x_n) J(x_1) \dots J(x_n) \quad (10.41)$$

in terms of so-called connected Green's functions,

$$G_c^{(n)}(x_1, \dots, x_n) = (-i)^{n-1} \frac{\delta^n W[J]}{\delta J(x_1) \dots \delta J(x_n)} \Big|_{J=0}, \quad (10.42)$$

which are denoted

$$G_c^{(n)}(x_1, \dots, x_n) = \text{diagram}$$

Specific examples are

$$G_c^{(0)} = \text{diagram} \quad (\text{vacuum bubble})$$

$$G_c^{(1)}(x_1) = \text{diagram} \quad (\text{tadpole})$$

$$G_c^{(2)}(x_1, x_2) = \text{diagram} \quad (\text{connected propagator})$$

To see that $Z[J] = \exp(iW[J])$ contains all non-connected diagrams can be performed inductively, but is easily illustrated by writing down the first few terms. If

$$iW[J] = \text{blob} + i \int d^4x_1 \{ \text{cross} \text{---} \text{blob} \} + \frac{i^2}{2!} \int d^4x_1 d^4x_2 \{ \text{cross} \text{---} \text{blob} \text{---} \text{cross} \},$$

then

$$\begin{aligned} \exp(iW[J]) &= 1 + iW[J] + \frac{(iW[J])^2}{2!} + \frac{(iW[J])^3}{3!} + \dots \\ &= 1 \\ &\quad + \text{blob} + i \int d^4x_1 \{ \text{cross} \text{---} \text{blob} \} + \frac{i^2}{2!} \int d^4x_1 d^4x_2 \{ \text{cross} \text{---} \text{blob} \text{---} \text{cross} \} \\ &\quad + \frac{1}{2!} \left\{ \begin{array}{l} \text{blob} \\ \text{blob} \end{array} \right\} + i \int d^4x_1 \left\{ \begin{array}{l} \text{cross} \text{---} \text{blob} \\ \text{blob} \end{array} \right\} + \frac{i^2}{2!} \int d^4x_1 d^4x_2 \left\{ \begin{array}{l} \text{cross} \text{---} \text{blob} \text{---} \text{cross} \\ \text{blob} \\ \text{cross} \text{---} \text{blob} \\ \text{blob} \text{---} \text{cross} \end{array} \right\} \\ &\quad + \frac{1}{3!} \left\{ \begin{array}{l} \text{blob} \\ \text{blob} \\ \text{blob} \end{array} \right\} + i \int d^4x_1 \left\{ \begin{array}{l} \text{cross} \text{---} \text{blob} \\ \text{blob} \\ \frac{1}{2!} \left\{ \begin{array}{l} \text{blob} \\ \text{blob} \end{array} \right\} \end{array} \right\} + \frac{i^2}{2!} \int d^4x_1 d^4x_2 \left\{ \begin{array}{l} \text{cross} \text{---} \text{blob} \text{---} \text{cross} \\ \text{blob} \\ \frac{1}{2!} \left\{ \begin{array}{l} \text{blob} \\ \text{blob} \end{array} \right\} \\ \text{cross} \text{---} \text{blob} \\ \text{blob} \text{---} \text{cross} \\ \text{blob} \end{array} \right\} \\ &\quad + \dots \end{aligned}$$

Notes:

- (i) In $G^{(n)}$ the connected Green functions $G_c^{(n)}$ appear with particular combinatorial factors (accounted for in the definition of Feynman rules to be discussed later).
- (ii) Note that $Z[0] = \exp(iW[0])$ appears as a multiplicative factor, in the expansion of which all vacuum blobs are contained,

$$Z[0] = 1 + \text{blob} + \frac{1}{2!} \left\{ \begin{array}{l} \text{blob} \\ \text{blob} \end{array} \right\} + \dots$$

These can be divided out. In the expansion of $Z[J]/Z[0]$ one (by definition) has the so-called *source connected* Green's functions $G_{sc}^{(n)}$.

(iii) $G_{sc}^{(1)} = G_c^{(1)}$.

(iv) If

$$\left. \frac{\delta Z[J]}{\delta J(x)} \right|_{J=0} = i \langle 0 | \phi(x) | 0 \rangle = 0,$$

i.e. the vacuum expectation value of the field $\phi(x)$ is zero, implying the absence of tadpoles, then $G_{sc}^{(n)} = G_c^{(n)}$ for $n \leq 3$.

For the free scalar theory, considered explicitly, we have

$$W_0[J] = -\frac{1}{2} \int d^4x d^4y J(x) \Delta_F(x-y) J(y), \quad (10.43)$$

implying as expected as the only nonzero connected Green's function

$$G_c^{(2)}(x_1 - x_2) = (-i) \frac{\delta^2 W[J]}{\delta J(x_1) \delta J(x_2)} \Big|_{J=0} = G_0^{(2)}(x_1 - x_2) = i \Delta_F(x_1 - x_2). \quad (10.44)$$

For the interacting theory, one finds for the connected 4-point Green's function at order g ,

$$G_c^{(4,g^1)}(x_1, x_2, x_3, x_4) = \begin{array}{c} x_1 \quad \quad x_3 \\ \quad \diagdown \quad \diagup \\ \quad \quad z \\ \quad \diagup \quad \diagdown \\ x_2 \quad \quad x_4 \end{array} \quad (10.45)$$

as the only surviving diagram (see exercises).

10.3 Interactions and the S-matrix

The S-matrix

The S-matrix transforms initial state free particle states (*in-states*) $|\alpha; in\rangle = |p_1, \dots, p_n; in\rangle$ into final state particle states (*out-states*) $|\beta; out\rangle = |p'_1, \dots, p'_m; out\rangle$ (suppressing except momenta all other quantum numbers),

$$\begin{aligned} S_{\beta\alpha} &\equiv \langle \beta; out | \alpha; in \rangle \Leftrightarrow \langle \beta; in | S = \langle \beta; out | \\ &\Leftrightarrow |\alpha; in\rangle = S |\alpha; out\rangle. \end{aligned} \quad (10.46)$$

The properties of the S-matrix are

(1) The vacuum is invariant or $|S_{00}| = 1$.

Proof: $\langle 0; in | S = \langle 0; out | = e^{i\varphi_0} \langle 0; in |$ (choose $\varphi_0 = 0$).

(2) The one-particle state is invariant (conservation of energy and momentum; translation invariance), $\langle p; in | S | p'; in \rangle = \langle p; out | p'; in \rangle = \langle p; in | p'; in \rangle = \langle p; out | p'; out \rangle = \langle p | p' \rangle$.

(3) S is unitary (it conserves the scalar product from initial to final state).

Proof: $\langle \alpha; in | S = \langle \alpha; out |$ and $S^\dagger |\alpha; in\rangle = |\alpha; out\rangle$,

thus $\langle \beta; in | S S^\dagger |\alpha; in\rangle = \langle \beta; out | \alpha; out \rangle = \delta_{\alpha\beta} \leftrightarrow S S^\dagger = 1$.

Next, this will be translated to the action on fields. Also for free field (Heisenberg) operators a distinction is made between ϕ_{in} and ϕ_{out} . In line with the consideration of the generating functional representing the vacuum to vacuum amplitude we consider fields ϕ_{in} , ϕ_{out} and the interpolating field $\phi(x)$,

$$\begin{array}{ccc} t = -\infty & & t = +\infty \\ \phi_{in}(x) & \phi(x) & \phi_{out}(x) \end{array}$$

where ϕ_{in} and ϕ_{out} transform under the Poincaré group as scalar fields and satisfy the homogeneous Klein-Gordon equation with the physical mass M , e.g.

$$(\partial_\mu \partial^\mu + M^2) \phi_{in}(x) = 0, \quad (10.47)$$

while ϕ satisfies the inhomogeneous Klein-Gordon equation with the bare mass M_0 (this is the mass appearing in the lagrangian \mathcal{L}_0),

$$(\partial_\mu \partial^\mu + M_0^2) \phi(x) = J(x). \quad (10.48)$$

The fact that ϕ_{in} and ϕ_{out} satisfy the homogeneous Klein-Gordon equation with the physical mass M implies that they create particles and antiparticles as discussed, e.g.

$$\phi_{in}(x) = \int \frac{d^3k}{(2\pi)^3 2E} [a(k) f_k(x) + a^\dagger(k) f_k^*(x)]. \quad (10.49)$$

The field ϕ can be expressed in $\phi_{in/out}$ using retarded or advanced Green's functions,

$$(\partial_\mu \partial^\mu + M^2) \phi(x) = J(x) + (M^2 - M_0^2) \phi(x) = \tilde{J}(x) \quad (10.50)$$

$$\begin{aligned} \phi(x) &= \sqrt{Z} \phi_{in}(x) - \int d^4y \Delta_R(x-y) \tilde{J}(y) \\ &= \sqrt{Z} \phi_{in}(x) - \int d^4y d^4z \Delta_R(x-y) K(y,z) \phi(z) \end{aligned} \quad (10.51)$$

$$\phi(x) = \sqrt{Z} \phi_{out}(x) - \int d^4y d^4z \Delta_A(x-y) K(y,z) \phi(z), \quad (10.52)$$

where Z is a constant, which in a later stage (renormalization) will become more important.

Although the above, as it stands, implies the strong (operator) convergence $\phi(x) \rightarrow \sqrt{Z} \phi_{in}(x)$, this can actually not be used as it would imply $[\phi(x), \phi(y)] = Z [\phi_{in}(x), \phi_{in}(y)] = iZ \Delta(x-y)$, a causality condition that can be proven to imply the absence of interactions. The convergence therefore must be weakened to

$$\langle \alpha | \phi^f(t) \beta \rangle \xrightarrow{t \rightarrow -\infty} \sqrt{Z} \langle \alpha | \phi_{in}^f(t) \beta \rangle \quad (10.53)$$

for normalizable states $|\alpha\rangle$ and $|\beta\rangle$ and $\phi^f(t) \equiv \int d^3(x) f^*(x) i \overleftrightarrow{\partial}_0 \phi(x)$ with f a normalizable solution of the Klein-Gordon equation (wave packet). Considering plane waves one sees from

$$\begin{aligned} \langle 0 | \phi(x) | p \rangle &= \lim_{t \rightarrow \infty} \sqrt{Z} \langle 0 | \phi_{out}(x) | p \rangle = \sqrt{Z} e^{-i p \cdot x} \\ &= \lim_{t \rightarrow -\infty} \sqrt{Z} \langle 0 | \phi_{in}(x) | p \rangle = \sqrt{Z} e^{-i p \cdot x} \end{aligned}$$

that identical normalization of (single-particle) plane waves in initial and final state implies the same wave function normalization Z for in and out fields.

The relation between S-matrix and in- and out-fields is: $\phi_{in}(x) = S \phi_{out}(x) S^{-1}$.

Proof:

$$\langle \beta; out | \phi_{out} = \left\{ \begin{array}{l} \langle \beta; in | \phi_{in} S \\ \langle \beta; in | S \phi_{out} \end{array} \right\} \rightarrow \phi_{in} S = S \phi_{out}(x).$$

Finally we check that as expected S does not spoil Poincaré invariance, i.e. S is invariant under Poincaré transformations: $U(\Lambda, a) S U^{-1}(\Lambda, a) = S$.

Proof: $\phi_{in}(\Lambda x + a) = U \phi_{in}(x) U^{-1} = U S \phi_{out}(x) S^{-1} U^{-1} = U S U^{-1} U \phi_{out}(x) U^{-1} U S^{-1} U^{-1} = U S U^{-1} \phi_{out}(\Lambda x + a) U S^{-1} U^{-1} \Leftrightarrow U S U^{-1} = S$.

The relation between S and Z[J]

To establish this relation, the source term $J\phi$ in the lagrangian density is treated in the interaction picture. The time evolution operator (see Eq. 9.62) is then

$$U_T^{T'}[J] = \mathcal{T} \exp \left(i \int_T^{T'} d^4x J(x) \phi(x) \right), \quad (10.54)$$

and satisfies the property

$$\frac{\delta U_T^{T'}[J]}{\delta J(x)} = i U_t^{T'}[J] \phi(x) U_T^t[J], \quad (10.55)$$

or for $U[J] \equiv U_{-\infty}^{\infty}[J]$

$$\begin{aligned} \frac{\delta U[J]}{\delta J(x)} &= i U_t^{\infty}[J] \phi(x) U_{-\infty}^t[J] \\ &\xrightarrow{t \rightarrow \infty} i \phi(x) U[J] \rightarrow i \sqrt{Z} \phi_{out}(x) U[J] \\ &\xrightarrow{t \rightarrow -\infty} i U[J] \phi(x) \rightarrow i \sqrt{Z} U[J] \phi_{in}(x). \end{aligned} \quad (10.56)$$

Since $\delta U[J]/\delta J(x)$ satisfies the same equations as $\phi(x)$ and we know the limits, we can, just as we did for $\phi(x)$, express it in terms of advanced and retarded Green's functions,

$$\begin{aligned} \frac{\delta U[J]}{\delta J(x)} &= i \sqrt{Z} U[J] \phi_{in}(x) - \int d^4 y d^4 z \Delta_R(x-y) K(y,z) \frac{\delta U[J]}{\delta J(z)} \\ &= i \sqrt{Z} \phi_{out}(x) U[J] - \int d^4 y d^4 z \Delta_A(x-y) K(y,z) \frac{\delta U[J]}{\delta J(z)}. \end{aligned}$$

Taking the difference between the two expressions

$$\begin{aligned} &\sqrt{Z} (\phi_{out}(x) U[J] - U[J] \phi_{in}(x)) \\ &= i \int d^4 y d^4 z (\Delta_R(x-y) - \Delta_A(x-y)) K(y,z) \frac{\delta U[J]}{\delta J(z)}, \end{aligned} \quad (10.57)$$

or

$$[\phi_{in}(x), S U[J]] = \frac{i}{\sqrt{Z}} \int d^4 y d^4 z \Delta(x-y) K(y,z) \frac{\delta}{\delta J(z)} S U[J]. \quad (10.58)$$

In order to find a solution to this equation note that, with the use of the Baker-Campbell-Hausdorff formula $e^{-B} A e^{+B} = A + [A, B]$ (for the case that $[A, B]$ is a c-number), one has

$$[A, e^B] = [A, B] e^B, \quad (10.59)$$

$$[A, e^B e^C] = [A, B + C] e^B e^C, \quad (10.60)$$

provided $[A, B]$ and $[A, C]$ are c-numbers. Thus applied to the field $\phi(x) = \phi_+(x) + \phi_-(x)$,

$$\begin{aligned} &[\phi(x), e^{\int d^4 x \phi_-(x) f(x)} e^{\int d^4 y \phi_+(y) f(y)}] \\ &= \int d^4 y [\phi(x), \phi(y)] f(y) e^{\int d^4 z \phi_-(z) f(z)} e^{\int d^4 z \phi_+(z) f(z)}, \end{aligned}$$

i.e.

$$[\phi(x), : e^{\int d^4 z \phi(z) f(z)} :] = i \int d^4 y \Delta(x-y) f(y) : e^{\int d^4 z \phi(z) f(z)} :, \quad (10.61)$$

where the normal ordered expression $: e^{\int \phi f} :$ is used, which is equal to the expression $e^{\int \phi_- f} e^{\int \phi_+ f}$ in which the creation operators are placed left of annihilation operators. Thus

$$S U[J] =: \exp \left(\frac{1}{\sqrt{Z}} \int d^4 x d^4 y \phi_{in}(x) K(x,y) \frac{\delta}{\delta J(y)} \right) : F[J], \quad (10.62)$$

where $F[J]$ is some (arbitrary) functional. Noting that $\langle 0 | e^A | 0 \rangle = 1$ it follows that

$$F[J] = \langle 0 | S U[J] | 0 \rangle = \langle 0 | U[J] | 0 \rangle = \langle 0_{out} | 0_{in} \rangle \propto Z[J], \quad (10.63)$$

while for $J = 0$ one has $U[0] = 1$, i.e.

$$S =: \exp\left(\frac{1}{\sqrt{Z}} \int d^4x d^4y \phi_{in}(x) K(x, y) \frac{\delta}{\delta J(y)}\right) : \frac{Z[J]}{Z[0]} \Big|_{J=0}. \quad (10.64)$$

Therefore, an S-matrix element between momentum eigenstates in initial and final states is found by considering those source-connected Green's functions (action of $\delta/\delta J$ on $Z[J]/Z[0]$) where the external sources $J(x_i)$ are replaced by the particle wave functions (which are the result of acting with $\phi_{in}(x_i)$ on momentum eigenstates). Note that the Green's function connecting the external point x_i with the bubble is annihilated by $K(x, y)$.

Usually we are interested in the part of the S-matrix describing the scattering,

$$S_{fi} = \delta_{fi} - i(2\pi)^4 \delta^4(P_i - P_f) \mathcal{M}_{fi}, \quad (10.65)$$

which is obtained considering only connected diagrams.

Explicitly, using that

$$\phi_{in}(x) = \int \frac{d^3k}{(2\pi)^3 2E} [a(k) f_k(x) + a^\dagger(k) f_k^*(x)].$$

we get the Lehman-Symanzik-Zimmerman (LSZ) reduction formula,

$$\begin{aligned} \langle p', \dots | S | p, \dots \rangle &= \dots \delta \dots + \left(\frac{1}{\sqrt{Z}}\right)^n \int d^4x' \dots d^4x \dots f_{p'}^*(x') \dots i\overrightarrow{K}_{x'} \dots \\ &\quad \times G_c^{(n)}(x', \dots, x, \dots) i\overleftarrow{K}_x \dots f_p(x) \dots, \end{aligned} \quad (10.66)$$

where $G_c^{(n)}$ is the connected Green's function, $i\overrightarrow{K}_{x'} = (\square_{x'} + M^2)$ precisely annihilating an external propagator $i\Delta_F$ in the Green's function.

Next we introduce the Fourier transform (after extracting a momentum conserving delta function coming from translation invariance, see exercise 10.2),

$$(2\pi)^4 \delta^4(p_1 + \dots + p_n) G^{(n)}(p_1, \dots, p_n) = \int \prod_{i=1}^n d^4x_i e^{ip_i \cdot x_i} G^{(n)}(x_1, \dots, x_n), \quad (10.67)$$

and the (amputated) Green's functions

$$\Gamma^{(n)}(p_1, \dots, p_n) = \left(\prod_{j=1}^n \frac{-i}{\Delta(p_j)}\right) G^{(n)}(p_1, \dots, p_n), \quad (10.68)$$

where $\Delta(p)$ is the Fourier transform of the (full) propagator $\Delta_F(x)$. It is straightforward to check that the S-matrix element now precisely is given by the amputated Green's function multiplied with the momentum space wave functions of the particles in initial and final state (by which we refer to the quantities multiplying the plane wave $e^{\pm ip \cdot x}$ in the field expansion, i.e. 1 for scalar case, $u(p)$, $v(p)$, $\bar{u}(p)$ and $\bar{v}(p)$ for fermions and $\epsilon_\mu^{(\lambda)}(p)$ for vector fields).

10.4 Feynman rules

The real scalar field

The procedure to obtain the matrix element is commonly summarized by a set of rules known as *Feynman rules*. They start with the *propagator* ($i\Delta_F(k)$) in momentum space, which is determined by the inverse of the operator found in the quadratic term in the lagrangian i.e.

$$\overrightarrow{\mathbf{k}} = \frac{i}{k^2 - M^2 + i\epsilon}$$

(in fact the inverse of the operator found in the quadratic term is for real scalar fields also multiplied by a factor 2, which cancels the factor 1/2 in the quadratic piece; the factor 2 corresponds to the two-points Green's function having two identical ends).

For the interaction terms in the lagrangian, to be precise $i\mathcal{L}_I$ vertices in momentum space are introduced,

$$\begin{array}{c} \diagup \\ \diagdown \end{array} = -i g$$


(multiplied with 4! corresponding to the allowed number of permutations of identical particles). At these vertices each line can be assigned a momentum, but overall momentum conservation at a vertex is understood.

Corresponding to external particles wave functions are introduced

$$\begin{array}{cc} \longrightarrow \mathbf{k} \text{ (blue semi-circle)} & 1 \\ \text{(blue semi-circle)} \longrightarrow \mathbf{k} & 1 \end{array}$$

In order to calculate the connected amplitude $-i\mathcal{M}_{fi}$ appearing in the S-matrix element these ingredients are combined using Eqs 10.66 and 10.27, which is summarized in the following rules:

(Rule 1) Start with external legs (incoming particles/outgoing particles) and draw all possible topologically different connected diagrams, for example up to order g^2 the scattering of two neutral spin 0 particles (real scalar field) is described by



(Rule 2) The contribution of each diagram is obtained by multiplying the contributions from propagators, vertices and external particle wave functions in that diagram. Note that in calculating amputated Green's functions external lines are neglected, or calculating full Green's functions external lines are treated as propagators.

(Rule 3) Carry out the integration over all internal momenta (keeping track of momentum conservation at all vertices!)

(Rule 4) Add a symmetry factor 1/S corresponding to permutation of internal lines and vertices (keeping external lines fixed). If problems arise go back to the defining expression for the generating function in 10.27.


For the symmetry factor consider the examples (given in G. 't Hooft and M. Veltman, *Diagrammar*) in the case of the interaction terms

$$\mathcal{L}_I(\phi) = -\frac{f}{3!} \phi^3 - \frac{g}{4!} \phi^4. \tag{10.69}$$

The vertices are:

$$\begin{array}{cc} \begin{array}{c} | \\ \diagdown \\ \diagup \end{array} = -i f & \begin{array}{c} \diagup \\ \diagdown \end{array} = -i g \end{array}$$

Consider first the lowest order self-energy diagram,




Draw two points corresponding to the two vertices and draw in each of these points the lines coming out of the vertices:



Now count in how many ways the lines can be connected with the same topological result. External line 1 can be attached in six, after that line 2 in three ways. Then there are two ways to connect the remaining lines such that the desired diagram results. Divide by the permutational factors of the vertices, which have been included in the definition of vertices (here 3! for each vertex). Finally divide by the number of permutation of the points that have identical vertices (here 2!). The total result is

$$\frac{1}{S} = \frac{6 \times 3 \times 2}{3! 3! 2!} = \frac{1}{2}.$$


As a second example consider the diagram



There are three vertices,



After connecting line 1 (6 ways) and line 2 (4 ways) we have



leaving $6 \times 3 \times 2$ ways to connect the rest as to get the desired topology. Dividing by vertex factors and permutations of identical vertices, the result is

$$\frac{1}{S} = \frac{6 \times 4 \times 6 \times 3 \times 2}{3! 3! 4! 2!} = \frac{1}{2}.$$

Complex scalar fields

The case of complex scalar fields can be considered as two independent fields, or equivalently as independent fields ϕ and ϕ^* . The generating functional in the interacting case can be written as

$$\begin{aligned} Z[J, J^*] &= \int \mathcal{D}\phi \mathcal{D}\phi^* \exp \left(i \int d^4x [\phi^* (-\partial_\mu \partial^\mu - M^2) \phi + \mathcal{L}_I(\phi) + J^* \phi + J \phi^*] \right) \\ &= \exp \left(i \int d^4z \mathcal{L}_I \left(\frac{1}{i} \frac{\delta}{\delta J(z)}, \frac{1}{i} \frac{\delta}{\delta J^*(z)} \right) \right) \\ &\quad \exp \left(- \int d^4x d^4y J^*(x) i\Delta_F(x-y) J(y) \right). \end{aligned} \tag{10.70}$$

In Feynman diagrams the propagator is still given by $i\Delta_F(k)$, but it connects a source with its complex conjugate and therefore is oriented, denoted

$$\begin{array}{c} \longrightarrow \mathbf{k} \\ \longrightarrow \mathbf{k} \end{array} = \frac{i}{k^2 - M^2 + i\epsilon}$$

Note that in this case the propagator does not have identical ends, i.e. there is no combinatorial factor like in the scalar case.

Dirac fields

For fermions the generating functional is given by

$$\begin{aligned} Z[\eta, \bar{\eta}] &= \int \mathcal{D}\bar{\psi} \mathcal{D}\psi \exp \left(i \int d^4x [\bar{\psi} (i\cancel{\partial} - M) \psi + \mathcal{L}_I(\psi) + \bar{\eta}\psi + \bar{\psi}\eta] \right) \\ &= \exp \left(i \int d^4z \mathcal{L}_I \left(\frac{1}{i} \frac{\delta}{\delta\eta(z)}, -\frac{1}{i} \frac{\delta}{\delta\bar{\eta}(z)} \right) \right) \\ &\quad \exp \left(- \int d^4x d^4y \bar{\eta}(x) iS_F(x-y) \eta(y) \right), \end{aligned} \quad (10.71)$$

where iS_F is the Feynman propagator for fermions, which is the solution of $(i\cancel{\partial} - M)S_F(x) = -\delta^4(x)$ (i.e. minus the inverse of the operator appearing in the quadratic piece) and is given by

$$\begin{aligned} iS_F(x-y) &= \langle 0 | \mathcal{T} \bar{\psi}(x) \psi(y) | 0 \rangle = \\ &= (i\cancel{\partial}_x + M) i\Delta_F(x-y) \end{aligned} \quad (10.72)$$


$$= \int \frac{d^4p}{(2\pi)^4} e^{-ip \cdot x} \frac{\not{p} + M}{p^2 - M^2 + i\epsilon}, \quad (10.73)$$

and the (oriented) propagator in Feynman diagrams involving fermions is

$$i \xrightarrow{\mathbf{p}} j = \left(\frac{i}{\not{p} - M + i\epsilon} \right)_{ji} = \frac{i(\not{p} + M)_{ji}}{p^2 - M^2 + i\epsilon}$$

The time ordered functions are obtained by functional derivatives from $Z(\eta, \bar{\eta})$, but the anticommuting properties of Grassmann variables imply some additional minus sign in Feynman diagrams, namely


(Rule 5) Feynman diagrams which only differ by exchanging identical fermions in initial or final state have a relative minus sign, e.g. in $e^-e^- \rightarrow e^-e^-$ scattering (Møller scattering) the lowest order contribution is



(see next section for $e^-e^-\gamma$ vertex).

(Rule 6) Each closed fermion loop gets a sign -1 .

The latter rule is illustrated in the example of an interaction term $\mathcal{L}_I = : g\bar{\psi}(x)\psi(x)\phi(x) :$ in an interacting theory with fermionic and scalar fields. The two-points Green's function $\langle 0 | \mathcal{T} \phi(x) \phi(y) | 0 \rangle$ contains a fermionic loop contribution,



which arises from the quadratic term in $\exp(i \int d^4z \mathcal{L}_I)$,

$$-\frac{g^2}{2} \int dz dz' \frac{\delta^2}{\delta\bar{\eta}(z) \delta\eta(z)} \frac{\delta^2}{\delta\bar{\eta}(z') \delta\eta(z')}$$

and the quadratic term in $Z_0[\eta, \bar{\eta}]$,





$$-\frac{1}{2} \int dx dy dx' dy' \bar{\eta}(x) S(x-y) \eta(y) \bar{\eta}(x') S(x'-y') \eta(y').$$

The result is

$$\propto g^2 S(z - z') S(z' - z) = -g^2 iS(z - z') iS(z' - z),$$

which contains an extra minus sign as compared to a bosonic loop.

The wave functions for fermions are given by

	$u_i(p)$	incoming fermion
	$\bar{u}_i(p)$	outgoing fermion
	$\bar{v}_i(p)$	incoming antifermion
	$v_i(p)$	outgoing antifermion

In writing down the expressions for Feynman diagrams one has to be aware that the wave functions and the propagators have more than one component. It is necessary to start at the end of a fermion line (with the above arrow convention an outgoing fermion wave function \bar{u} or an incoming antifermion wave function \bar{v}) and keep on following that line, writing down the propagators till the beginning of the line (incoming fermion u or outgoing antifermion v) has been reached. As an example some scattering processes in quantum electrodynamics will be discussed in the next section after the introduction of the Feynman rules for vector fields.

Vector fields and Quantum Electrodynamics



As the most important example of vector fields consider the lagrangian density for quantum electrodynamics (QED),

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{\lambda}{2} (\partial \cdot A)^2 + \bar{\psi}(i\cancel{\partial} - M)\psi - e \bar{\psi} \gamma^\mu \psi A_\mu. \tag{10.74}$$

In addition to the fermion propagator and fermion wave functions discussed in the previous section we have the photon propagator¹

$$\mu \text{ } \overset{\vec{k}}{\sim} \text{ } \nu = -i \left[\frac{g_{\mu\nu}}{k^2 + i\epsilon} - \left(1 - \frac{1}{\lambda}\right) \frac{k_\mu k_\nu}{(k^2 + i\epsilon)^2} \right],$$

where the $k_\mu k_\nu$ terms in the case where the photon couples to a conserved current (such as $\bar{\psi} \gamma^\mu \psi$) will not contribute. Particular choices of λ are $\lambda = 1$ (Feynman propagator or Feynman gauge) and $\lambda = \infty$ (Landau gauge). The wave functions are given by

	$\epsilon_\mu(k)$	incoming photon
	$\epsilon_\mu^*(k)$	outgoing photon

The vertex for the coupling of photon to the electron is given by

$$\begin{array}{c} \mu \\ | \\ \text{---} \\ / \quad \backslash \\ i \quad \quad j \end{array} = -i e (\gamma_\mu)_{ji}.$$


¹For a massive vector boson inversion of the quadratic term including the Lorentz constraint $\propto \frac{\lambda}{2} (\partial_\mu A^\mu)^2$ leads to the propagator

$$i D_{\mu\nu}(k) = i \left[\frac{-g_{\mu\nu}}{k^2 - M^2 + i\epsilon} + \left(1 - \frac{1}{\lambda}\right) \frac{k_\mu k_\nu}{(k^2 - M^2 + i\epsilon)(k^2 - \frac{1}{\lambda} M^2 + i\epsilon)} \right].$$

10.5 Some examples

$e\mu$ scattering

The first example is the electromagnetic scattering of an electron and a muon. To lowest order ($\propto \alpha = e^2/4\pi$) only one diagram contributes. The diagram and momenta and the commonly used invariants (Mandelstam variables) for a $2 \rightarrow 2$ scattering process are



$$\begin{aligned}
 s &= (k+p)^2 = m^2 + M^2 + 2k \cdot p \\
 &= (k'+p')^2 = m^2 + M^2 + 2k' \cdot p' \\
 t &= (k-k')^2 = q^2 = 2m^2 - 2k \cdot k' \\
 &= (p-p')^2 = 2M^2 - 2p \cdot p' \\
 u &= (k-p')^2 = m^2 + M^2 - 2k \cdot p' \\
 &= (k'-p)^2 = m^2 + M^2 - 2k' \cdot p \\
 s + t + u &= \sum m_i^2 = 2m^2 + 2M^2
 \end{aligned}$$

The scattering amplitude is given by

$$-i\mathcal{M} = \bar{u}(k', s_3)(-ie)\gamma^\mu u(k, s_1) \frac{-ig_{\mu\nu}}{q^2} \bar{u}(p', s_4)(-ie)\gamma^\nu u(p, s_2), \quad (10.75)$$

Note that the $q^\mu q^\nu$ term in the photon propagator are irrelevant because the photon couples to a conserved current. If we are interested in the scattering process of an unpolarized initial state and we are not interested in the spins in the final state we need $|\mathcal{M}|^2$ summed over spins in the final state (\sum_{s_3, s_4}) and averaged over spins in the initial state ($1/2 \times 1/2 \times \sum_{s_1, s_2}$) which can be written as (see also chapter 4)

$$\begin{aligned}
 |\mathcal{M}|^2 &= \frac{1}{4} \sum_{s_1, s_2, s_3, s_4} |\bar{u}(k', s_3)\gamma^\mu u(k, s_1) \frac{e^2}{q^2} \bar{u}(p', s_4)\gamma_\mu u(p, s_2)|^2 \\
 &= \frac{e^4}{q^4} L_{\mu\nu}^{(m)} L^{\mu\nu (M)}, \quad (10.76)
 \end{aligned}$$

[using that $(\bar{u}(k')\gamma_\mu u(k))^* = \bar{u}(k)\gamma_\mu u(k')$], where

$$\begin{aligned}
 L_{\mu\nu}^{(m)} &= \frac{1}{2} \sum_{s, s'} \bar{u}(k', s')\gamma_\mu u(k, s) \bar{u}(k, s)\gamma_\nu u(k', s') \\
 &= \frac{1}{2} \text{Tr}[(\not{k}' + m)\gamma_\mu(\not{k} + m)\gamma_\nu] \\
 &= 2[k_\mu k'_\nu + k_\nu k'_\mu - g_{\mu\nu}(k \cdot k' - m^2)] \\
 &= 2k_\mu k'_\nu + 2k_\nu k'_\mu + q^2 g_{\mu\nu} \quad (10.77)
 \end{aligned}$$

Combining $L_{\mu\nu}^{(m)}$ and $L^{\mu\nu (M)}$ one obtains


$$L_{\mu\nu}^{(m)} L^{\mu\nu (M)} = 2[s^2 + u^2 + 4t(M^2 + m^2) - 2(M^2 + m^2)^2] \quad (10.78)$$

and

$$\left| \frac{\mathcal{M}}{4\pi} \right|^2 = \frac{2\alpha^2}{t^2} [s^2 + u^2 + 4t(M^2 + m^2) - 2(M^2 + m^2)^2]. \quad (10.79)$$

$e^-e^+ \rightarrow \mu^-\mu^+$ scattering

The second example is the annihilation of an electron pair and creation of a muon pair. To lowest order ($\propto \alpha = e^2/4\pi$) only one diagram contributes. The diagram, the masses, momenta and invariants are



$$s = (k + k')^2 = (p + p')^2$$

$$t = (k - p)^2 = (k' - p')^2$$

$$u = (k - p')^2 = (k' - p)^2$$

The scattering amplitude squared (spins summed and averaged) is given by

$$\begin{aligned} |\mathcal{M}|^2 &= \frac{1}{4} \sum_{s_1, s_2, s_3, s_4} |\bar{u}(p, s_3) \gamma^\mu v(p', s_4) \frac{e^2}{s} \bar{v}(k', s_2) \gamma_\mu u(k, s_1)|^2 \\ &= \frac{e^4}{s^2} \left(\frac{1}{2} \sum_{s_1, s_2} \bar{v}(k', s_2) \gamma_\mu u(k, s_1) \bar{u}(k, s_1) \gamma_\nu v(k', s_2) \right) \\ &\quad \times \left(\frac{1}{2} \sum_{s_3, s_4} \bar{u}(p, s_3) \gamma^\mu v(p', s_4) \bar{v}(p', s_4) \gamma^\nu u(p, s_3) \right) \\ &= \frac{e^4}{s^2} \left(\frac{1}{2} \text{Tr}(\not{k} - m) \gamma_\mu (\not{k}' + m) \gamma_\nu \right) \left(\frac{1}{2} \text{Tr}(\not{p} + M) \gamma^\mu (\not{p}' - M) \gamma^\nu \right) \\ &= 4 \frac{e^4}{s^2} \left(k_\mu k'_\nu + k_\nu k'_\mu - \frac{1}{2} g_{\mu\nu} s \right) \left(p^\mu p'^\nu + p^\nu p'^\mu - \frac{1}{2} g^{\mu\nu} s \right) \end{aligned}$$

and

$$\left| \frac{\mathcal{M}}{4\pi} \right|^2 = \frac{2\alpha^2}{s^2} [t^2 + u^2 + 4s(M^2 + m^2) - 2(M^2 + m^2)^2]. \quad (10.80)$$

Note the similarity in the amplitudes for $e\mu$ scattering and $e^-e^+ \rightarrow \mu^-\mu^+$. Basically the same diagram is calculated and the result is the same after the interchange of $s \leftrightarrow t$. This is known as crossing symmetry. Similarly, for instance Møller scattering ($e^-e^- \rightarrow e^-e^-$) and Bhabha scattering ($e^-e^+ \rightarrow e^-e^+$) are related using crossing symmetry.

Exercises**Exercise 10.1**

- Give in diagrammatic notation the full Green functions $G^{(4)}(x_1, \dots, x_4)$ for the interacting case in ϕ^4 -theory to first order in the coupling constant g as obtained from the full expression for $Z[J]$ in section 10.2.
- Use the definition of the source-connected Green function $G_{sc}^{(4)}$ and show that indeed source-connected diagrams survive.
- Do the same for the connected Green function $G_c^{(4)}$.

Exercise 10.2

(a) Show that the translation properties of the fields and the vacuum imply

$$G^{(n)}(x_1 + a, \dots, x_n + a) = G^{(n)}(x_1, \dots, x_n).$$

(b) Show (by using x_1 as shift-variable) that this implies that

$$\int \prod_{i=1}^n d^4x_i e^{i p_i \cdot x_i} G^{(n)}(x_1, \dots, x_n) \propto (2\pi)^4 \delta^4(p_1 + \dots + p_n),$$

hence we can write

$$\int \prod_{i=1}^n d^4x_i e^{i p_i \cdot x_i} G^{(n)}(x_1, \dots, x_n) \equiv (2\pi)^4 \delta^4(p_1 + \dots + p_n) G^{(n)}(p_1, \dots, p_n),$$

which means overall momentum conservation in Green functions in momentum space.

Exercise 10.3

Show that the combinatorial factors found using the rules given in section 10.3 reproduce for the diagrams



precisely the combinatorial factors that appear in Eqs 10.37 and 10.38.

Exercise 10.4

(a) Write down the Feynman diagrams contributing to electron-electron scattering, $e(p_1) + e(p_2) \rightarrow e(p'_1) + e(p'_2)$ in lowest order in α . It is of the form $-i \mathcal{M} = A_1 - A_2$.

(b) Calculate the quadratic pieces and interference terms,

$$|\mathcal{M}|^2 = T_{11} + T_{22} - T_{12} - T_{21},$$

in the amplitude ($T_{ij} = A_i^* A_j$). Express the contributions in invariants s , t and u . Show that the amplitude is symmetric under the interchange of $t \leftrightarrow u$.

Chapter 11

Scattering theory

11.1 kinematics in scattering processes

Phase space

The 1-particle state is denoted $|p\rangle$. It is determined by the energy-momentum four vector $p = (E, \mathbf{p})$ which satisfies $p^2 = E^2 - \mathbf{p}^2 = m^2$. A physical state has positive energy. The phase space is determined by the weight factors assigned to each state in the summation or integration over states, i.e. the 1-particle phase space is

$$\int \frac{d^3\mathbf{p}}{(2\pi)^3 2E} = \int \frac{d^4p}{(2\pi)^4} \theta(p^0) (2\pi) \delta(p^2 - m^2), \quad (11.1)$$

(proven in Chapter 2). This is generalized to the multi-particle phase space

$$d\mathcal{R}(p_1, \dots, p_n) = \prod_{i=1}^n \frac{d^3p_i}{(2\pi)^3 2E_i}, \quad (11.2)$$

and the *reduced phase space element* by

$$d\mathcal{R}(s, p_1, \dots, p_n) = (2\pi)^4 \delta^4(P - \sum_i p_i) d\mathcal{R}(p_1, \dots, p_n), \quad (11.3)$$

which is useful because the total 4-momentum of the final state usually is fixed by overall momentum conservation. Here s is the invariant mass of the n-particle system, $s = (p_1 + \dots + p_n)^2$. It is a useful quantity, for instance for determining the threshold energy for the production of a final state $1 + 2 + \dots + n$. In the CM frame the threshold value for s obviously is

$$s_{\text{threshold}} = \left(\sum_{i=1}^n m_i \right)^2. \quad (11.4)$$

For two particle states $|p_a, p_b\rangle$ we start with the four vectors $p_a = (E_a, \mathbf{p}_a)$ and $p_b = (E_b, \mathbf{p}_b)$ satisfying $p_a^2 = m_a^2$ and $p_b^2 = m_b^2$, and the total momentum four-vector $P = p_a + p_b$. For two particles, the quantity

$$s = P^2 = (p_a + p_b)^2, \quad (11.5)$$

is referred to as the invariant mass squared. Its square root, \sqrt{s} is for obvious reasons known as the center of mass (CM) energy.

To be specific let us consider two frequently used frames. The first is the CM system. In that case

$$p_a = (E_a^{\text{cm}}, \mathbf{q}), \quad (11.6)$$

$$p_b = (E_b^{\text{cm}}, -\mathbf{q}). \quad (11.7)$$

It is straightforward to prove that the unknowns in the particular system can be expressed in the invariants (m_a , m_b and s). Prove that

$$|\mathbf{q}| = \sqrt{\frac{(s - m_a^2 - m_b^2)^2 - 4m_a^2 m_b^2}{4s}} = \sqrt{\frac{\lambda(s, m_a^2, m_b^2)}{4s}}, \quad (11.8)$$

$$E_a^{\text{cm}} = \frac{s + m_a^2 - m_b^2}{2\sqrt{s}}, \quad (11.9)$$

$$E_b^{\text{cm}} = \frac{s - m_a^2 + m_b^2}{2\sqrt{s}}. \quad (11.10)$$

The function $\lambda(s, m_a^2, m_b^2)$ is a function symmetric in its three arguments, which in the specific case also can be expressed as $\lambda(s, m_a^2, m_b^2) = 4(p_a \cdot p_b)^2 - 4p_a^2 p_b^2$.

The second frame considered explicitly is the so-called target rest frame in which one of the particles (called the target) is at rest. In that case

$$p_a = (E_a^{\text{trf}}, \mathbf{p}_a^{\text{trf}}), \quad (11.11)$$

$$p_b = (m_b, \mathbf{0}), \quad (11.12)$$

Also in this case one can express the energy and momentum in the invariants. Prove that

$$E_a^{\text{trf}} = \frac{s - m_a^2 - m_b^2}{2m_b}, \quad (11.13)$$

$$|\mathbf{p}_a^{\text{trf}}| = \frac{\sqrt{\lambda(s, m_a^2, m_b^2)}}{2m_b}. \quad (11.14)$$

One can, for instance, use the first relation and the abovementioned threshold value for s to calculate the threshold for a specific n -particle final state in the target rest frame,

$$E_a^{\text{trf}}(\text{threshold}) = \frac{1}{2m_b} \left(\left(\sum_i m_i \right)^2 - m_a^2 - m_b^2 \right). \quad (11.15)$$

Explicit calculation of the reduced two-body phase space element gives

$$\begin{aligned} d\mathcal{R}(s, p_1, p_2) &= \frac{1}{(2\pi)^2} \frac{d^3 p_1}{2E_1} \frac{d^3 p_2}{2E_2} \delta^4(P - p_1 - p_2) \\ &\stackrel{\text{CM}}{=} \frac{1}{(2\pi)^2} \frac{d^3 q}{4E_1 E_2} \delta(\sqrt{s} - E_1 - E_2) \\ &= \frac{1}{(2\pi)^2} d\Omega(\hat{q}) \frac{\mathbf{q}^2 d|\mathbf{q}|}{4E_1 E_2} \delta(\sqrt{s} - E_1 - E_2) \end{aligned}$$

which using $|\mathbf{q}| d|\mathbf{q}| = (E_1 E_2 / (E_1 + E_2)) d(E_1 + E_2)$ gives

$$\begin{aligned} d\mathcal{R}(s, p_1, p_2) &= \frac{|\mathbf{q}|}{(2\pi)^2} d\Omega(\hat{q}) \frac{d(E_1 + E_2)}{4(E_1 + E_2)} \delta(\sqrt{s} - E_1 - E_2) \\ &= \frac{|\mathbf{q}|}{4\pi\sqrt{s}} \frac{d\Omega(\hat{q})}{4\pi} = \frac{\sqrt{\lambda_{12}}}{8\pi s} \frac{d\Omega(\hat{q})}{4\pi}, \end{aligned} \quad (11.16)$$

where λ_{12} denotes $\lambda(s, m_1^2, m_2^2)$.

Kinematics of $2 \rightarrow 2$ scattering processes

The simplest scattering process is 2 particles in and 2 particles out. Examples appear in

$$\pi^- + p \rightarrow \pi^- + p \quad (11.17)$$

$$\rightarrow \pi^0 + n \quad (11.18)$$

$$\rightarrow \pi^+ + \pi^- + n \quad (11.19)$$

$$\rightarrow \dots \quad (11.20)$$

The various possibilities are referred to as different reaction channels, where the first is referred to as elastic channel and the set of all other channels as the inelastic channels. Of course there are not only 2-particle channels. The initial state, however, usually is a 2-particle state, while the final state often arises from a series of 2-particle processes combined with the decay of an intermediate particle (resonance).

Consider the process $a + b \rightarrow c + d$. An often used set of invariants are the Mandelstam variables,

$$s = (p_a + p_b)^2 = (p_c + p_d)^2 \quad (11.21)$$

$$t = (p_a - p_c)^2 = (p_b - p_d)^2 \quad (11.22)$$

$$u = (p_a - p_d)^2 = (p_b - p_c)^2 \quad (11.23)$$

which are not independent as $s + t + u = m_a^2 + m_b^2 + m_c^2 + m_d^2$. The variable s is always larger than the minimal value $(m_a + m_b)^2$. A specific reaction channel starts contributing at the threshold value (Eq. 11.4). Instead of the scattering angle, which for the above $2 \rightarrow 2$ process in the case of azimuthal symmetry is defined as $\hat{p}_a \cdot \hat{p}_c = \cos \theta$ one can use in the CM the invariant

$$t \equiv (p_a - p_c)^2 \stackrel{\text{CM}}{=} m_a^2 + m_c^2 - 2 E_a E_c + 2 q q' \cos \theta_{\text{cm}},$$

with $q = \sqrt{\lambda_{ab}/4s}$ and $q' = \sqrt{\lambda_{cd}/4s}$. The minimum and maximum values for t correspond to θ_{cm} being 0 or 180 degrees,

$$\begin{aligned} t_{\min}^{\max} &= m_a^2 + m_c^2 - 2 E_a E_c \pm 2 q q' \\ &= m_a^2 + m_c^2 - \frac{(s + m_a^2 - m_b^2)(s + m_c^2 - m_d^2)}{2s} \pm \frac{\sqrt{\lambda_{ab}\lambda_{cd}}}{2s}. \end{aligned} \quad (11.24)$$

Using the relation between t and $\cos \theta_{\text{cm}}$ it is straightforward to express $d\Omega_{\text{cm}}$ in dt , $dt = 2 q q' d \cos \theta_{\text{cm}}$ and obtain for the two-body phase space element

$$d\mathcal{R}(s, p_c, p_d) = \frac{q'}{4\pi\sqrt{s}} \frac{d\Omega_{\text{cm}}}{4\pi} = \frac{\sqrt{\lambda_{cd}}}{8\pi s} \frac{d\Omega_{\text{cm}}}{4\pi} \quad (11.25)$$

$$= \frac{dt}{8\pi\sqrt{\lambda_{ab}}} = \frac{dt}{16\pi q\sqrt{s}}. \quad (11.26)$$

Kinematics of inclusive hard scattering processes

In high energy (hard) scattering processes, usually many particles are produced. In inclusive measurements no particles are detected in the final state, in exclusive measurements all particles are detected. Consider the 1-particle inclusive case, in which one particle is detected, $H_1 + H_2 \rightarrow h + X$. At high energies, there is usually a preferred direction, for instance the momenta of incoming (colliding) hadrons and it is useful to use for the produced particle rapidity as a variable. Writing

$$E = m_T \cosh y, \quad p_z = m_T \sinh y, \quad p_T = (p_x, p_y), \quad (11.27)$$

with $m_T^2 = m^2 - p_T^2 = m^2 + p_x^2 + p_y^2$, the rapidity y is defined

$$y = \frac{1}{2} \ln \left(\frac{E + p_z}{E - p_z} \right) = \ln \left(\frac{E + p_z}{m_T} \right) = \tanh^{-1} \left(\frac{p_z}{E} \right).$$

It is convenient because under a boost (along z) with velocity β it changes as $y \rightarrow y - \tanh^{-1} \beta$, which means that rapidity distributions dN/dy maintain their shape. For large energies and not too small scattering angles ($\theta \gg 1/\gamma$) y is approximately equal to the pseudo-rapidity η ,

$$\eta = -\ln(\tan(\theta/2)) = \tanh^{-1}(\cos \theta), \quad (11.28)$$

which (only involving angles) is easier to determine. The one-particle phase space in terms of these variables becomes


$$\frac{d^3 p}{(2\pi)^3 2E} = \frac{dy d^2 p_T}{2(2\pi)^3} = \frac{dy d|p_T|^2 d\phi}{16\pi^2 2\pi} = \frac{|\mathbf{p}|}{E} \frac{d\eta d|p_T|^2 d\phi}{16\pi^2 2\pi}, \quad (11.29)$$

with at high energies the factor $|\mathbf{p}|/E \simeq 1$.

11.2 Crossing symmetry

In the previous chapter, we have seen that the amplitudes for the processes $e^- \mu^- \rightarrow e^- \mu^-$ and the process $e^- e^+ \rightarrow \mu^+ \mu^-$ are simply related by an interchange of the variables s and t . This is known as crossing symmetry.

Given a two-to-two scattering process $ab \rightarrow cd$ one can relate the processes



They are referred to as s-channel, t-channel and u-channel processes respectively. With the momenta defined as in the figures above one has for all these processes the same amplitude $\mathcal{M}(s, t, u)$ with

$$\begin{aligned} s &= (p_1 + p_2)^2 = (p_3 + p_4)^2, \\ t &= (p_1 - p_3)^2 = (p_2 - p_4)^2, \\ u &= (p_1 - p_4)^2 = (p_2 - p_3)^2. \end{aligned}$$

These variables are precisely the Mandelstam variables for the s-channel process ($ab \rightarrow cd$). For the t-channel process ($a\bar{c} \rightarrow \bar{b}d$) one has

$$\begin{aligned} s_t &= (p_a + p_{\bar{c}})^2 = t, \\ t_t &= (p_a - p_{\bar{b}})^2 = s, \\ u_t &= (p_a - p_d)^2 = u, \end{aligned}$$

while for the u-channel process ($a\bar{d} \rightarrow c\bar{b}$) one has

$$\begin{aligned} s_u &= (p_a + p_{\bar{d}})^2 = u, \\ t_u &= (p_a - p_c)^2 = t, \\ u_u &= (p_a - p_{\bar{b}})^2 = s. \end{aligned}$$

Analyticity of the field theoretically calculated result implies


$$\mathcal{M}_{a\bar{c} \rightarrow \bar{b}d}(s_t = t, t_t = s, u_t = u) = \mathcal{M}_{ab \rightarrow cd}(s, t, u), \quad (11.30)$$

$$\mathcal{M}_{a\bar{d} \rightarrow c\bar{b}}(s_u = u, t_u = s, u_u = s) = \mathcal{M}_{ab \rightarrow cd}(s, t, u). \quad (11.31)$$

One can also phrase it in the following way: one has a single analytic function $\mathcal{M}_{ab \rightarrow cd}(s, t, u)$ that represents physical amplitudes in the physical regions for three scattering processes. To see this one can make a two-dimensional plot for the variables s , t and u . This is a consequence of the constraint $s + t + u = M_a^2 + M_b^2 + M_c^2 + M_d^2$. To find the physical regions one looks for the boundaries of

$$\begin{aligned}\cos\theta_s &= \frac{1}{\sqrt{\lambda_{ab}^s \lambda_{cd}^s}} [s(t-u) + (M_a^2 - M_b^2)(M_c^2 - M_d^2)], \\ \cos\theta_t &= \frac{1}{\sqrt{\lambda_{ac}^t \lambda_{bd}^t}} [t(s-u) + (M_a^2 - M_c^2)(M_b^2 - M_d^2)], \\ \cos\theta_u &= \frac{1}{\sqrt{\lambda_{ad}^u \lambda_{cb}^u}} [u(t-s) + (M_a^2 - M_d^2)(M_c^2 - M_b^2)].\end{aligned}$$

This defines the boundaries of the physical regions, shown below for the case of equal masses.



11.3 Cross sections and lifetimes

Scattering process

For a scattering process $a + b \rightarrow c + \dots$ (consider for convenience the rest frame for the target, say b) the cross section $\sigma(a + b \rightarrow c + \dots)$ is defined as the proportionality factor in

$$\frac{N_c}{T} = \sigma(a + b \rightarrow c + \dots) \cdot N_b \cdot \text{flux}(a),$$

where V and T indicate the volume and the time in which the experiment is performed, N_c/T indicates the number of particles c detected in the scattering process, N_b indicates the number of (target) particles b , which for a density ρ_b is given by $N_b = \rho_b \cdot V$, while the flux of the beam particles a is $\text{flux}(a) = \rho_a \cdot v_a^{\text{trf}}$. The proportionality factor has the dimension of area and is called the cross section, i.e.

$$\sigma = \frac{N}{T \cdot V} \frac{1}{\rho_a \rho_b v_a^{\text{trf}}}. \quad (11.32)$$

Although this at first sight does not look covariant, it is. N and $T \cdot V$ are covariant. Using $\rho_a^{\text{trf}} = \rho_a^{(0)} \cdot \gamma_a = \rho_a^{(0)} \cdot E_a^{\text{lab}}/m_a$ (where $\rho_a^{(0)}$ is the rest frame density) and $v_a^{\text{lab}} = p_a^{\text{lab}}/E_a^{\text{lab}}$ we have

$$\rho_a \rho_b v_a^{\text{trf}} = \frac{\rho_a^{(0)} \rho_b^{(0)}}{4 m_a m_b} 2\sqrt{\lambda_{ab}}$$

or with $\rho_a^{(0)} = 2 m_a$,

$$\sigma = \frac{1}{2\sqrt{\lambda_{ab}}} \frac{N}{T \cdot V}. \quad (11.33)$$

Decay of particles

For the decay of particle a one has macroscopically

$$\frac{dN}{dt} = -\Gamma N, \quad (11.34)$$

i.e. the amount of decaying particles is proportional to the number of particles with proportionality factor the em decay width Γ . From the solution

$$N(t) = N(0) e^{-\Gamma t} \quad (11.35)$$

one knows that the decay time $\tau = 1/\Gamma$. Microscopically one has

$$\frac{N_{\text{decay}}}{T} = N_a \cdot \Gamma$$

or

$$\Gamma = \frac{N}{T \cdot V} \frac{1}{\rho_a}. \quad (11.36)$$

This quantity is not covariant, as expected. The decay time for moving particles τ is related to the decay time in the rest frame of that particle (the proper decay time τ_0) by $\tau = \gamma \tau_0$. For the (proper) decay width one thus has

$$\Gamma_0 = \frac{1}{2 m_a} \frac{N}{T \cdot V}. \quad (11.37)$$

Fermi's Golden Rule

In both the scattering cross section and the decay constant the quantity N/TV appears. For this we employ in essence *Fermi's Golden rule* stating that when the S -matrix element is written as

$$S_{fi} = \delta_{fi} - (2\pi)^4 \delta^4(P_i - P_f) i \mathcal{M}_{fi} \quad (11.38)$$

(in which we can calculate $-i \mathcal{M}_{fi}$ using Feynman diagrams), the number of scattered or decayed particles is given by

$$N = |(2\pi)^4 \delta^4(P_i - P_f) i \mathcal{M}_{fi}|^2 d\mathcal{R}(p_1, \dots, p_n). \quad (11.39)$$

One of the δ functions can be rewritten as $T \cdot V$ (remember the normalization of plane waves),

$$\begin{aligned} & |(2\pi)^4 \delta^4(P_i - P_f)|^2 \\ &= (2\pi)^4 \delta^4(P_i - P_f) \int_{V,T} d^4x e^{i(P_i - P_f) \cdot x} \\ &= (2\pi)^4 \delta^4(P_i - P_f) \int_{V,T} d^4x = V \cdot T (2\pi)^4 \delta^4(P_i - P_f). \end{aligned}$$

(Using normalized wave packets these somewhat ill-defined manipulations can be made more rigorous). The result is

$$\frac{N}{T \cdot V} = |\mathcal{M}_{fi}|^2 d\mathcal{R}(s, p_1, \dots, p_n). \quad (11.40)$$

Combining this with the expressions for the width or the cross section one obtains for the decay width

$$\Gamma = \frac{1}{2m} \int d\mathcal{R}(m^2, p_1, \dots, p_n) |\mathcal{M}|^2 \quad (11.41)$$

$$\stackrel{\text{2-body decay}}{=} \frac{q}{32\pi^2 m^2} \int d\Omega |\mathcal{M}|^2. \quad (11.42)$$

The *differential cross section* (final state not integrated over) is given by

$$d\sigma = \frac{1}{2\sqrt{\lambda_{ab}}} |\mathcal{M}_{fi}|^2 d\mathcal{R}(s, p_1, \dots, p_n), \quad (11.43)$$

and for instance for two particles

$$d\sigma = \frac{q'}{q} \left| \frac{\mathcal{M}(s, \theta_{\text{cm}})}{8\pi\sqrt{s}} \right|^2 d\Omega_{\text{cm}} = \frac{\pi}{\lambda_{ab}} \left| \frac{\mathcal{M}(s, t)}{4\pi} \right|^2 dt. \quad (11.44)$$

This can be used to get the full expression for $d\sigma/dt$ for $e\mu$ and e^+e^- scattering, for which the amplitudes squared have been calculated in the previous chapter. The amplitude $-\mathcal{M}/8\pi\sqrt{s}$ is the one to be compared with the quantum mechanical scattering amplitude $f(E, \theta)$, for which one has $d\sigma/d\Omega = |f(E, \theta)|^2$. The sign difference comes from the (conventional) sign in relation between S and quantummechanical and relativistic scattering amplitude, respectively.

11.4 Unitarity condition

The unitarity of the S -matrix, i.e.

$$(S^\dagger)_{fn} S_{ni} = \delta_{fi}$$

implies for the scattering matrix \mathcal{M} ,

$$[\delta_{fn} + i(2\pi)^4 \delta^4(P_f - P_n) (\mathcal{M}^\dagger)_{fn}] [\delta_{ni} - i(2\pi)^4 \delta^4(P_i - P_n) \mathcal{M}_{ni}] = \delta_{fi},$$

or

$$-i [\mathcal{M}_{fi} - (\mathcal{M}^\dagger)_{fi}] = - \sum_n (\mathcal{M}^\dagger)_{fn} (2\pi)^4 \delta^4(P_i - P_n) \mathcal{M}_{ni}. \quad (11.45)$$

Since the amplitudes also depend on all momenta the full result for two-particle intermediate states is (in CM, see 11.25)

$$-i [\mathcal{M}_{fi} - (\mathcal{M}^\dagger)_{fi}] = - \sum_n \int d\Omega(\hat{q}_n) \mathcal{M}_{nf}^*(\mathbf{q}_f, \mathbf{q}_n) \frac{q_n}{16\pi^2 \sqrt{s}} \mathcal{M}_{ni}(\mathbf{q}_i, \mathbf{q}_n). \quad (11.46)$$

Partial wave expansion

Often it is useful to make a partial wave expansion for the amplitude $\mathcal{M}(s, \theta)$ or $\mathcal{M}(\mathbf{q}_i, \mathbf{q}_f)$,

$$\mathcal{M}(s, \theta) = -8\pi\sqrt{s} \sum_\ell (2\ell + 1) M_\ell(s) P_\ell(\cos \theta), \quad (11.47)$$

(in analogy with the expansion for $f(E, \theta)$ in quantum mechanics; note the sign and $\cos \theta = \hat{q}_i \cdot \hat{q}_f$). Inserted in the unitarity condition for \mathcal{M} ,

$$i \left[\frac{\mathcal{M}}{8\pi\sqrt{s}} - \frac{\mathcal{M}^\dagger}{8\pi\sqrt{s}} \right]_{fi} = \sum_n \int d\Omega_n \frac{\mathcal{M}_{nf}^*}{8\pi\sqrt{s}} \frac{q_n}{2\pi} \frac{\mathcal{M}_{ni}}{8\pi\sqrt{s}},$$

we obtain

$$\text{LHS} = -i \sum_{\ell} (2\ell + 1) P_{\ell}(\hat{q}_i \cdot \hat{q}_f) \left((M_{\ell})_{fi} - (M_{\ell}^{\dagger})_{fi} \right),$$

while for the RHS use is made of

$$P_{\ell}(\hat{q} \cdot \hat{q}') = \sum_m \frac{4\pi}{2\ell + 1} Y_m^{(\ell)}(\hat{q}) Y_m^{(\ell)*}(\hat{q}')$$

and the orthogonality of the $Y_m^{(\ell)}$ functions to prove that

$$\text{RHS} = 2 \sum_n \sum_{\ell} (2\ell + 1) P_{\ell}(\hat{q}_i \cdot \hat{q}_f) (M_{\ell}^{\dagger})_{fn} q_n (M_{\ell})_{ni},$$

i.e.

$$-i \left((M_{\ell})_{fi} - (M_{\ell}^{\dagger})_{fi} \right) = 2 \sum_n (M_{\ell}^{\dagger})_{fn} q_n (M_{\ell})_{ni}. \quad (11.48)$$

If only one channel is present this simplifies to

$$-i (M_{\ell} - M_{\ell}^*) = 2 q M_{\ell}^* M_{\ell}, \quad (11.49)$$

or $\text{Im } M_{\ell} = q |M_{\ell}|^2$, which allows writing

$$M_{\ell}(s) = \frac{S_{\ell}(s) - 1}{2i q} = \frac{e^{2i \delta_{\ell}(s)} - 1}{2i q}, \quad (11.50)$$

where $S_{\ell}(s)$ satisfies $|S_{\ell}(s)| = 1$ and $\delta_{\ell}(s)$ is the phase shift.

In general a given channel has $|S_{\ell}(s)| \leq 1$, parametrized as $S_{\ell}(s) = \eta_{\ell}(s) \exp(2i \delta_{\ell}(s))$. Using

$$\sigma = \int d\Omega \frac{q'}{q} \left| \frac{\mathcal{M}(s, \theta_{\text{cm}})}{8\pi \sqrt{s}} \right|^2,$$

in combination with the partial wave expansion for the amplitudes \mathcal{M} and the orthogonality of the Legendre polynomials immediately gives for the elastic channel,

$$\begin{aligned} \sigma_{\text{el}} &= 4\pi \sum_{\ell} (2\ell + 1) |M_{\ell}(s)|^2 \\ &= \frac{4\pi}{q^2} \sum_{\ell} (2\ell + 1) \left| \frac{\eta_{\ell} e^{2i \delta_{\ell}} - 1}{2i} \right|^2, \end{aligned} \quad (11.51)$$

and for the case that this is the only channel (purely elastic scattering, $\eta = 1$) the result

$$\sigma_{\text{el}} = \frac{4\pi}{q^2} \sum_{\ell} (2\ell + 1) \sin^2 \delta_{\ell}. \quad (11.52)$$

From the imaginary part of $\mathcal{M}(s, 0)$, the total cross section can be determined. Show that

$$\begin{aligned} \sigma_T &= \frac{4\pi}{q} \sum_{\ell} (2\ell + 1) \text{Im } M_{\ell}(s) \\ &= \frac{2\pi}{q^2} \sum_{\ell} (2\ell + 1) (1 - \eta_{\ell} \cos 2\delta_{\ell}). \end{aligned} \quad (11.53)$$

The difference is the inelastic cross section,

$$\sigma_{\text{inel}} = \frac{\pi}{q^2} \sum_{\ell} (2\ell + 1) (1 - \eta_{\ell}^2). \quad (11.54)$$

Note that the total cross section is maximal in the case of full absorption, $\eta = 0$, in which case, however, $\sigma_{\text{el}} = \sigma_{\text{inel}}$.

We note that unitarity is generally broken in a finite order calculation. For instance the tree-level calculation of Feynman diagrams at order e^2 was real. The product of amplitudes is of order e^4 , but still real. To check unitarity we need the amplitude at order e^4 . We can also invert the situation and use unitarity to find the imaginary part of an amplitude at some order from a lower order calculation. This is what we will do to discuss decay widths.

11.5 Unstable particles

For a stable particle the propagator is

$$i\Delta(k) = \frac{i}{k^2 - M^2 + i\epsilon} \quad (11.55)$$

(note that we have disregarded spin). The prescription for the pole structure, i.e. one has poles at $k^0 = \pm(E_k - i\epsilon)$ where $E_k = +\sqrt{\mathbf{k}^2 + M^2}$ guarantees the correct behavior, specifically one has for $t > 0$ that the Fourier transform is

$$\int dk^0 e^{-ik^0 t} \Delta(k) \propto \int dk^0 \frac{e^{-ik^0 t}}{(k^0 - E_k + i\epsilon)(k^0 + E_k - i\epsilon)} \stackrel{(t>0)}{\propto} e^{-iE_k t},$$

i.e. $\propto U(t, 0)$, the time-evolution operator. For an unstable particle one expects that

$$U(t, 0) \propto e^{-i(E - i\Gamma/2)t},$$

such that $|U(t, 0)|^2 = e^{-\Gamma t}$. This is achieved with a propagator

$$i\Delta_R(k) = \frac{i}{k^2 - M^2 + iM\Gamma} \quad (11.56)$$

(again disregarding spin). The quantity Γ is precisely the width for unstable particles. This is (somewhat sloppy!) seen by considering the (amputated) 1PI two-point vertex

$$\Gamma^{(2)} = \frac{-i}{\Delta}$$

as the amplitude $-i\mathcal{M}$ for scattering a particle into itself (forward!) through the decay channels as intermediate states. The unitarity condition then states

$$\begin{aligned} 2 \text{Im} \Delta_R^{-1}(k) &= \sum_n \int d\mathcal{R}(p_1, \dots, p_n) \mathcal{M}_{Rn}^\dagger (2\pi)^4 \delta^4(k - P_n) \mathcal{M}_{nR} \\ &= 2M \sum_n \Gamma_n = 2M\Gamma. \end{aligned} \quad (11.57)$$

This shows that Γ is the width of the resonance, which is given by a sum of the partial widths into the different channels. It is important to note that the physical width of a particle is the imaginary part of the two-point vertex at $s = M^2$.

For the amplitude in a scattering process going through a resonance, it is straightforward to write down the partial wave amplitude,

$$(q M_\ell)_{ij}(s) = \frac{-M \sqrt{\Gamma_i \Gamma_j}}{s - M^2 + iM\Gamma}. \quad (11.58)$$

(Prove this using the unitarity condition for partial waves). From this one sees that a resonance has the same shape in all channels but different strength. Limiting ourselves to a resonance in one channel, it is furthermore easy to prove that the cross section is given by

$$\sigma_{\text{el}} = \frac{4\pi}{q^2} (2\ell + 1) \frac{M^2 \Gamma^2}{(s - M^2)^2 + M^2 \Gamma^2}, \quad (11.59)$$

reaching the unitarity limit for $s = M^2$, where furthermore $\sigma_{\text{inel}} = 0$. This characteristic shape of a resonance is called the Breit-Wigner shape. The half-width of the resonance is $M\Gamma$. The phase shift in the resonating channel near the resonance is given by

$$\tan \delta_\ell(s) = \frac{M\Gamma}{M^2 - s}, \quad (11.60)$$

showing that the phase shift at resonance rises through $\delta = \pi/2$ with a 'velocity' $\partial\delta/\partial s = 1/M\Gamma$, i.e. a fast change in the phase shift for a narrow resonance. Note that because of the presence of a background the phase shift at resonance may actually be shifted.

Three famous resonances are:

- The Δ -resonance seen in pion-nucleon scattering. Its mass is $M = 1232$ MeV, its width $\Gamma = 120$ MeV. At resonance the cross section $\sigma_T(\pi^+p)$ is about 210 mb. The cross section $\sigma_T(\pi^-p)$ also shows a resonance with the same width with a value of about 70 mb. This implies that the resonance has spin $J = 3/2$ (decaying in a P-wave ($\ell = 1$) pion-nucleon state) and isospin $I = 3/2$ (the latter under the assumption that isospin is conserved for the strong interactions).
- The J/ψ resonance in e^+e^- scattering. This is a narrow resonance discovered in 1974. Its mass is $M = 3096.88$ MeV, the full width is $\Gamma = 88$ keV, the partial width into e^+e^- is $\Gamma_{ee} = 5.26$ keV.
- The Z^0 resonance in e^+e^- scattering with $M = 91.2$ GeV, $\Gamma = 2.49$ GeV. Essentially this resonance can decay into quark-antiquark pairs or into pairs of charged leptons. All these decays can be seen and leave an 'invisible' width of 498 MeV, which is attributed to neutrinos. Knowing that each neutrino contributes about 160 MeV (see next chapter), one can reconstruct the resonance shape for different numbers of neutrino species. Three neutrinos explain the resonance shape. The cross section at resonance is about 30 nb.

Exercises

Exercise 11.1

Show that the cross section for electron-electron scattering (exercise 10.4) can be written as

$$\frac{d\sigma}{dt} = \frac{4\pi\alpha^2}{s(s - 4m^2)} \{f(t, u) + g(t, u) + f(u, t) + g(u, t)\},$$

with

$$\begin{aligned} f(t, u) &= \frac{1}{t^2} \left[\frac{1}{2}(s^2 + u^2) + 4m^2(t - m^2) \right] \\ g(t, u) &= \frac{2}{tu} \left[\left(\frac{1}{2}s - m^2\right) \left(\frac{1}{2}s - 3m^2\right) \right] \end{aligned}$$

Exercise 11.2

Calculate the flux factor,

$$\frac{1}{\rho_a^{\text{cm}} \rho_b^{\text{cm}} |v_a^{\text{cm}} - v_b^{\text{cm}}|},$$

in the center of mass system for two colliding particles a and b,

Exercise 11.3

Complete the calculation of the decay width and lifetime of the pion by including the appropriate phase space factors and using the calculation of the squared matrix element in Exc. 4.8. Compare the results with the experimental π^- lifetime $\tau = 2.6 \times 10^{-8}$ s and the relative decay widths $\Gamma(\pi^- \rightarrow \mu^- \bar{\nu}_\mu) = 0.999877\Gamma$ and $\Gamma(\pi^- \rightarrow e^- \bar{\nu}_e) = 0.000123\Gamma$. Calculate now f_π , a quantity which we will encounter in the next chapter (section 12.2).

Exercise 11.4

The Δ resonance in $N\pi$ scattering appears in one partial wave (a P-wave) as a pole at the position $s = M_\Delta^2 - iM_\Delta\Gamma_\Delta$. As discussed for unstable particles, we have here already used unitarity to find a finite imaginary part. Show that unitarity can be used one step further to fix also the numerator of the amplitude near the pole,

$$M_1(s) = \frac{-M_\Delta\Gamma_\Delta/q}{s - M_\Delta^2 + iM_\Delta\Gamma_\Delta},$$

when Γ_Δ is approximately constant near this pole.

Chapter 12

The standard model

12.1 Non-abelian gauge theories

In chapter 10 we have considered quantum electrodynamics as an example of a gauge theory. The photon field A_μ was introduced as to render the lagrangian invariant under local gauge transformations. The extension to non-abelian gauge theories is straightforward. The symmetry group is a Lie-group G generated by generators T_a , which satisfy commutation relations

$$[T_a, T_b] = i c_{abc} T_c, \quad (12.1)$$

with c_{abc} known as the *structure constants* of the group. For a compact Lie-group they are antisymmetric in the three indices. In an abelian group the structure constants would be zero (for instance the trivial example of $U(1)$). Consider a field transforming under the group,

$$\phi(x) \longrightarrow \underline{U}(\theta) \phi(x) = e^{i\theta^a L_a} \phi(x) \stackrel{\text{inf}}{\equiv} (1 + i\theta^a L_a) \phi(x) \quad (12.2)$$

where L_a is a representation matrix for the representation to which ϕ belongs, i.e. for a three-component field (written as $\vec{\phi}$) under an $SO(3)$ or $SU(2)$ symmetry transformation,

$$\vec{\phi} \longrightarrow \underline{U}(\vec{\theta}) \vec{\phi} = e^{i\vec{\theta} \cdot \vec{L}} \vec{\phi} \stackrel{\text{inf}}{\equiv} \vec{\phi} - \vec{\theta} \times \vec{\phi}. \quad (12.3)$$

The complication arises (as in the abelian case) when one considers for a lagrangian density $\mathcal{L}(\phi, \partial_\mu \phi)$ the behavior of $\partial_\mu \phi$ under a *local* gauge transformation, $\underline{U}(\theta) = e^{i\theta^a(x)L_a}$,

$$\phi(x) \longrightarrow \underline{U}(\theta)\phi(x), \quad (12.4)$$

$$\partial_\mu \phi(x) \longrightarrow \underline{U}(\theta)\partial_\mu \phi(x) + (\partial_\mu \underline{U}(\theta)) \phi(x). \quad (12.5)$$

Introducing as many gauge fields as there are generators in the group, which are conveniently combined in the matrix valued field $\underline{W}_\mu = W_\mu^a L_a$, one defines

$$\underline{D}_\mu \phi(x) \equiv (\partial_\mu - ig \underline{W}_\mu) \phi(x), \quad (12.6)$$

and one obtains after transformation

$$\underline{D}_\mu \phi(x) \longrightarrow \underline{U}(\theta)\partial_\mu \phi(x) + (\partial_\mu \underline{U}(\theta)) \phi(x) - ig \underline{W}'_\mu \underline{U}(\theta)\phi(x).$$

Requiring that $\underline{D}_\mu \phi$ transforms as $\underline{D}_\mu \phi \rightarrow \underline{U}(\theta) \underline{D}_\mu \phi$ (or $\underline{D}_\mu \rightarrow \underline{U}(\theta) \underline{D}_\mu \underline{U}^{-1}(\theta)$) gives

$$\underline{D}_\mu \phi(x) \longrightarrow \underline{U}(\theta)\partial_\mu \phi(x) - ig \underline{U}(\theta) \underline{W}_\mu \phi(x),$$

which implies

$$\underline{W}'_\mu = \underline{U}(\theta) \underline{W}_\mu \underline{U}^{-1}(\theta) - \frac{i}{g} (\partial_\mu \underline{U}(\theta)) \underline{U}^{-1}(\theta), \quad (12.7)$$

or infinitesimal

$$W'^a_\mu = W^a_\mu - c_{abc} \theta^b W^c_\mu + \frac{1}{g} \partial_\mu \theta^a = W^a_\mu + \frac{1}{g} D_\mu \theta^a.$$

It is necessary to introduce the free lagrangian density for the gauge fields just like the term $-(1/4)F_{\mu\nu}F^{\mu\nu}$ in QED. For abelian fields $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu = (i/g)[D_\mu, D_\nu]$ is gauge invariant. In the nonabelian case $\partial_\mu \underline{W}_\nu^a - \partial_\nu \underline{W}_\mu^a$ does not provide a gauge invariant candidate for $\underline{G}_{\mu\nu} = G_{\mu\nu}^a L_a$, as can be checked easily. Generalizing the expression in terms of the covariant derivatives, provides a gauge invariant definition for $\underline{G}_{\mu\nu}$. We have

$$\underline{G}_{\mu\nu} = \frac{i}{g} [D_\mu, D_\nu] = \partial_\mu \underline{W}_\nu - \partial_\nu \underline{W}_\mu - ig [\underline{W}_\mu, \underline{W}_\nu], \quad (12.8)$$

with for the explicit fields

$$G_{\mu\nu}^a = \partial_\mu W_\nu^a - \partial_\nu W_\mu^a + g c_{abc} W_\mu^b W_\nu^c, \quad (12.9)$$

transforming like

$$\underline{G}_{\mu\nu} \rightarrow \underline{U}(\theta) \underline{G}_{\mu\nu} \underline{U}^{-1}(\theta). \quad (12.10)$$

The gauge-invariant lagrangian density is now constructed as

$$\mathcal{L}(\phi, \partial_\mu \phi) \longrightarrow \mathcal{L}(\phi, D_\mu \phi) - \frac{1}{2} \text{Tr} \underline{G}_{\mu\nu} \underline{G}^{\mu\nu} = \mathcal{L}(\phi, D_\mu \phi) - \frac{1}{4} G_{\mu\nu}^a G^{\mu\nu a} \quad (12.11)$$

with the standard normalization $\text{Tr}(L_a L_b) = \frac{1}{2} \delta_{ab}$. Note that the gauge fields must be massless, as a mass term $\propto M_W^2 W_\mu^a W^{\mu a}$ would break gauge invariance.

QCD, an example of a nonabelian gauge theory

As an example of a nonabelian gauge theory consider *quantum chromodynamics* (QCD), the theory describing the interactions of the colored quarks. The existence of an extra degree of freedom for each species of quarks is evident for several reasons, e.g. the necessity to have an antisymmetric wave function for the Δ^{++} particle consisting of three *up* quarks (each with charge $+(2/3)e$). With the quarks belonging to the fundamental (three-dimensional) representation of $SU(3)_C$, i.e. having three components in color space

$$\psi = \begin{pmatrix} \psi_r \\ \psi_g \\ \psi_b \end{pmatrix},$$

the wave function of the baryons (such as nucleons and deltas) form a singlet under $SU(3)_C$,

$$|\text{color}\rangle = \frac{1}{\sqrt{6}} (|rgb\rangle - |grb\rangle + |gbr\rangle - |bgr\rangle + |brg\rangle - |rbg\rangle). \quad (12.12)$$

The nonabelian gauge theory that is obtained by making the 'free' quark lagrangian, for one specific species (flavor) of quarks just the Dirac lagrangian for an elementary fermion,

$$\mathcal{L} = i \bar{\psi} \not{\partial} \psi - m \bar{\psi} \psi,$$

invariant under local $SU(3)_C$ transformations has proven to be a good candidate for the microscopic theory of the strong interactions. The representation matrices for the quarks and antiquarks in the fundamental representation are given by

$$F_a = \frac{\lambda_a}{2} \quad \text{for quarks,}$$

$$F_a = -\frac{\lambda_a^*}{2} \quad \text{for antiquarks,}$$




Figure 12.1: The vectors belonging to internal space located at each point in (one-dimensional) space

which satisfy commutation relations $[F_a, F_b] = i f_{abc} F_c$ in which f_{abc} are the (completely antisymmetric) structure constants of $SU(3)$ and where the matrices λ_a are the eight Gell-Mann matrices¹. The (locally) gauge invariant lagrangian density is

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu}^a F^{\mu\nu a} + i \bar{\psi} D \psi - m \bar{\psi} \psi, \tag{12.13}$$

with

$$D_\mu \psi = \partial_\mu \psi - ig A_\mu^a F_a \psi, \\ F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g c_{abc} A_\mu^b A_\nu^c.$$

Note that the term $i \bar{\psi} D \psi = i \bar{\psi} \not{\partial} \psi + g \bar{\psi} A^a F_a \psi = i \bar{\psi} \not{\partial} \psi + j^{\mu a} A_\mu^a$ with $j^{\mu a} = g \bar{\psi} \gamma^\mu F_a \psi$ describes the interactions of the gauge bosons A_μ^a (gluons) with the color current of the quarks (this is again precisely the Noether current corresponding to color symmetry transformations). Note furthermore that the lagrangian terms for the gluons contain interaction terms corresponding to vertices with three gluons and four gluons due to the nonabelian character of the theory. For writing down the complete set of Feynman rules it is necessary to account for the gauge symmetry in the quantization procedure. This will lead (depending on the choice of gauge conditions) to the presence of ghost fields. (For more details see e.g. Ryder, chapter 7.)

A geometric picture of gauge theories

A geometric picture of gauge theories is useful for comparison with general relativity and topological considerations (such as we have seen in the Aharonov-Bohm experiment). Consider the space $\prod_x {}^x G$ (called a fibre bundle). At each space-time point x there is considered to be a copy of an internal space G (say spin or isospin). In each of these spaces a reference frame is defined. ${}^x \psi(x)$ denotes a field vector $\psi(x)$ which belongs to a representation of G , i.e. forms a vector in the internal space (see fig. 12.1). The superscript x denotes that it is expressed with respect to the frame at point x , i.e. the basis of ${}^x G$. Let fields $A_\mu^a(x)$ determine the 'parallel displacements' in the internal space, i.e. connect

¹The Gell-Mann matrices are the eight traceless hermitean matrices generating $SU(3)$ transformations,

$$\lambda_1 = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix} \quad \lambda_2 = \begin{pmatrix} & -i \\ i & \end{pmatrix} \quad \lambda_3 = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix} \\ \lambda_4 = \begin{pmatrix} & 1 \\ & 1 \end{pmatrix} \quad \lambda_5 = \begin{pmatrix} & -i \\ & i \end{pmatrix} \quad \lambda_6 = \begin{pmatrix} & & 1 \\ & 1 & \\ & & 1 \end{pmatrix} \\ \lambda_7 = \begin{pmatrix} & -i \\ i & \end{pmatrix} \quad \lambda_8 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & & \\ & 1 & \\ & & -2 \end{pmatrix}$$

the basis for xG and ${}^{x+dx}G$,

$${}^{x+dx}\psi(x) = (1 + ig dx^\mu A_\mu^a(x) T_a)^x \psi(x) \quad (12.14)$$

$$= (1 + ig dx^\mu \underline{A}_\mu(x))^x \psi(x), \quad (12.15)$$

which connects two identical vectors, but expresses them with respect to different bases.

If there is a 'true' difference in the vector $\psi(x)$ and $\psi(x + dx)$ it is denoted with the covariant derivative connecting the vectors expressed with respect to the same basis, i.e.

$${}^{x+dx}\psi(x + dx) = (1 + dx^\mu \underline{D}_\mu)^{x+dx} \psi(x), \quad (12.16)$$

which in the presence of the 'connection' \underline{A}_μ differs from the total change between ${}^x\psi(x)$ and ${}^{x+dx}\psi(x + dx)$,

$${}^{x+dx}\psi(x + dx) = (1 + dx^\mu \partial_\mu)^x \psi(x). \quad (12.17)$$

The three equations given so far immediately give

$$\underline{D}_\mu = \partial_\mu - ig \underline{A}_\mu(x). \quad (12.18)$$

We note that local gauge invariance requires that we can modify all local systems with a (local) unitary transformation $S(x)$. The relation in Eq. 12.16, should be independent of such transformations, requiring that the 'connection' $\underline{A}_\mu(x)$ is such that $\underline{D}_\mu \rightarrow S(x)\underline{D}_\mu S^{-1}(x)$.

A 'constant' vector that only rotates because of the arbitrary definitions of local frames satisfies $\underline{D}_\mu \psi(x) = 0$, i.e.

$$[\partial_\mu - ig \underline{A}_\mu(x)] \psi(x) = 0$$

or considering a path $x^\mu(s)$ from a fixed origin (0) to point x ,

$$\frac{dx^\mu}{ds} [\partial_\mu - ig \underline{A}_\mu(x(s))] \psi(x) = 0,$$

which is solved by


$$\begin{aligned} \frac{d\psi(s)}{ds} &= ig \underline{A}_\mu(s) \psi(s) \frac{dx^\mu}{ds}, \\ \psi(s) &= \mathcal{P} \exp \left(ig \int_0^s ds' \frac{dx^\mu}{ds'} \underline{A}_\mu(s') \right) \psi(0), \end{aligned}$$

which is the path-ordered integral denoted

$$\psi(x) = \mathcal{P} e^{ig \int_P dx^\mu \underline{A}_\mu(x)} \psi(0). \quad (12.19)$$

This gives rise to a (path dependent) phase in each point.

In principle such a phase in a given point is not observable. However, if two different paths to the same point give different phases the effects can be observed. What is this physical effect by which the 'connection' \underline{A}_μ can be observed? For this consider the phase around a closed loop,



For a vector, the real change thus is given by

$$\begin{aligned} \psi(x) &= (1 - dy^\rho \underline{D}_\rho)(1 - dx^\sigma \underline{D}_\sigma)(1 + dy^\nu \underline{D}_\nu)(1 + dx^\mu \underline{D}_\mu) \psi(x) \\ &= (1 + dx^\mu dy^\nu [\underline{D}_\mu, \underline{D}_\nu]) \psi(x) \\ &= (1 - ig d\sigma^{\mu\nu} \underline{G}_{\mu\nu}) \psi(x), \end{aligned}$$

where $\underline{G}_{\mu\nu} = (i/g)[\underline{D}_\mu, \underline{D}_\nu]$. Similarly as the definition of the covariant derivative the effect is thus frame-independent and we have the transformation law $\underline{G}_{\mu\nu} \rightarrow S(x)\underline{G}_{\mu\nu}S^{-1}(x)$. In geometric language the effect on parallel transport of a vector depends on the 'curvature' $\underline{G}_{\mu\nu}$. Only if this quantity is nonzero a physically observable effect of \underline{A}_μ exists. If it is zero one has $\oint dx^\mu \underline{A}_\mu(x) = 0$, and equivalently \underline{A}_μ can be considered as a pure gauge effect, which means that by an appropriate transformation $S(x)$ it can be gauged away (see the example of Aharonov-Bohm effect).

12.2 Spontaneous symmetry breaking

In this section we consider the situation that the groundstate of a physical system is degenerate. Consider as an example a ferromagnet with an interaction hamiltonian of the form

$$H = - \sum_{i>j} J_{ij} \mathbf{S}_i \cdot \mathbf{S}_j,$$

which is rotationally invariant. If the temperature is high enough the spins are oriented randomly and the (macroscopic) ground state is spherically symmetric. If the temperature is below a certain critical temperature ($T < T_c$) the kinetic energy is no longer dominant and the above hamiltonian prefers a lowest energy configuration in which all spins are parallel. In this case there are many possible groundstates (determined by a fixed direction in space). This characterizes spontaneous symmetry breaking, the groundstate itself appears degenerate. As there can be one and only one groundstate, this means that there is more than one possibility for the groundstate. Nature will choose one, usually being (slightly) prejudiced by impurities, external magnetic fields, i.e. in reality a not perfectly symmetric situation.

Nevertheless, we can disregard those 'perturbations' and look at the ideal situation, e.g. a theory for a scalar degree of freedom (a scalar field) having three (real) components,

$$\vec{\phi} = \begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{pmatrix},$$

with a lagrangian density of the form

$$\mathcal{L} = \frac{1}{2} \partial_\mu \vec{\phi} \cdot \partial^\mu \vec{\phi} - \underbrace{\frac{1}{2} m^2 \vec{\phi} \cdot \vec{\phi} - \frac{1}{4} \lambda (\vec{\phi} \cdot \vec{\phi})^2}_{-V(\vec{\phi})}. \quad (12.20)$$

The potential $V(\vec{\phi})$ is shown in fig. 12.2. Classically the (time-independent) ground state is found for




Figure 12.2: The symmetry-breaking 'potential' in the lagrangian for the case that $m^2 < 0$.

a constant field ($\nabla\vec{\phi} = 0$) and the condition

$$\left. \frac{\partial V}{\partial \vec{\phi}} \right|_{\vec{\varphi}_c} = 0 \quad \longrightarrow \quad \vec{\varphi}_c \cdot \vec{\varphi}_c = 0 \quad \text{or} \quad \vec{\varphi}_c \cdot \vec{\varphi}_c = -\frac{m^2}{\lambda} \equiv F^2,$$

the latter only forming a minimum for $m^2 < 0$. In this situation one speaks of *spontaneous symmetry breaking*. The classical groundstate appears degenerate. Any constant field with 'length' F is a possible groundstate. The presence of a nonzero value for the classical groundstate value of the field will have an effect when the field is quantized. A quantum field theory has only *one* nondegenerate groundstate $|0\rangle$. Writing the field $\vec{\phi}$ as a sum of a classical and a quantum field, $\vec{\phi} = \vec{\varphi}_c + \vec{\phi}_{\text{quantum}}$ where for the (operator-valued) coefficients in the quantum field one wants $\langle 0|c^\dagger = c|0\rangle = 0$, so one has

$$\langle 0|\text{vecphi}_{\text{quantum}}|0\rangle = 0 \quad \text{and hence} \quad \langle 0|\vec{\phi}|0\rangle = \vec{\varphi}_c. \quad (12.21)$$

Stability of the action requires the classical groundstate $\vec{\varphi}_c$ to have a well-defined value (which can be nonzero), while the quadratic terms must correspond with non-negative masses. In the case of degeneracy, therefore a choice must be made, say

$$\langle 0|\vec{\phi}|0\rangle = \begin{pmatrix} 0 \\ 0 \\ F \end{pmatrix}. \quad (12.22)$$

The situation now is the following. The original lagrangian contained an $\text{SO}(3)$ invariance under (length conserving) rotations among the three fields, while the lagrangian including the nonzero groundstate expectation value chosen by nature, has less symmetry. It is only invariant under rotations around the 3-axis.

It is appropriate to redefine the field as

$$\vec{\phi} = \begin{pmatrix} \varphi_1 \\ \varphi_2 \\ F + \eta \end{pmatrix}, \quad (12.23)$$

such that $\langle 0|\varphi_1|0\rangle = \langle 0|\varphi_2|0\rangle = \langle 0|\eta|0\rangle = 0$. The field along the third axis plays a special role because of the choice of the vacuum expectation value. In order to see the consequences for the particle spectrum of the theory we construct the lagrangian in terms of the fields φ_1 , φ_2 and η . It is sufficient to do this to second order in the fields as the higher (cubic, etc.) terms constitute interaction terms. The result is

$$\begin{aligned} \mathcal{L} &= \frac{1}{2} (\partial_\mu \varphi_1)^2 + \frac{1}{2} (\partial_\mu \varphi_2)^2 + \frac{1}{2} (\partial_\mu \eta)^2 - \frac{1}{2} m^2 (\varphi_1^2 + \varphi_2^2) \\ &\quad - \frac{1}{2} m^2 (F + \eta)^2 - \frac{1}{4} \lambda (\varphi_1^2 + \varphi_2^2 + F^2 + \eta^2 + 2F\eta)^2 \end{aligned} \quad (12.24)$$

$$= \frac{1}{2} (\partial_\mu \varphi_1)^2 + \frac{1}{2} (\partial_\mu \varphi_2)^2 + \frac{1}{2} (\partial_\mu \eta)^2 + m^2 \eta^2 + \dots \quad (12.25)$$

Therefore there are *2 massless scalar particles*, corresponding to the number of broken generators (in this case rotations around 1 and 2 axis) and *1 massive scalar particle* with mass $m_\eta^2 = -2m^2$. The massless particles are called *Goldstone bosons*.

Realization of symmetries

In this section we want to discuss a bit more formal the two possible ways that a symmetry can be implemented. They are known as the *Weyl mode* or the *Goldstone mode*:

Weyl mode. In this mode the lagrangian *and* the vacuum are both invariant under a set of symmetry transformations generated by Q^a , i.e. for the vacuum $Q^a|0\rangle = 0$. In this case the spectrum is described

by degenerate representations of the symmetry group. Known examples are rotational symmetry and the fact that the spectrum shows multiplets labeled by angular momentum ℓ (with members labeled by m). The generators Q^a (in that case the rotation operators L_z, L_x and L_y or instead of the latter two L_+ and L_-) are used to label the multiplet members or transform them into one another. A bit more formal, if the generators Q^a generate a symmetry, i.e. $[Q^a, H] = 0$, and $|a\rangle$ and $|a'\rangle$ belong to the same multiplet (there is a Q^a such that $|a'\rangle = Q^a|a\rangle$) then $H|a\rangle = E_a|a\rangle$ implies that $H|a'\rangle = E_a|a'\rangle$, i.e. a and a' are degenerate states.

Goldstone mode. In this mode the lagrangian is invariant but $Q^a|0\rangle \neq 0$ for a number of generators. This means that they are operators that create states from the vacuum, denoted $|\pi^a(k)\rangle$. As the generators for a symmetry are precisely the zero-components of a conserved current $J_\mu^a(x)$ integrated over space, there must be a nonzero expectation value $\langle 0|J_\mu^a(x)|\pi^a(k)\rangle$. Using translation invariance and as k_μ is the only four vector on which this matrix element could depend one may write

$$\langle 0|J_\mu^a(x)|\pi^b(k)\rangle = f_\pi k_\mu e^{ik \cdot x} \delta_{ab} \quad (f_\pi \neq 0) \quad (12.26)$$

for all the states labeled by a corresponding to 'broken' generators. Taking the derivative,

$$\langle 0|\partial^\mu J_\mu^a(x)|\pi^b(k)\rangle = f_\pi k^2 e^{ik \cdot x} \delta_{ab} = f_\pi m_{\pi^a}^2 e^{ik \cdot x} \delta_{ab}. \quad (12.27)$$

If the transformations in the lagrangian give rise to a symmetry the Noether currents are conserved, $\partial^\mu J_\mu^a = 0$, irrespective of the fact if they annihilate the vacuum, and one must have $m_{\pi^a} = 0$, i.e. a massless Goldstone boson for each 'broken' generator. Note that for the fields $\pi^a(x)$ one would have the relation $\langle 0|\pi^a(x)|\pi^a(k)\rangle = e^{ik \cdot x}$, suggesting the stronger relation $\partial^\mu J_\mu^a(x) = f_\pi m_{\pi^a}^2 \pi^a(x)$.

Chiral symmetry

An example of spontaneous symmetry breaking is chiral symmetry breaking in QCD. Neglecting at this point the local color symmetry, the lagrangian for the quarks consists of the free Dirac lagrangian for each of the types of quarks, called flavors. Including a sum over the different flavors (up, down, strange, etc.) one can write

$$\mathcal{L} = \bar{\psi}(i\cancel{\partial} - M)\psi, \quad (12.28)$$

where ψ is extended to a vector in *flavor space* and M is a diagonal matrix,

$$\psi = \begin{pmatrix} \psi_u \\ \psi_d \\ \vdots \end{pmatrix}, \quad M = \begin{pmatrix} m_u & & \\ & m_d & \\ & & \ddots \end{pmatrix} \quad (12.29)$$

(Note that each of the entries in the vector for ψ is a 4-component Dirac spinor). This lagrangian density then is invariant under unitary (vector) transformations in the flavor space,

$$\psi \longrightarrow e^{i\vec{\alpha} \cdot \vec{T}} \psi, \quad (12.30)$$

which for instance including only two flavors form an $SU(2)_V$ symmetry (isospin symmetry) generated by the Pauli matrices, $\vec{T} = \vec{\tau}/2$. The conserved currents corresponding to this symmetry transformation are found directly using Noether's theorem (see chapter 6),

$$\vec{V}^\mu = \bar{\psi} \gamma^\mu \vec{T} \psi. \quad (12.31)$$

Using the Dirac equation, it is easy to see that one gets

$$\partial_\mu \vec{V}^\mu = i \bar{\psi} [M, \vec{T}] \psi. \quad (12.32)$$

Furthermore $\partial_\mu \vec{V}^\mu = 0 \iff [M, \vec{T}] = 0$. From group theory (Schur's theorem) one knows that the latter can only be true, if in flavor space M is proportional to the unit matrix, $M = m \cdot 1$. I.e. $SU(2)_V$ (isospin) symmetry is good if the up and down quark masses are identical. This situation, both are very small, is what happens in the real world. This symmetry is realized in the Weyl mode with the spectrum of QCD showing an almost perfect isospin symmetry, e.g. a doublet (isospin 1/2) of nucleons, proton and neutron, with almost degenerate masses ($M_p = 938.3 \text{ MeV}/c^2$ and $M_n = 939.6 \text{ MeV}/c^2$), but also a triplet (isospin 1) of pions, etc.

There exists another set of symmetry transformations, so-called axial vector transformations,

$$\psi \longrightarrow e^{i \vec{\alpha} \cdot \vec{T} \gamma_5} \psi, \quad (12.33)$$

which for instance including only two flavors form $SU(2)_A$ transformations generated by the Pauli matrices, $\vec{T} \gamma_5 = \vec{\tau} \gamma_5 / 2$. Note that these transformations also work on the spinor indices. The currents corresponding to this symmetry transformation are again found using Noether's theorem,

$$\vec{A}^\mu = \bar{\psi} \gamma^\mu \vec{T} \gamma_5 \psi. \quad (12.34)$$

Using the Dirac equation, it is easy to see that one gets

$$\partial_\mu \vec{A}^\mu = i \bar{\psi} \{M, \vec{T}\} \gamma_5 \psi. \quad (12.35)$$

In this case $\partial_\mu \vec{A}^\mu = 0$ will be true if the quarks have zero mass, which is approximately true for the up and down quarks. Therefore the world of up and down quarks describing pions, nucleons and atomic nuclei has not only an isospin or vector symmetry $SU(2)_V$ but also an axial vector symmetry $SU(2)_A$. This combined symmetry is what one calls *chiral symmetry*.

That the massless theory has this symmetry can also be seen by writing it down for the so-called lefthanded and righthanded fermions, $\psi_{R/L} = \frac{1}{2}(1 \pm \gamma_5)\psi$, in terms of which the Dirac lagrangian density looks like

$$\mathcal{L} = i \bar{\psi}_L \not{\partial} \psi_L + i \bar{\psi}_R \not{\partial} \psi_R - \bar{\psi}_R M \psi_L - \bar{\psi}_L M \psi_R. \quad (12.36)$$

If the mass is zero the lagrangian is split into two disjunct parts for L and R showing that there is a direct product $SU(2)_L \otimes SU(2)_R$ symmetry, generated by $\vec{T}_{R/L} = \frac{1}{2}(1 \pm \gamma_5)\vec{T}$, which is equivalent to the V-A symmetry. This symmetry, however, is by nature not realized in the Weyl mode. How can we see this. The chiral fields ψ_R and ψ_L are transformed into each other under parity. Therefore realization in the Weyl mode would require that all particles come double with positive and negative parity, or, stated equivalently, parity would not play a role in the world. We know that mesons and baryons (such as the nucleons) have a well-defined parity that is conserved.

The conclusion is that the original symmetry of the lagrangian is spontaneously broken and as the vector part of the symmetry is the well-known isospin symmetry, nature has chosen the path

$$SU(2)_L \otimes SU(2)_R \implies SU(2)_V,$$

i.e. the lagrangian density is invariant under left (L) and right (R) rotations independently, while the groundstate is only invariant under isospin rotations ($R = L$). From the number of broken generators it is clear that one expects three massless Goldstone bosons, for which the field (according to the discussion above) has the same behavior under parity, etc. as the quantity $\partial_\mu A^\mu(x)$, i.e. (leaving out the flavor structure) the same as $\bar{\psi} \gamma_5 \psi$, i.e. behaves as a pseudoscalar particle (spin zero, parity minus). In the real world, where the quark masses are not completely zero, chiral symmetry is not perfect. Still the basic fact that the generators acting on the vacuum give a nonzero result (i.e. $f_\pi \neq 0$ remains, but the fact that the symmetry is not perfect and the right hand side of Eq. 12.35 is nonzero, gives also rise to a nonzero mass for the Goldstone bosons according to Eq. 12.27. The Goldstone bosons of QCD are the pions for which $f_\pi = 93 \text{ MeV}$ and which have a mass of $m_\pi \approx 138 \text{ MeV}/c^2$, much smaller than any of the other mesons or baryons.

12.3 The Higgs mechanism

The Higgs mechanism occurs when spontaneous symmetry breaking happens in a gauge theory where gauge bosons have been introduced in order to assure the local symmetry. Considering the same example with rotational symmetry ($SO(3)$) as for spontaneous symmetry breaking of a scalar field (Higgs field) with three components, made into a gauge theory,

$$\mathcal{L} = -\frac{1}{4} \vec{G}_{\mu\nu} \cdot \vec{G}^{\mu\nu} + \frac{1}{2} D_\mu \vec{\phi} \cdot D^\mu \vec{\phi} - V(\vec{\phi}), \quad (12.37)$$

where

$$D_\mu \vec{\phi} = \partial_\mu \vec{\phi} - ig W_\mu^a L_a \vec{\phi}. \quad (12.38)$$

Since the explicit (adjoint, in this case three-dimensional) representation reads $(L_a)_{ij} = -i \epsilon_{aij}$, one sees that the fields \vec{W}_μ and $\vec{G}_{\mu\nu}$ also can be represented as three-component fields,

$$D_\mu \vec{\phi} = \partial_\mu \vec{\phi} + g \vec{W}_\mu \times \vec{\phi}, \quad (12.39)$$

$$\vec{G}_{\mu\nu} = \partial_\mu \vec{W}_\nu - \partial_\nu \vec{W}_\mu + g \vec{W}_\mu \times \vec{W}_\nu. \quad (12.40)$$

The symmetry is broken in the same way as before and the same choice for the vacuum,

$$\vec{\varphi}_c = \langle 0 | \vec{\phi} | 0 \rangle = \begin{pmatrix} 0 \\ 0 \\ F \end{pmatrix}.$$

is made. The difference comes when we reparametrize the field $\vec{\phi}$. We have the possibility to perform local gauge transformations. Therefore we can always rotate the field ϕ into the 3-direction in order to simplify the calculation, i.e.

$$\vec{\phi} = \begin{pmatrix} 0 \\ 0 \\ \phi_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ F + \eta \end{pmatrix}. \quad (12.41)$$

Explicitly one then has

$$D_\mu \vec{\phi} = \partial_\mu \vec{\phi} + g \vec{W}_\mu \times \vec{\phi} = \begin{pmatrix} gF W_\mu^2 + g W_\mu^2 \eta \\ -gF W_\mu^1 - g W_\mu^1 \eta \\ \partial_\mu \eta \end{pmatrix},$$

which gives for the lagrangian density up to quadratic terms

$$\begin{aligned} \mathcal{L} &= -\frac{1}{4} \vec{G}_{\mu\nu} \cdot \vec{G}^{\mu\nu} + \frac{1}{2} D_\mu \vec{\phi} \cdot D^\mu \vec{\phi} - \frac{1}{2} m^2 \vec{\phi} \cdot \vec{\phi} - \frac{\lambda}{4} (\vec{\phi} \cdot \vec{\phi})^2 \\ &= -\frac{1}{4} (\partial_\mu \vec{W}_\nu - \partial_\nu \vec{W}_\mu) \cdot (\partial^\mu \vec{W}^\nu - \partial^\nu \vec{W}^\mu) - \frac{1}{2} g^2 F^2 (W_\mu^1 W^{\mu 1} + W_\mu^2 W^{\mu 2}) \\ &\quad + \frac{1}{2} (\partial_\mu \eta)^2 + m^2 \eta^2 + \dots, \end{aligned} \quad (12.42)$$

from which one reads off that the particle content of the theory consists of one massless gauge boson (W_μ^3), two massive bosons (W_μ^1 and W_μ^2 with $M_W = gF$) and a massive scalar particle (η with $m_\eta^2 = -2m^2$). The latter is a spin 0 particle (real scalar field) called a Higgs particle. Note that the number of massless gauge bosons (in this case one) coincides with the number of generators corresponding to the remaining symmetry (in this case rotations around the 3-axis), while the number of massive gauge bosons coincides with the number of 'broken' generators.

One may wonder about the degrees of freedom, as in this case there are no massless Goldstone bosons. Initially there are 3 massless gauge fields (each, like a photon, having two independent spin components) and three scalar fields (one degree of freedom each), thus 9 independent degrees of freedom. After symmetry breaking the same number (as expected) comes out, but one has 1 massless gauge field (2), 2 massive vector fields or spin 1 bosons (2×3) and one scalar field (1), again 9 degrees of freedom.

12.4 The standard model $SU(2)_W \otimes U(1)_Y$

The symmetry ideas discussed before play an essential role in the standard model that describes the elementary particles, the quarks (up, down, etc.), the leptons (elektrons, muons, neutrinos, etc.) and the gauge bosons responsible for the strong, electromagnetic and weak forces. In the standard model one starts with a very simple basic lagrangian for (massless) fermions which exhibits more symmetry than observed in nature. By introducing gauge fields and breaking the symmetry a more complex lagrangian is obtained, that gives a good description of the physical world. The procedure, however, implies certain nontrivial relations between masses and mixing angles that can be tested experimentally and so far are in excellent agreement with experiment.

The lagrangian for the leptons consists of three families each containing an elementary fermion (electron e^- , muon μ^- or tau τ^-), its corresponding neutrino (ν_e , ν_μ and ν_τ) and their antiparticles. As they are massless, left- and righthanded particles, $\psi_{R/L} = \frac{1}{2}(1 \pm \gamma_5)\psi$ decouple. For the neutrino only a lefthanded particle (and righthanded antiparticle) exist. Thus

$$\mathcal{L}^{(f)} = i\bar{e}_R\partial e_R + i\bar{e}_L\partial e_L + i\bar{\nu}_{eL}\partial\nu_{eL} + (\mu, \tau). \quad (12.43)$$

One introduces a (weak) $SU(2)_W$ symmetry under which e_R forms a singlet, while the lefthanded particles form a doublet, i.e.

$$L = \begin{pmatrix} \nu_e \\ e_L \end{pmatrix} \quad \text{with } T_W = \frac{1}{2} \text{ and } T_W^3 = \begin{cases} +1/2 \\ -1/2 \end{cases}$$

and

$$R = e_R \quad \text{with } T_W = 0 \text{ and } T_W^3 = 0.$$

Thus the lagrangian density is

$$\mathcal{L}^{(f)} = i\bar{L}\partial L + i\bar{R}\partial R, \quad (12.44)$$

which has an $SU(2)_W$ symmetry under transformations $e^{i\vec{\alpha}\cdot\vec{T}_W}$, explicitly

$$L \xrightarrow{SU(2)_W} e^{i\vec{\alpha}\cdot\vec{\tau}/2} L, \quad (12.45)$$

$$R \xrightarrow{SU(2)_W} R. \quad (12.46)$$

One notes that the charges of the leptons can be obtained as $Q = T_W^3 - 1/2$ for lefthanded particles and $Q = T_W^3 - 1$ for righthanded particles. This is written as

$$Q = T_W^3 + \frac{Y_W}{2}, \quad (12.47)$$

and Y_W is considered as an operator that generates a $U(1)_Y$ symmetry, under which the lefthanded and righthanded particles with $Y_W(L) = -1$ and $Y_W(R) = -2$ transform with $e^{i\beta Y_W/2}$, explicitly

$$L \xrightarrow{U(1)_Y} e^{-i\beta/2} L, \quad (12.48)$$

$$R \xrightarrow{U(1)_Y} e^{-i\beta} R. \quad (12.49)$$

Next the $SU(2)_W \otimes U(1)_Y$ symmetry is made into a local symmetry introducing gauge fields \vec{W}_μ and B_μ in the covariant derivative $D_\mu = \partial_\mu + i g \vec{W}_\mu \cdot \vec{T}_W + i g' B_\mu Y_W/2$, explicitly

$$D_\mu L = \partial_\mu L - \frac{i}{2} g \vec{W}_\mu \cdot \vec{\tau} L + \frac{i}{2} g' B_\mu L, \quad (12.50)$$

$$D_\mu R = \partial_\mu R + i g' B_\mu R, \quad (12.51)$$

where \vec{W}_μ is a triplet of gauge bosons with $T_W = 1$, $T_W^3 = \pm 1$ or 0 and $Y_W = 0$ (thus $Q = T_W^3$) and B_μ is a singlet under $SU(2)_W$ ($T_W = T_W^3 = 0$) and also has $Y_W = 0$. Putting this in leads to

$$\mathcal{L}^{(f)} = \mathcal{L}^{(f1)} + \mathcal{L}^{(f2)}, \quad (12.52)$$

$$\begin{aligned} \mathcal{L}^{(f1)} &= i\bar{R}\gamma^\mu(\partial_\mu + ig'B_\mu)R + i\bar{L}\gamma^\mu(\partial_\mu + \frac{i}{2}g'B_\mu - \frac{i}{2}g\vec{W}_\mu \cdot \vec{\tau})L \\ \mathcal{L}^{(f2)} &= -\frac{1}{4}(\partial_\mu\vec{W}_\nu - \partial_\nu\vec{W}_\mu + g\vec{W}_\mu \times \vec{W}_\nu)^2 - \frac{1}{4}(\partial_\mu B_\nu - \partial_\nu B_\mu)^2. \end{aligned}$$

In order to break the symmetry to the symmetry of the physical world, the $U(1)_Q$ symmetry (generated by the charge operator), a complex Higgs field

$$\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{2}}(\theta_2 + i\theta_1) \\ \frac{1}{\sqrt{2}}(\theta_4 - i\theta_3) \end{pmatrix} \quad (12.53)$$

with $T_W = 1/2$ and $Y_W = 1$ is introduced, with the following lagrangian density consisting of a symmetry breaking piece and a coupling to the fermions,

$$\mathcal{L}^{(h)} = \mathcal{L}^{(h1)} + \mathcal{L}^{(h2)}, \quad (12.54)$$

where

$$\begin{aligned} \mathcal{L}^{(h1)} &= (D_\mu\phi)^\dagger(D^\mu\phi) \underbrace{-m^2\phi^\dagger\phi - \lambda(\phi^\dagger\phi)^2}_{-V(\phi)}, \\ \mathcal{L}^{(h2)} &= -G_e(\bar{L}\phi R + \bar{R}\phi^\dagger L), \end{aligned}$$

and

$$D_\mu\phi = (\partial_\mu - \frac{i}{2}g\vec{W}_\mu \cdot \vec{\tau} - \frac{i}{2}g'B_\mu)\phi. \quad (12.55)$$

The Higgs potential $V(\phi)$ is chosen such that it gives rise to spontaneous symmetry breaking with $\phi^\dagger\phi = -m^2/2\lambda \equiv v^2/2$. For the classical field the choice $\theta_4 = v$ is made, which assures with the choice of Y_W of the Higgs field assures that Q generates the remaining $U(1)$ symmetry. Using *local* gauge invariance θ_i for $i = 1, 2$ and 3 may be eliminated (the necessary $SU(2)_W$ rotation is precisely $e^{-i\vec{\theta}(x)\cdot\vec{\tau}}$), leading to the parametrization

$$\phi(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + h(x) \end{pmatrix} \quad (12.56)$$

and

$$D_\mu\phi = \begin{pmatrix} \frac{ig}{2} \left(\frac{W_\mu^1 - iW_\mu^2}{\sqrt{2}} \right) (v + h) \\ \frac{1}{\sqrt{2}} \partial_\mu h - \frac{i}{2} \left(\frac{gW_\mu^3 - g'B_\mu}{\sqrt{2}} \right) (v + h) \end{pmatrix}. \quad (12.57)$$

Up to cubic terms, this leads to the lagrangian

$$\begin{aligned} \mathcal{L}^{(h1)} &= \frac{1}{2}(\partial_\mu h)^2 + m^2 h^2 + \frac{g^2 v^2}{8} [(W_\mu^1)^2 + (W_\mu^2)^2] \\ &\quad + \frac{v^2}{8} (gW_\mu^3 - g'B_\mu)^2 + \dots \end{aligned} \quad (12.58)$$

$$\begin{aligned} &= \frac{1}{2}(\partial_\mu h)^2 + m^2 h^2 + \frac{g^2 v^2}{8} [(W_\mu^+)^2 + (W_\mu^-)^2] \\ &\quad + \frac{(g^2 + g'^2) v^2}{8} (Z_\mu)^2 + \dots, \end{aligned} \quad (12.59)$$

where the quadratically appearing gauge fields that are furthermore eigenstates of the charge operator are

$$W_\mu^\pm = \frac{1}{\sqrt{2}} (W_\mu^1 \pm i W_\mu^2), \quad (12.60)$$

$$Z_\mu = \frac{g W_\mu^3 - g' B_\mu}{\sqrt{g^2 + g'^2}} \equiv \cos \theta_W W_\mu^3 - \sin \theta_W B_\mu, \quad (12.61)$$

$$A_\mu = \frac{g' W_\mu^3 + g B_\mu}{\sqrt{g^2 + g'^2}} \equiv \sin \theta_W W_\mu^3 + \cos \theta_W B_\mu, \quad (12.62)$$

and correspond to three massive particle fields (W^\pm and Z^0) and one massless field (photon γ) with

$$M_W^2 = \frac{g^2 v^2}{4}, \quad (12.63)$$

$$M_Z^2 = \frac{g^2 v^2}{4 \cos^2 \theta_W} = \frac{M_W^2}{\cos^2 \theta_W}, \quad (12.64)$$

$$M_\gamma^2 = 0. \quad (12.65)$$

The weak mixing angle is related to the ratio of coupling constants, $g'/g = \tan \theta_W$.


The coupling of the fermions to the physical gauge bosons are contained in $\mathcal{L}^{(f1)}$ giving

$$\begin{aligned} \mathcal{L}^{(f1)} &= i \bar{e} \gamma^\mu \partial_\mu e + i \bar{\nu}_e \gamma^\mu \partial_\mu \nu_e - g \sin \theta_W \bar{e} \gamma^\mu e A_\mu \\ &+ \frac{g}{\cos \theta_W} \left(\sin^2 \theta_W \bar{e}_R \gamma^\mu e_R - \frac{1}{2} \cos 2\theta_W \bar{e}_L \gamma^\mu e_L + \frac{1}{2} \bar{\nu}_e \gamma^\mu \nu_e \right) Z_\mu \\ &+ \frac{g}{\sqrt{2}} (\bar{\nu}_e \gamma^\mu e_L W_\mu^- + \bar{e}_L \gamma^\mu \nu_e W_\mu^+). \end{aligned} \quad (12.66)$$

From the coupling to the photon, we can read off

$$e = g \sin \theta_W = g' \cos \theta_W. \quad (12.67)$$

The coupling of electrons or muons to their respective neutrinos, for instance in the amplitude for the decay of the muon



is given by

$$\begin{aligned} -i \mathcal{M} &= -\frac{g^2}{2} (\bar{\nu}_\mu \gamma^\rho \mu_L) \frac{-i g_{\rho\sigma} + \dots}{k^2 + M_W^2} (\bar{e}_L \gamma^\sigma \nu_e) \\ &\approx i \frac{g^2}{8 M_W^2} \underbrace{(\bar{\nu}_\mu \gamma_\rho (1 - \gamma_5) \mu)}_{(j_L^{(\mu)\dagger})_\rho} \underbrace{(\bar{e} \gamma^\rho (1 - \gamma_5) \nu_e)}_{(j_L^{(e)})_\rho} \end{aligned} \quad (12.68)$$

$$\equiv i \frac{G_F}{\sqrt{2}} (j_L^{(\mu)\dagger})_\rho (j_L^{(e)})^\rho, \quad (12.69)$$

the good old four-point interaction introduced by Fermi to explain the weak interactions, i.e. one has the relation

$$\frac{G_F}{\sqrt{2}} = \frac{g^2}{8 M_W^2} = \frac{e^2}{8 M_W^2 \sin^2 \theta_W} = \frac{1}{2 v^2}. \quad (12.70)$$

In this way the parameters g , g' and v determine a number of experimentally measurable quantities, such as

$$e^2/4\pi \approx 1/137, \quad (12.71)$$

$$G_F = 1.1664 \times 10^{-5} \text{ GeV}^{-2}, \quad (12.72)$$

$$\sin^2 \theta_W = 0.2312, \quad (12.73)$$

$$M_W = 80.40 \text{ GeV}, \quad (12.74)$$

$$M_Z = 91.19 \text{ GeV}. \quad (12.75)$$

The coupling of the Z^0 to fermions is given by $(g/\cos\theta_W)\gamma^\mu$ multiplied with

$$T_W^3 \frac{1}{2}(1 - \gamma_5) - \sin^2 \theta_W Q \equiv \frac{1}{2} C_V - \frac{1}{2} C_A \gamma_5, \quad (12.76)$$

with

$$C_V = T_W^3 - 2 \sin^2 \theta_W Q, \quad (12.77)$$

$$C_A = T_W^3. \quad (12.78)$$

From this coupling it is straightforward to calculate the partial width for Z^0 into a fermion-antifermion pair,

$$\Gamma(Z^0 \rightarrow f\bar{f}) = \frac{M_Z}{48\pi} \frac{g^2}{\cos^2 \theta_W} (C_V^2 + C_A^2). \quad (12.79)$$


For the electron, muon or tau, leptons with $C_V = -1/2 + 2 \sin^2 \theta_W \approx -0.05$ and $C_A = -1/2$ we calculate $\Gamma(e^+e^-) \approx 78.5 \text{ MeV}$ (exp. $\Gamma_e \approx \Gamma_\mu \approx \Gamma_\tau \approx 83 \text{ MeV}$). For each neutrino species (with $C_V = 1/2$ and $C_A = 1/2$ one expects $\Gamma(\bar{\nu}\nu) \approx 155 \text{ MeV}$. Comparing this with the total width into (invisible!) channels, $\Gamma_{invisible} = 480 \text{ MeV}$ one sees that three families of (light) neutrinos are allowed. Actually including corrections corresponding to higher order diagrams the agreement for the decay width into electrons can be calculated much more accurately and the number of allowed (light) neutrinos turns to be even closer to three.

The masses of the fermions and the coupling to the Higgs particle are contained in $\mathcal{L}^{(h2)}$. With the chosen vacuum expectation value for the Higgs field, one obtains

$$\begin{aligned} \mathcal{L}^{(h2)} &= -\frac{G_e v}{\sqrt{2}} (\bar{e}_L e_R + \bar{e}_R e_L) - \frac{G_e}{\sqrt{2}} (\bar{e}_L e_R + \bar{e}_R e_L) h \\ &= -m_e \bar{e}e - \frac{m_e}{v} \bar{e}eh. \end{aligned} \quad (12.80)$$

First, the mass of the electron comes from the spontaneous symmetry breaking but is not predicted (it is in the coupling G_e). The coupling to the Higgs particle is weak as the value for v calculated e.g. from the M_W mass is about 250 GeV, i.e. m_e/v is extremely small.

Finally we want to say something about the weak properties of the quarks, as appear for instance in the decay of the neutron or the decay of the Λ (quark content uds),






Figure 12.3: Appropriate Y_W and T_W^3 assignments of quarks, leptons, their antiparticles and the electroweak gauge bosons as appearing in each family. The electric charge Q is then fixed, $Q = T_W^3 + Y_W/2$ and constant along specific diagonals as indicated in the figure. The pattern is actually intriguing, suggesting an underlying larger unifying symmetry group, for which $SU(5)$ or $SO(10)$ are actually nice candidates. We will not discuss this any further in this chapter.

The quarks also turn out to fit into doublets of $SU(2)_W$ for the lefthanded species and into singlets for the righthanded quarks. As shown in Fig. 12.3, this requires particular Y_W - T_W^3 assignments to get the charges right.

A complication arises for quarks (and as we will discuss in the next section in more detail also for leptons) as it are not the 'mass' eigenstates that appear in the weak isospin doublets but linear combinations of them,

$$\begin{pmatrix} u \\ d' \end{pmatrix}_L \quad \begin{pmatrix} c \\ s' \end{pmatrix}_L \quad \begin{pmatrix} t \\ b' \end{pmatrix}_L,$$

where

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix}_L = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}_L \tag{12.81}$$

This mixing allows all quarks with $T_W^3 = -1/2$ to decay into an up quark, but with different strength. Comparing neutron decay and Λ decay one can get an estimate of the mixing parameter V_{us} in the so-called Cabibbo-Kobayashi-Maskawa mixing matrix. Decay of B-mesons containing b-quarks allow estimate of V_{ub} , etc. In principle one complex phase is allowed in the most general form of the CKM matrix, which can account for the (observed) CP violation of the weak interactions. This is only true if the mixing matrix is at least three-dimensional, i.e. CP violation requires three generations. The magnitudes of the entries in the CKM matrix are nicely represented using the so-called Wolfenstein parametrization

$$V = \begin{pmatrix} 1 - \frac{1}{2}\lambda^2 & \lambda & \lambda^3 A(\rho - i\eta) \\ -\lambda & 1 - \frac{1}{2}\lambda^2 & \lambda^2 A \\ \lambda^3 A(1 - \rho - i\eta) & -\lambda^2 A & 1 \end{pmatrix} + \mathcal{O}(\lambda^4)$$

with $\lambda \approx 0.227$, $A \approx 0.82$ and $\rho \approx 0.22$ and $\eta \approx 0.34$. The imaginary part $i\eta$ gives rise to CP violation in decays of \bar{K} and B-mesons (containing s and b quarks, respectively).

12.5 Family mixing in the Higgs sector and neutrino masses

The quark sector

Allowing for the most general (Dirac) mass generating term in the lagrangian one starts with

$$\mathcal{L}^{(h2,q)} = -\overline{Q}_L \phi \Lambda_d D_R - \overline{D}_R \Lambda_d^\dagger \phi^\dagger Q_L - \overline{Q}_L \phi^c \Lambda_u U_R - \overline{U}_R \Lambda_u^\dagger \phi^{c\dagger} Q_L \quad (12.82)$$

where we include now the three lefthanded quark doublets in Q_L , the three righthanded quarks with charge $+2/3$ in U_R and the three righthanded quarks with charges $-1/3$ in D_R , each of these containing the three families, e.g. $\overline{U}_R = \begin{pmatrix} \overline{u}_R & \overline{c}_R & \overline{t}_R \end{pmatrix}$. The Λ_u and Λ_d are complex matrices in the 3×3 family space. The Higgs field is still limited to one complex doublet. Note that we need the conjugate Higgs field to get a $U(1)_Y$ singlet in the case of the charge $+2/3$ quarks, for which we need the appropriate weak isospin doublet

$$\phi^c = \begin{pmatrix} \phi^{0*} \\ -\phi^- \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} v+h \\ 0 \end{pmatrix}.$$

For the (squared) complex matrices we can find positive eigenvalues,

$$\Lambda_u \Lambda_u^\dagger = V_u G_u^2 V_u^\dagger, \quad \text{and} \quad \Lambda_d \Lambda_d^\dagger = V_d G_d^2 V_d^\dagger, \quad (12.83)$$

where V_u and V_d are unitary matrices, allowing us to write

$$\Lambda_u = V_u G_u W_u^\dagger \quad \text{and} \quad \Lambda_d = V_d G_d W_d^\dagger, \quad (12.84)$$

with G_u and G_d being real and positive and W_u and W_d being different unitary matrices. Thus one has

$$\mathcal{L}^{(h2,q)} \implies -\overline{D}_L V_d M_d W_d^\dagger D_R - \overline{D}_R W_d M_d V_d^\dagger D_L - \overline{U}_L V_u M_u W_u^\dagger U_R - \overline{U}_R W_u M_u V_u^\dagger U_L \quad (12.85)$$

with $M_u = G_u v / \sqrt{2}$ (diagonal matrix containing m_u, m_c and m_t) and $M_d = G_d v / \sqrt{2}$ (diagonal matrix containing m_d, m_s and m_b). One then reads off that starting with the family basis as defined via the left doublets that the mass eigenstates (and states coupling to the Higgs field) involve the righthanded states $U_R^{\text{mass}} = W_u^\dagger U_R$ and $D_R^{\text{mass}} = W_d^\dagger D_R$ and the lefthanded states $U_L^{\text{mass}} = V_u^\dagger U_L$ and $D_L^{\text{mass}} = V_d^\dagger D_L$. Working with the mass eigenstates one simply sees that the weak current coupling to the W^\pm becomes $\overline{U}_L \gamma^\mu D_L = \overline{U}_L^{\text{mass}} \gamma^\mu V_u^\dagger V_d D_L^{\text{mass}}$, i.e. the weak mass eigenstates are

$$D_L' = D_L^{\text{weak}} = V_u^\dagger V_d D_L^{\text{mass}} = V_{\text{CKM}} D_L^{\text{mass}}, \quad (12.86)$$

the unitary CKM-matrix introduced above in an ad hoc way.

The lepton sector (massless neutrinos)

For a lepton sector with a lagrangian density of the form

$$\mathcal{L}^{(h2,\ell)} = -\overline{L} \phi \Lambda_e E_R - \overline{E}_R \Lambda_e^\dagger \phi^\dagger L, \quad (12.87)$$

in which

$$L = \begin{pmatrix} N_L \\ E_L \end{pmatrix}$$

is a weak doublet containing the three families of neutrinos (N_L) and charged leptons (E_L) and E_R is a three-family weak singlet, we find massless neutrinos. As before, one can write $\Lambda_e = V_e G_e W_e^\dagger$ and we find

$$\mathcal{L}^{(h2,\ell)} \implies -M_e (\overline{E}_L V_e W_e^\dagger E_R - \overline{E}_R W_e V_e^\dagger E_L), \quad (12.88)$$

with $M_e = G_e v / \sqrt{2}$ the diagonal mass matrix with masses m_e, m_μ and m_τ . The mass fields $E_R^{\text{mass}} = W_e^\dagger E_R$, $E_L^{\text{mass}} = V_e^\dagger E_L$. For the (massless) neutrino fields we just can redefine fields into $N_L^{\text{mass}} = V_e^\dagger N_L$, since the weak current is the only place where they show up. The W -current then becomes $\overline{E}_L \gamma^\mu N_L = \overline{E}_L^{\text{mass}} \gamma^\mu N_L^{\text{mass}}$, i.e. there is no family mixing for massless neutrinos.

The lepton sector (massive Dirac neutrinos)

In principle a massive Dirac neutrino could be accounted for by a lagrangian of the type

$$\mathcal{L}^{(h2,\ell)} = -\bar{L}\phi\Lambda_e E_R - \bar{E}_R\Lambda_e^\dagger\phi^\dagger L - \bar{L}\phi^c\Lambda_n N_R - \bar{N}_R\Lambda_n^\dagger\phi^{c\dagger} L \quad (12.89)$$

with three righthanded neutrinos added to the previous case, decoupling from all known interactions. Again we continue as before now with matrices $\Lambda_e = V_e G_e W_e^\dagger$ and $\Lambda_n = V_n G_n W_n^\dagger$, and obtain

$$\mathcal{L}^{(h2,\ell)} \implies -\bar{E}_L V_e M_e W_e^\dagger E_R - \bar{E}_R W_e M_e V_e^\dagger E_L - \bar{N}_L V_n M_n W_n^\dagger N_R - \bar{N}_R W_n M_n V_n^\dagger N_L. \quad (12.90)$$

We note that there are mass fields $E_R^{\text{mass}} = W_e^\dagger E_R$, $E_L^{\text{mass}} = V_e^\dagger E_L$, $N_L^{\text{mass}} = V_n^\dagger N_L$ and $N_R^{\text{mass}} = W_n^\dagger N_R$ and the weak current becomes $\bar{E}_L \gamma^\mu N_L = \bar{E}_L^{\text{mass}} \gamma^\mu V_e^\dagger V_n N_L^{\text{mass}}$. Working with the mass eigenstates for the charged leptons we see that the weak eigenstates for the neutrinos are $N_L^{\text{weak}} = V_e^\dagger N_L$ with the relation to the mass eigenstates for the lefthanded neutrinos given by

$$N_L' = N_L^{\text{weak}} = V_e^\dagger V_n N_L^{\text{mass}} = U_{\text{PMNS}}^\dagger N_L^{\text{mass}}, \quad (12.91)$$

with $U_{\text{PMNS}} = V_n^\dagger V_e$ known as the Pontecorvo-Maki-Nakagawa-Sakata mixing matrix.

For neutrino's this matrix is parametrized in terms of three angles θ_{ij} with $c_{ij} = \cos \theta_{ij}$ and $s_{ij} = \sin \theta_{ij}$ and one angle δ ,

$$U_{\text{PMNS}} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13} e^{i\delta} \\ 0 & 1 & 0 \\ -s_{13} e^{i\delta} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (12.92)$$

a parametrization that in principle also could have been used for quarks. In this case, it is particularly useful because θ_{12} is essentially determined by solar neutrino oscillations requiring $\Delta m_{12}^2 \approx 8 \times 10^{-5} \text{ eV}^2$ (convention $m_2 > m_1$), while θ_{23} then is determined by atmospheric neutrino oscillations requiring $|\Delta m_{23}^2| \approx 2.5 \times 10^{-3} \text{ eV}^2$. The mixing is intriguingly close to the Harrison-Perkins-Scott tri-bimaximal mixing matrix

$$U_{\text{HPS}} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \sqrt{1/2} & -\sqrt{1/2} \\ 0 & \sqrt{1/2} & \sqrt{1/2} \end{pmatrix} \begin{pmatrix} \sqrt{2/3} & \sqrt{1/3} & 0 \\ -\sqrt{1/3} & \sqrt{2/3} & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} \sqrt{2/3} & \sqrt{1/3} & 0 \\ -\sqrt{1/6} & \sqrt{1/3} & -\sqrt{1/2} \\ -\sqrt{1/6} & \sqrt{1/3} & \sqrt{1/2} \end{pmatrix}. \quad (12.93)$$

The lepton sector (massive Majorana fields)

An even simpler option than sterile righthanded Dirac neutrinos, is to add in Eq. 12.88 a Majorana mass term for the (lefthanded) neutrino mass eigenstates,

$$\mathcal{L}^{\text{mass},\nu} = -\frac{1}{2} (M_L \bar{N}_L^c N_L + M_L^* \bar{N}_L N_L^c), \quad (12.94)$$

although this option is not attractive as it violates the electroweak symmetry. The way to circumvent this is to introduce as in the previous section righthanded neutrinos, with for the righthanded sector a mass term M_R ,

$$\mathcal{L}^{\text{mass},\nu} = -\frac{1}{2} (M_R \bar{N}_R N_R^c + M_R^* \bar{N}_R^c N_R). \quad (12.95)$$

In order to have more than a completely decoupled sector, one must for the neutrinos as well as charged leptons, couple the right- and lefthanded species through Dirac mass terms coming from the coupling to the Higgs sector as in the previous section. Thus (disregarding family structure) one has two Majorana neutrinos, one being massive. For the charged leptons there cannot exist a Majorana mass term as this would break the U(1) electromagnetic symmetry. For the leptons, the left- and righthanded species then just form a Dirac fermion.

For the neutrino sector, the massless and massive Majorana neutrinos, coupled by a Dirac mass term, are equivalent to two decoupled Majorana neutrinos (see below). If the Majorana mass $M_R \gg M_D$ one actually obtains in a natural way one Majorana neutrino with a very small mass. This is called the see-saw mechanism (outlined below).

For these light Majorana neutrinos one has, as above, a unitary matrix relating them to the weak eigenstates. Absorption of phases in the states is not possible for Majorana neutrinos, however, hence the mixing matrix becomes

$$V_{\text{PMNS}} = U_{\text{PMNS}} K \quad \text{with} \quad K = \begin{pmatrix} e^{i\alpha_1/2} & 0 & 0 \\ 0 & e^{i\alpha_2/2} & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (12.96)$$

containing three (CP-violating) phases (α_1 , α_2 and δ).

The see-saw mechanism

Consider (for one family $N = n$) the most general Lorentz invariant mass term for two independent Majorana spinors, Υ'_1 and Υ'_2 (satisfying $\Upsilon^c = \Upsilon$ and as discussed in chapter 6, $\Upsilon_L^c \equiv (\Upsilon_L)^c = \Upsilon_R$ and $\Upsilon_R^c = \Upsilon_L$). We use here the primes starting with the weak eigenstates. Actually, it is easy to see that this incorporates the Dirac case by considering the lefthanded part of Υ'_1 and the righthanded part of Υ'_2 as a Dirac spinor ψ . Thus

$$\Upsilon'_1 = n_L^c + n_L, \quad \Upsilon'_2 = n_R + n_R^c, \quad \psi = n_R + n_L. \quad (12.97)$$

As the most general mass term in the lagrangian density we have

$$\begin{aligned} \mathcal{L}^{\text{mass}} &= -\frac{1}{2} (M_L \overline{n_L^c} n_L + M_L^* \overline{n_L} n_L^c) - \frac{1}{2} (M_R \overline{n_R} n_R^c + M_R^* \overline{n_R^c} n_R) \\ &\quad - \frac{1}{2} (M_D \overline{n_L^c} n_R^c + M_D^* \overline{n_L} n_R) - \frac{1}{2} (M_D \overline{n_R} n_L + M_D^* \overline{n_R^c} n_L^c) \end{aligned} \quad (12.98)$$

$$= -\frac{1}{2} \begin{pmatrix} \overline{n_L^c} & \overline{n_R} \end{pmatrix} \begin{pmatrix} M_L & M_D \\ M_D & M_R \end{pmatrix} \begin{pmatrix} n_L \\ n_R^c \end{pmatrix} + \text{h.c.} \quad (12.99)$$

which for $M_D = 0$ is a pure Majorana lagrangian and for $M_L = M_R = 0$ and real M_D represents the Dirac case. The mass matrix can be written as

$$M = \begin{pmatrix} M_L & |M_D| e^{i\phi} \\ |M_D| e^{i\phi} & M_R \end{pmatrix} \quad (12.100)$$

taking M_L and M_R real and non-negative. This choice is possible without loss of generality because the phases can be absorbed into Υ'_1 and Υ'_2 (real must be replaced by hermitean if one includes families). This is a mixing problem with a symmetric (complex) mass matrix leading to two (real) mass eigenstates. The diagonalization is analogous to what was done for the Λ -matrices and one finds $U M U^T = M_0$ with a (unitary) matrix U , which implies $U^* M^\dagger U^\dagger = U^* M^* U^\dagger = M_0$ and a 'normal' diagonalization of the (hermitean) matrix MM^\dagger ,

$$U (MM^\dagger) U^\dagger = M_0^2, \quad (12.101)$$

Thus one obtains from

$$MM^\dagger = \begin{pmatrix} M_L^2 + |M_D|^2 & |M_D| (M_L e^{-i\phi} + M_R e^{+i\phi}) \\ |M_D| (M_L e^{+i\phi} + M_R e^{-i\phi}) & M_R^2 + |M_D|^2 \end{pmatrix}, \quad (12.102)$$

the eigenvalues

$$M_{1/2}^2 = \frac{1}{2} \left[M_L^2 + M_R^2 + 2|M_D|^2 \pm \sqrt{(M_L^2 - M_R^2)^2 + 4|M_D|^2 (M_L^2 + M_R^2 + 2M_L M_R \cos(2\phi))} \right], \quad (12.103)$$

and we are left with two decoupled Majorana fields Υ_1 and Υ_2 , related via

$$\begin{pmatrix} \Upsilon_{1L} \\ \Upsilon_{2L} \end{pmatrix} = U^* \begin{pmatrix} n_L \\ n_R^c \end{pmatrix}, \quad \begin{pmatrix} \Upsilon_{1R} \\ \Upsilon_{2R} \end{pmatrix} = U \begin{pmatrix} n_L^c \\ n_R \end{pmatrix}. \quad (12.104)$$

for each of which one finds the lagrangians

$$\mathcal{L} = \frac{1}{4} \overline{\Upsilon}_i i \not{\partial} \Upsilon_i - \frac{1}{2} M_i \overline{\Upsilon}_i \Upsilon_i \quad (12.105)$$

for $i = 1, 2$ with real masses M_i . For the situation $M_L = 0$ and $M_R \gg M_D$ (taking M_D real) one finds $M_1 \approx M_D^2/M_R$ and $M_2 \approx M_R$.

Exercises

Exercise 12.1

Consider the case of the Weyl mode for symmetries. Prove that if the generators Q^a generate a symmetry, i.e. $[Q^a, H] = 0$, and $|a\rangle$ and $|a'\rangle$ belong to the same multiplet (there is a Q^a such that $|a'\rangle = Q^a|a\rangle$) then $H|a\rangle = E_a|a\rangle$ implies that $H|a'\rangle = E_a|a'\rangle$, i.e. a and a' are degenerate states.

Exercise 12.2

Derive for the vector and axial vector currents, $\vec{V}^\mu = \bar{\psi} \gamma^\mu \vec{T} \psi$ and $\vec{A}^\mu = \bar{\psi} \gamma^\mu \gamma_5 \vec{T} \psi$

$$\begin{aligned} \partial_\mu \vec{V}^\mu &= i \bar{\psi} [M, \vec{T}] \psi, \\ \partial_\mu \vec{A}^\mu &= i \bar{\psi} \{M, \vec{T}\} \gamma_5 \psi. \end{aligned}$$

Exercise 12.3

(a) The coupling of the Z^0 particle to fermions is described by the vertex

$$-i \frac{g}{2 \cos \theta_W} \left(C_V^f \gamma^\mu - C_A^f \gamma^\mu \gamma_5 \right),$$

with

$$\begin{aligned} C_V &= T_W^3 - 2Q \sin^2 \theta_W, \\ C_A &= T_W^3. \end{aligned}$$

Write down the matrix element squared (averaged over initial spins and summed over final spins) for the decay of the Z^0 . Neglect the masses of fermions and use the fact that the sum over polarizations is

$$\sum_{\lambda=1}^3 \epsilon_\mu^{(\lambda)}(p) \epsilon_\nu^{(\lambda)*}(p) = -g_{\mu\nu} + \frac{p_\mu p_\nu}{M^2}.$$

to calculate the width $\Gamma(Z^0 \rightarrow f\bar{f})$,

$$\Gamma(Z^0 \rightarrow f\bar{f}) = \frac{M_Z}{48\pi} \frac{g^2}{\cos^2 \theta_W} \left(C_V^{f2} + C_A^{f2} \right).$$

- (b) Calculate the width to electron-positron pair, $\Gamma(Z^0 \rightarrow e^+e^-)$, and the width to a pair of neutrino's, $\Gamma(Z^0 \rightarrow \nu_e\bar{\nu}_e)$. The mass of the Z^0 is $M_Z = 91$ GeV, the weak mixing angle is given by $\sin^2 \theta_W = 0.231$.

Exercise 12.4

Calculate the lifetime $\tau = 1/\Gamma$ for the top quark (t) assuming that the dominant decay mode is

$$t \rightarrow b + W^+.$$

In the standard model this coupling is described by the vertex

$$\frac{-ig}{2\sqrt{2}}(\gamma^\mu - \gamma^\mu\gamma_5).$$

The masses are $m_t \approx 175$ GeV, $m_b \approx 5$ GeV and $M_W \approx 80$ GeV.

Exercise 12.5

Show that the coupling to the Higgs (W^+W^-h , ZZh , hhh and e^+e^-h) are proportional to the mass squared (bosons) or mass (fermions) of the particles. Note that you can find the answer without explicit construction of the interaction terms in the lagrangian.

Exercise 12.6

Check that the use of the Wolfenstein parametrization in the CKM matrix indeed gives a unitary matrix, at least up to a high (which?) order in λ .

Exercise 12.7

In this exercise two limits are investigated for the two-Majorana case.

- (a) Calculate for the special choice $M_L = M_R = 0$ and M_D real, the mass eigenvalues and show that the mixing matrix is

$$U = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix}$$

which enables one to rewrite the Dirac field in terms of Majorana spinors. Give the explicit expressions that relate ψ and ψ^c with Υ_1 and Υ_2 .

- (b) A more interesting situation is $0 = M_L < |M_D| \ll M_R$, which leads to the so-called see-saw mechanism. Calculate the eigenvalues $M_L = 0$ and $M_R = M_X$. Given that neutrino masses are of the order of 0.05 eV, what is the mass M_X if we take for M_D the electroweak symmetry breaking scale v (about 250 GeV).